

# Gold-Catalyzed Nitrene Transfer to Activated Alkynes: Formation of $\alpha,\beta$ -Unsaturated Amidines


Chaoqun Li and Liming Zhang\*

Department of Chemistry and Biochemistry, University of California, Santa Barbara, California 93106, United States

zhang@chem.ucsb.edu

Received January 26, 2011

## ABSTRACT



A gold-catalyzed intermolecular nitrene transfer to alkynes was developed for the first time, revealing a new mode of nitrene transfer and providing a novel access to versatile  $\alpha$ -imino metal carbenes. Various mild nitrene-transfer reagents were examined, and iminopyridinium ylides especially those based on 3,5-dichloropyridine proved to be highly effective. With activated alkynes such as  $N$ -alkynyl oxazolidinones as substrates,  $\alpha,\beta$ -unsaturated amidines were formed in mostly good yields.

A rather unique and potent reactivity of gold catalysis is the capability of the metal to form gold carbenes via  $\gamma$ -eliminations. Although there are still some ambiguities regarding discrete reaction mechanisms,<sup>1</sup> this strategy has been applied successfully in intramolecular transformations using tethered oxygen-delivering oxidants such as

sulfoxides,<sup>2</sup>  $N$ -oxides,<sup>1b,3</sup> nitrones,<sup>4</sup> and epoxide<sup>5</sup> where their oxygenless parts (i.e., Y) behave as nucleofuges (Scheme 1A). Attempts to realize intermolecular versions of this strategy using external sulfoxides<sup>6</sup> led to 3,3-rearrangements instead, and the intermediacy of  $\alpha$ -oxo gold carbenes was ruled out both by experiments and via calculations. We discovered, however, that pyridine  $N$ -oxides and quinoline  $N$ -oxides were suitable external oxidants for the generation of these reactive gold intermediates intermolecularly, thus offering valuable opportunities to replace hazardous and potentially explosive diazo ketones with benign alkynes in gold carbene chemistry (Scheme 1B). We have applied this strategy successfully

(1) (a) Li, C.-W.; Pati, K.; Lin, G.-Y.; Sohel, S. M. A.; Hung, H.-H.; Liu, R.-S. *Angew. Chem., Int. Ed.* **2010**, 10.1002/anie.201004647. (b) Cui, L.; Zhang, G.; Peng, Y.; Zhang, L. *Org. Lett.* **2009**, 11, 1225–1228. (c) Cui, L.; Ye, L.; Zhang, L. *Chem. Commun.* **2010**, 46, 3351–3353.

(2) (a) Shapiro, N. D.; Toste, F. D. *J. Am. Chem. Soc.* **2007**, 129, 4160–4161. (b) Li, G.; Zhang, L. *Angew. Chem., Int. Ed.* **2007**, 46, 5156–5159. (c) Davies, P. W.; Albrecht, S. J. C. *Angew. Chem., Int. Ed.* **2009**, 48, 8372–8375. (d) Davies, P. W. *Pure Appl. Chem.* **2010**, 82, 1537–1544.

(3) Cui, L.; Peng, Y.; Zhang, L. *J. Am. Chem. Soc.* **2009**, 131, 8394–8395.


(4) (a) Yeom, H. S.; Lee, Y.; Lee, J. E.; Shin, S. *Org. Biomol. Chem.* **2009**, 7, 4744–4752. (b) Yeom, H. S.; Lee, Y.; Jeong, J.; So, E.; Hwang, S.; Lee, J. E.; Lee, S. S.; Shin, S. *Angew. Chem., Int. Ed.* **2010**, 49, 1611–1614. (c) Yeom, H. S.; Lee, J. E.; Shin, S. *Angew. Chem., Int. Ed.* **2008**, 47, 7040–7043.

(5) (a) Lin, G.-Y.; Li, C.-W.; Hung, S.-H.; Liu, R.-S. *Org. Lett.* **2008**, 10, 5059–5062. (b) Li, C.-W.; Lin, G.-Y.; Liu, R.-S. *Chem.—Eur. J.* **2010**, 16, 5803–5811. (c) Hashmi, A. S. K.; Buehrle, M.; Salathe, R.; Bats, J. W. *Adv. Synth. Catal.* **2008**, 350, 2059–2064.


(6) Cuenca, A. B.; Montserrai, S.; Hossain, K. M.; Mancha, G.; Lledos, A.; Medio-Simon, M.; Ujaque, G.; Asensio, G. *Org. Lett.* **2009**, 11, 4906–4909.

**Scheme 1.** Gold-Catalyzed Oxygen/Nitrene Transfer to Alkyne: Concept


**A. Intramolecular oxidation** (Y = sulfide, imine, *tert*-amine)



**B. Intermolecular oxidation** (Y = pyridine, quinoline)




**C. Intermolecular nitrene transfer – this work**



to rather expedient syntheses of several classes of functional molecules.<sup>7</sup>

With all the studies so far focused on oxidation using oxygen-delivering oxidants, there is surprisingly little attention paid to nitrene-delivering oxidants. While a tethered azido group<sup>8</sup> and a tethered sulfilimine (one example only)<sup>2a</sup> could be considered as intramolecular cases, no gold-catalyzed intermolecular nitrene transfer to alkynes has been reported (Scheme 1C). Several features of this mode of nitrene transfer are notable: (1) the nitrene moiety is delivered via an outer sphere attack, and no gold nitrene complex<sup>9</sup> is involved; this new mode of nitrene transfer is different from many known nitrene transfer reactions;<sup>10</sup> (2) it would make alkynes equivalent to  $\alpha$ -diazo imines, which are difficult to access. It is known that  $\alpha$ -diazo imines readily cyclize into 1,2,3-triazoles although in the presence of electron-withdrawing substituents at N-1 the reversed process could be utilized to generate  $\alpha$ -imino rhodium carbenes.<sup>11</sup> Herein, we disclose our effort in synthesizing and evaluating various external nitrene-transfer reagents in generating  $\alpha$ -imino gold carbenes directly from C–C triple bonds, which led to efficient synthesis of  $\alpha$ ,  $\beta$ -unsaturated amidines.

At the outset, a range of nitrene-transfer reagents were synthesized for screening. As shown in Figure 1, they



**Figure 1.** Selected nitrene-transfer reagents (NTR).

include sulfilimines **1** and **2** derived from dibenzothiophene,<sup>12</sup> cyclic sulfilimine **3**,<sup>13</sup> iminoiodinane **4**, which is soluble in organic solvents,<sup>14</sup> and iminopyridinium ylides **5–8**,<sup>15</sup> and iminoquinolinium ylide **9**.<sup>15</sup>

After having little success with aliphatic internal alkynes, we turned our attention to activated alkynes, specifically *N*-alkynyloxazolidinones,<sup>7c,16</sup> which are readily available from terminal alkynes via several convenient methods.<sup>17</sup> Table 1 shows some of the reaction optimization results. To our delight, expected amidine **11** was indeed formed when various nitrene-transfer reagents were used. While sulfilimines **1** and **2** were mostly ineffective, cyclic sulfilimine **3** gave good yields with a range of different gold catalysts including IPrAuNTf<sub>2</sub> (entry 5) and (F<sub>5</sub>C<sub>6</sub>)<sub>3</sub>PAuNTf<sub>2</sub> (entry 6). While the result with IPrAuNTf<sub>2</sub> was excellent, the fact that the propylsulfanyl group remained in the product prompted us to search further for oxidants that deliver a simple tosyl group. While soluble iminoiodinane **4** was less effective, iminopyridinium ylides offered mostly good to excellent yields (entries 8–13). Iminopyridinium ylide **8** derived from 3,5-dichloropyridine proved to be the most effective and led to the formation of **11** in excellent yields regardless of the electrophilic nature of the gold catalyst (comparing entries 11, 12, and 13). Notably, steric bulk on the oxidant was detrimental, and the 2-Me group in ylide **7** decreased the reaction yield from 65% to 25% (comparing entries 8 and 10). For the same reason, iminoquinolinium ylide **9** was a poor oxidant (entry 14).

(7) (a) Ye, L.; Cui, L.; Zhang, G.; Zhang, L. *J. Am. Chem. Soc.* **2010**, *132*, 3258–3259. (b) Ye, L.; He, W.; Zhang, L. *J. Am. Chem. Soc.* **2010**, *132*, 8550–8551. (c) Lu, B.; Li, C.; Zhang, L. *J. Am. Chem. Soc.* **2010**, *132*, 14070–14072.

(8) (a) Gorin, D. J.; Davis, N. R.; Toste, F. D. *J. Am. Chem. Soc.* **2005**, *127*, 11260–11261. (b) Hiroya, K.; Matsumoto, S.; Ashikawa, M.; Ogiwara, K.; Sakamoto, T. *Org. Lett.* **2006**, *8*, 5349–5352.

(9) (a) Li, Z.; Ding, X.; He, C. *J. Org. Chem.* **2006**, *71*, 5876–5880. (b) Li, Z.; Capretto, D. A.; Rahaman, R. O.; He, C. *J. Am. Chem. Soc.* **2007**, *129*, 12058–12059.

(10) (a) Fantauzzi, S.; Caselli, A.; Gallo, E. *Dalton Trans.* **2009**, 5434–5443. (b) Diaz-Requejo, M. M.; Perez, P. J. *Chem. Rev.* **2008**, *108*, 3379–3394. (c) Mueller, P.; Fruit, C. *Chem. Rev.* **2003**, *103*, 2905–2919.

(11) (a) Horneff, T.; Chuprakov, S.; Chernyak, N.; Gevorgyan, V.; Fokin, V. V. *J. Am. Chem. Soc.* **2008**, *130*, 14972–14974. (b) Chuprakov, S.; Kwok, S. W.; Zhang, L.; Lercher, L.; Fokin, V. V. *J. Am. Chem. Soc.* **2009**, *131*, 18034–18035. (c) Chuprakov, S.; Hwang, F. W.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2007**, *46*, 4757–4759.

(12) Desikan, V.; Liu, Y.; Toscano, J. P.; Jenks, W. S. *J. Org. Chem.* **2008**, *73*, 4398–4414.


(13) Chen, C.-S.; Tan, C.-M.; Huang, C.-H.; Chang, L.-C.; Wang, J.-P.; Cheng, F.-C.; Chern, J.-W. *Bioorg. Med. Chem.* **2010**, *18*, 597–604.

(14) Macikenas, D.; Skrzypczak-Jankun, E.; Protasiewicz, J. D. *J. Am. Chem. Soc.* **1999**, *121*, 7164–7165.

(15) Gangapuram, M.; Redda, K. K. *J. Heterocycl. Chem.* **2009**, *46*, 309–316.

(16) (a) Davies, P. W.; Cremonesi, A.; Martin, N. *Chem. Commun.* **2011**, 47, 379–381. (b) Couty, S.; Meyer, C.; Cossy, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 6726–6730. (c) Evano, G.; Coste, A.; Jouvin, K. *Angew. Chem., Int. Ed.* **2010**, *49*, 2840–2859. (d) De, K. K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, *110*, 5064–5106.

(17) (a) Kohnen, A. L.; Dunetz, J. R.; Danheiser, R. L. *Org. Synth.* **2007**, *84*, 88–101. (b) Sagamanova, I. K.; Kurtz, K. C. M.; Hsung, R. P. *Org. Synth.* **2007**, *84*, 359–367. (c) Coste, A.; Couty, F.; Evano, G. *Org. Synth.* **2010**, *87*, 231–244.

**Table 1.** Reaction Optimization

entry <sup>a</sup>	NTR	gold catalyst	reaction conditions	conversion (%)	yield (%)	E/Z
1	<b>1</b>	Ph <sub>3</sub> PAuNTf <sub>2</sub>	rt, 12 h	80	10	2/1
2	<b>1</b>	IPrAuNTf <sub>2</sub>	rt, 12 h	20	<5	-
3	<b>1</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 12 h	100	35	1.5/1
4	<b>2</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 12 h	100	30	1.5/1
5	<b>3</b>	IPrAuNTf <sub>2</sub>	rt, 3 h	100	90	3.2/1
6	<b>3</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 12 h	90	70	1.5/1
7	<b>4</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 3 h	100	46	3/1
8	<b>5</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 18 h	80	65	2.5/1
9	<b>6</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 48 h	70	60	1.5/1
10	<b>7</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 48 h	50	25	1.5/1
11	<b>8</b>	(F <sub>5</sub> C <sub>6</sub> ) <sub>3</sub> PAuNTf <sub>2</sub>	rt, 12 h	100	90	1.5/1
12	<b>8</b>	IPrAuNTf <sub>2</sub>	rt, 3 h	100	95 <sup>b</sup>	4/1
13	<b>8</b>	<b>12<sup>c</sup></b>	rt, 3 h	100	95	4/1
14	<b>9</b>	IPrAuNTf <sub>2</sub>	rt, 48 h	30	10	3/1



<sup>a</sup>[**10**]=0.05 M. <sup>b</sup>92% isolated yield. <sup>c</sup>**12**=dichloro(2-picolinato)gold(III).

The amidine product **11** was formed as an inseparable mixture of two isomers. <sup>1</sup>H NMR indicated that they were isomeric at the C–C double bond<sup>2d</sup> and the *E*-isomer was major but only to a small extent. Interestingly, this lack of stereoselectivity was in stark contrast to our previous result with the same substrate, which was oxidized into an  $\alpha,\beta$ -unsaturated imide with a >50/1 *E/Z* ratio.<sup>7c</sup> The geometries of the C–N double bonds in both isomers were assigned as *E* and discussed later.

Our attempts to improve the *E/Z* selectivities of the C–C double bond were mostly unsuccessful. A notable feature of the optimized reaction conditions is the absence of an acid additive, which was required in our previous studies using pyridine *N*-oxides.<sup>7a,b</sup>

The reaction scope was then studied, and the results are shown in Table 2. Good to excellent yields were obtained with substrates substituted at the alkyne terminus by a primary alkyl group (entry 1) and several functionalized ones (entries 2–6). Functional groups such as chloro (entry 2), aryl (entry 3), ester (entry 4), MsO (entry 5), and ether (entry 6) were readily tolerated; however, proximally positioned HO and azido groups appeared to interfere with the desired reaction, and little or no desired product was observed. The allowance of MsO is noteworthy as it offers a reactive electrophilic site for further

(18) The product was formed likely via the following sequence:


**Table 2.** Scope Studies<sup>a</sup>

1	2	3
<b>14a</b> , 93% E/Z = 4/1	<b>14b</b> , 81% <sup>b</sup> E/Z = 2/1	<b>14c</b> , 88% E/Z = 3/1
4	5	6
<b>14d</b> , 79% <sup>b</sup> E/Z = 2/1	<b>14e</b> , 72% E/Z = 3/2	<b>14f</b> , 75% E/Z = 5/1
7	8	9 <sup>d</sup>
<b>14g</b> , 90%	<b>14h</b> , 76% <sup>c</sup>	<b>14i</b> , 85% E/Z = 5/1
10 <sup>d</sup>	11 <sup>d</sup>	12
<b>14j</b> , 88% E/Z = 4/1	<b>14k</b> , 79% E/Z = 5/1	<b>14l</b> , 62% <sup>e</sup> E/Z = 3/2

<sup>a</sup>[**13**] = 0.05 M. <sup>b</sup>(F<sub>5</sub>C<sub>6</sub>)<sub>3</sub>PAuNTf<sub>2</sub> was the catalyst. <sup>c</sup>15% of the ring-expansion product was formed. <sup>d</sup>The oxidant used was analogous to **8** with the only difference at the sulfonyl group. <sup>e</sup>Dichloro(2-picolinato)gold(III).

transformation while highlighting the mild nature of this nitrene transfer process. A cyclohexyl group was easily accommodated, leading to **14g** as the only isomer in 90% yield (entry 7). Similarly, an efficient reaction was observed with a cyclopentyl group (entry 8); however, besides the expected 1,2-C–H insertion, a ring expansion occurred as a minor event, leading to a cyclohexene product<sup>18</sup> in 15% yield. In addition to tosyl nitrene, other arenesulfonyl nitrenes were readily transferred to alkynes using analogous iminopyridium ylides regardless of the electronic properties of the arene (comparing entries 9 and 10) as well as its steric bulk (entry 11, Mts = 2,4,6-trimethylbenzenesulfonyl). While the oxazolidinone moiety appeared to be ideal for this nitrene transfer reaction, an *N*-phenyltoluenesulfonyl moiety, as shown in entry 12, could also play the role of activating the C–C triple bond toward this intermolecular nitrene transfer.

The geometry of the C–N double bond was difficult to assign even after rather extensive NOESY studies. Eventually we were able to secure a single crystal of the major component of compound **14j** (Figure 2), and its X-ray diffraction studies revealed that the C–N double bond has




**Figure 2.** ORTEP drawing of *E*-**14j**.

an *E*-configuration. Although the corresponding geometry of the minor isomer was not determined, we reasoned it should be the same as that of the major isomer as it is difficult to imagine that the geometries of both C–N and C–C double bonds would be paired in an exclusive manner (i.e., *E,E* and *Z,Z* only). The exclusive formation of the *E*-C–N double bond was also consistent with the fact that **14g** and **14h** did not have any observed geometric isomer. The reason for the selectivity, although not apparent, could be rationalized by conformational analysis. As shown in Scheme 2, assuming that gold carbene intermediate **B** is formed via a concerted elimination, stereoelectronic effects require bond alignments as outlined in conformation **A**. The preference for such a conformation can be reasoned by invoking the minimization of charge repulsion between the heteroatoms on the oxazolidinone ring and the sulfonyl oxygens. Accordingly, the geometries of the C–N double bonds in other cases were all tentatively assigned as *E*.

In conclusion, we have developed the first examples of gold-catalyzed intermolecular nitrene transfer to alkynes

**Scheme 2.** Rationale for the Formation of the *E*-C–N Double Bond



and realized a new nitrene transfer strategy that provides a novel and facile access to versatile  $\alpha$ -imino metal carbenes. Various mild nitrene-transfer reagents were examined, and iminopyridium ylides especially those based on 3,5-dichloropyridine turned out to be highly effective. With activated alkynes such as *N*-alkynyloxazolidinones as substrates,  $\alpha$ -imino gold carbene intermediates were readily generated and underwent facile 1,2-C–H insertions, leading to  $\alpha,\beta$ -unsaturated amidines in mostly good yields. Further studies to expand the scope to normal alkynes will be pursued in due course.

**Acknowledgment.** The authors thank the NIH and UCSB for general financial support and Dr. Guang Wu for his kind help with X-ray crystallography. L.Z. is an Alfred P. Sloan fellow.

**Note Added after ASAP Publication.** In the second line under Figure 1, “cyclic sulfilimine 2” was changed to “cyclic sulfilimine 3”; the corrected version reposted March 1, 2011.

**Supporting Information Available.** Experimental procedures, compound characterization and spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.