

# Selective Formation of 1,5-Substituted Sulfonyl Triazoles Using Acetylides and Sulfonyl Azides


Maria Elena Meza-Aviña, Mudita Kishor Patel, Cylivia B. Lee, Thomas J. Dietz, and Mitchell P. Croatt\*

Department of Chemistry and Biochemistry, University of North Carolina at Greensboro, Greensboro, North Carolina 27402, United States

mpcroatt@uncg.edu

Received March 15, 2011

## ABSTRACT



The reaction of acetylides with sulfonyl azides was found to selectively form 1,5-substituted sulfonyl triazoles. This reaction thus provides access to the regioisomeric product as compared to the popular copper-catalyzed azide–alkyne cycloaddition. The reaction is efficient and selective with a variety of alkyne sources and sulfonyl azides and can incorporate an additional electrophile to yield 1,4,5-trisubstituted sulfonyl triazoles.

The copper-catalyzed reaction of terminal alkynes with azides to yield 1,2,3-triazoles has deservedly received much attention in recent years.<sup>1</sup> This reaction regioselectively forms 1,4-substituted triazoles using alkyl, aryl, or sulfonyl azides.<sup>2</sup> Although it was reported that 1,4-substituted sulfonyl triazoles could be converted to 2,4-substituted

triazoles using amine bases,<sup>3</sup> there are no efficient methods to selectively convert terminal alkynes and sulfonyl azides to 1,5-substituted triazoles.<sup>4–6</sup> In 2004, there was an excellent report for the reaction of alkynyl Grignard reagents with carbon-substituted azides to yield 1,5-disubstituted triazoles and 1,4,5-trisubstituted triazoles.<sup>7</sup> Interestingly, no reactions were reported with sulfonyl azides despite the authors'

(1) For a recent set of reviews in this area, see the themed issue: *Chem. Soc. Rev.* **2010**, *39*, 1221–1408.

(2) (a) Yoo, E. J.; Ahlquist, M.; Kim, S. H.; Bae, I.; Fokin, V. V.; Sharpless, K. B.; Chang, S. *Angew. Chem., Int. Ed.* **2007**, *46*, 1730. (b) Wang, F.; Fu, H.; Jiang, Y.; Zhao, Y. *Adv. Synth. Catal.* **2008**, *350*, 1830.

(3) Yamauchi, M.; Miura, T.; Murakami, M. *Heterocycles* **2010**, *80*, 177.

(4) For a report of the conversion of aryl or alkyl azides to 1,5-triazoles, see: Boren, B. C.; Narayan, S.; Rasmussen, L. K.; Zhang, L.; Zhao, H.; Lin, Z.; Jia, G.; Fokin, V. V. *J. Am. Chem. Soc.* **2008**, *130*, 8923.

(5) For earlier reports of reactions of acetylides with sulfonyl azides with low efficiency or no yield reported, see: (a) Boyer, J. H.; Mack, C. H.; Goebel, N.; Morgan, L. R., Jr. *J. Org. Chem.* **1958**, *23*, 1051. (b) Robson, E.; Tedder, J. M.; Webster, B. *J. Chem. Soc.* **1963**, 1863. (c) Huisgen, R.; Knorr, R.; Möbius, L.; Szeimies, G. *Chem. Ber.* **1965**, *98*, 4014. (d) Helwig, R.; Hanack, M. *Chem. Ber.* **1985**, *118*, 1008.

(6) For the synthesis of 1,5-substituted sulfonyl triazoles from sulfonyl azides and phosphorus ylides, see: Harvey, G. R. *J. Org. Chem.* **1966**, *31*, 1587.


(7) (a) Krasinski, A.; Fokin, V. V.; Sharpless, K. B. *Org. Lett.* **2004**, *6*, 1237. For seminal work, see: (b) Akimova, G. S.; Chistokletov, V. N.; Petrov, A. A. *Zh. Org. Khim.* **1967**, *3*, 968. (c) Akimova, G. S.; Chistokletov, V. N.; Petrov, A. A. *Zh. Org. Khim.* **1967**, *3*, 2241. (d) Akimova, G. S.; Chistokletov, V. N.; Petrov, A. A. *Zh. Org. Khim.* **1968**, *4*, 389.

(8) For representative examples, see: (a) Raushel, J.; Fokin, V. V. *Org. Lett.* **2010**, *12*, 4952. (b) Chuprakov, S.; Kwok, S. W.; Zhang, L.; Lercher, L.; Fokin, V. V. *J. Am. Chem. Soc.* **2009**, *131*, 18034. (c) Yoo, E. J.; Ahlquist, M.; Bae, I.; Sharpless, K. B.; Fokin, V. V.; Chang, S. *J. Org. Chem.* **2008**, *73*, 5520. (d) Yoo, E. J.; Ahlquist, M.; Kim, S. H.; Bae, I.; Fokin, V. V.; Sharpless, K. B.; Chang, S. *Angew. Chem., Int. Ed.* **2007**, *46*, 1730.

expertise with this electrophilic azide source.<sup>8</sup> Additionally, the copper-catalyzed reaction with sulfonyl azides often forms products other than the typical triazoles.<sup>8</sup> Due to the significance of substituted triazoles in ligand design, drug discovery, and biochemical applications,<sup>1</sup> it is attractive to have a facile, regioselective, and efficient route to 1,5-substituted sulfonyl triazoles. In this report we describe an efficient method for the selective synthesis of 1,5-substituted sulfonyl triazoles from terminal alkynes and sulfonyl azides.

During our initial study of the reactivity of acetylides and sulfonyl azides, it was determined that the major product of the reaction with 1-heptyne was 1,5-triazole **3** (Scheme 1). This was true for both sulfonyl azides **2a** and **2b**. The structures were determined by comparison to the 1,4-triazoles formed by reaction with CuI and 2,6-lutidine along with X-ray crystallographic analysis of **3a** (see Supporting Information).

**Scheme 1.** Synthesis of 1,5-Substituted Triazoles



Although this reaction appears similar to the frequently employed copper-catalyzed azide–alkyne cycloaddition (CuAAC) and complementary ruthenium-catalyzed cycloaddition (RuAAC),<sup>4</sup> there are two main differences. First, the reaction described herein produces 1,5-substituted triazoles whereas the CuAAC yields 1,4-substituted triazoles. Second, this reaction works efficiently with sulfonyl azides, whereas the RuAAC has not been reported with sulfonyl azides. Based on these differences, it was decided to optimize this reaction and study its scope.

Optimization of the reaction conditions determined that if the reaction was not kept at  $-78^\circ\text{C}$ , mixtures of differentially substituted triazoles (mostly the 1,4-substituted triazoles) were isolated (Table 1, entry 1 versus 2; see the

**Table 1.** Optimization for Synthesis of 1,5-Triazoles **3**<sup>a</sup>

| entry          | heptyne (equiv)/<br>ArSO <sub>2</sub> N <sub>3</sub> (equiv) | base<br>(equiv)      | time<br>(h) | yield<br>of <b>3</b> <sup>b</sup> |
|----------------|--|----------------------|-------------|-----------------------------------|
| 1              | <b>1</b> (1.0)/ <b>2a</b> (1)                                | <i>n</i> -BuLi (1.2) | 2           | 64% <b>3a</b>                     |
| 2 <sup>c</sup> | <b>1</b> (1.0)/ <b>2a</b> (1)                                | <i>n</i> -BuLi (1.2) | 5           | 39% <b>3a</b>                     |
| 3              | <b>1</b> (1.0)/ <b>2a</b> (1)                                | NaHMDS (1.2)         | 2           | 23% <b>3a</b>                     |
| 4              | <b>1</b> (1.0)/ <b>2a</b> (1)                                | KHMDS (1.2)          | 2           | 27% <b>3a</b>                     |
| 5              | <b>1</b> (1.2)/ <b>2a</b> (1)                                | <i>n</i> -BuLi (1.2) | 0.3         | 90% <b>3a</b>                     |
| 6              | <b>1</b> (1.2)/ <b>2b</b> (1)                                | <i>n</i> -BuLi (1.2) | 0.3         | 67% <b>3b</b>                     |
| 7 <sup>d</sup> | <b>1</b> (1.2)/ <b>2c</b> (1)                                | <i>n</i> -BuLi (1.2) | 2           | 79% <b>3c</b>                     |

<sup>a</sup> Reaction conditions (unless otherwise noted): The indicated base was added to a solution of heptyne in THF at  $-78^\circ\text{C}$ . After 15 min, the indicated sulfonyl azide (1 equiv) was added, and the reaction was stirred for the time indicated. <sup>b</sup> Isolated yield. <sup>c</sup> The reaction temperature was increased to ambient temperature after 2 h. <sup>d</sup> **2c**, Ar = *p*-NO<sub>2</sub>-Ph.

Supporting Information for a more detailed optimization table). Interestingly, it was found that other counterions, including sodium and potassium, allowed for the formation of 1,5-triazoles; however the reactions were not as efficient, potentially due to the poor solubility of the anions at  $-78^\circ\text{C}$  (entry 1 versus 3 and 4). A very poor yield was observed with a Grignard reagent (not shown) which likely illustrates why the reactions described herein were not described in prior reports.<sup>7</sup> A final key reaction condition for the optimization was to use a slight excess of the alkyne, relative to the sulfonyl

**Table 2.** Alkyne Screen<sup>a</sup>


| entry          | substrate         | ArSO <sub>2</sub> N <sub>3</sub> | time   | yield <sup>b</sup>   | product |
|----------------|-------------------|----------------------------------|--------|----------------------|---------|
| 1              | Ph—C≡C—           | <b>2a</b>                        | 40 min | 79%<br>( <b>4a</b> ) |         |
| 2              | Ph—C≡C—           | <b>2b</b>                        | 25 min | 85%<br>( <b>4b</b> ) |         |
| 3              | Ph—C≡C—           | <b>2c</b>                        | 25 min | 87%<br>( <b>4c</b> ) |         |
| 4              |                   | <b>2a</b>                        | 45 min | 80%<br>( <b>5</b> )  |         |
| 5              |                   | <b>2a</b>                        | 25 min | 24%<br>( <b>6</b> )  |         |
| 6              |                   | <b>2a</b>                        | 25 min | 53%<br>( <b>7</b> )  |         |
| 7 <sup>c</sup> |                   | <b>2a</b>                        | 3 hr   | 0%<br>( <b>8</b> )   |         |
| 8              | TMS—C≡C—          | <b>2a</b>                        | 25 min | 14%<br>( <b>9</b> )  |         |
| 9              | <i>t</i> -Bu—C≡C— | <b>2a</b>                        | 45 min | 87%<br>( <b>10</b> ) |         |

<sup>a</sup> Reaction conditions (unless otherwise noted): 1.2 equiv of *n*-BuLi was added to a solution of the indicated alkyne (1.2 equiv) in THF at  $-78^\circ\text{C}$ . After 15 min, 1 equiv of the indicated sulfonyl azide was added, and the reaction was stirred for the time indicated at which point the sulfonyl azide was fully consumed as judged by TLC. <sup>b</sup> Isolated yield. <sup>c</sup> Quantitative recovery of starting alkyne and sulfonyl azide as judged by <sup>1</sup>H NMR of the crude reaction mixture. Ts = *p*-toluenesulfonyl; Tris = 2,4,6-triisopropylphenylsulfonyl; *p*-Ns = *p*-nitrophenylsulfonyl.

azide source (entry 5). With these optimal conditions, other sulfonyl azide sources (**2a** and **2c**) were examined and the reaction was found to maintain efficiency (entries 6 and 7).

With conditions in hand for the efficient transformation of heptyne into 1,5-triazoles, a variety of different alkynes were studied in the reaction (Table 2). Importantly, the reaction with phenylacetylene was also efficient with all three of the sulfonyl azide sources (entries 1–3). It was determined that electron-rich arenes worked better in this reaction than electron-poor arenes (entry 4 versus 6), although the more sterically hindered ortho-anisole substrate was not as efficient (entry 5). Although a nitro-group was tolerated on the sulfonyl azide, the nitro-substituted aryl acetylide did not yield any desired product and only starting material was isolated (entry 7). The silyl-substituted acetylide was low yielding for this reaction (entry 8), whereas the *tert*-butyl acetylide was highly efficient (entry 9).

**Scheme 2.** Proposed Mechanism for the Reactions



Based on the proposed reaction mechanism (Scheme 2), it was decided to explore the reactions of the triazole anion (C) with other electrophiles (formation of E). Using identical conditions as described earlier, the reactions were quenched with electrophiles other than a Brønsted acid (Table 3). Acid chlorides (entries 1–3) were able to yield the respective ketones with moderate efficiency. Unfortunately, the reactions with benzylchloroformate or a chlorosilane were not high yielding (entries 4 and 5), but the reaction with the iodomethane proceeded smoothly (entry 6). Unexpectedly, when the *p*-nitrophenylsulfonyl azide source (**2c**) was used in slight excess, the free triazole was isolated where the presumed anion (Scheme 2, C) reacted similarly to an intermolecular electrophilic aromatic substitution and subsequently cleaved the sulfur–nitrogen bond (Table 3, entry 7). These reactions formally represent highly regioselective cycloadditions between sulfonyl azides and internal alkynes, a reaction that is otherwise difficult.


It should be noted that the 1,5-substituted sulfonyl triazoles isomerized to the 1,4-triazoles to a moderate amount when being purified by silica gel chromatography, presumably via an acid-catalyzed pathway. To circumvent this, crystallization was utilized to isolate the pure 1,5-triazoles. Surprisingly, when the crystals of some of the 1,5-triazole products were stored in a  $-20\text{ }^{\circ}\text{C}$  freezer for extended periods (> 1 week), isomerization to the 1,4-triazoles still occurred. When the same 1,5-triazoles

**Table 3.** Electrophile Screen<sup>a</sup>

| entry          | Alkyne | ArSO <sub>2</sub> N <sub>3</sub> | electrophile           | yield <sup>b</sup> | product |
|----------------|--------|----------------------------------|------------------------|--------------------|---------|
| 1              | 1      | 2a                               | BzCl                   | 80% (11a)          |         |
| 2              | 1      | 2b                               | BzCl                   | 57% (11b)          |         |
| 3              | 1      | 2a                               | Cl-C(=O)- <i>n</i> -Pr | 66% (12)           |         |
| 4              | 1      | 2a                               | Cl-C(=O)-OBn           | 19% (13)           |         |
| 5              | 1      | 2a                               | TBSCl                  | 17% (14)           |         |
| 6 <sup>c</sup> | 1      | 2a                               | MeI                    | 64% (15)           |         |
| 7              | 1      | 2c                               | 2c                     | 52% (16)           |         |

<sup>a</sup> Reaction conditions (unless otherwise noted): 1.2 equiv of *n*-BuLi was added to a solution of the indicated alkyne (1.2 equiv) in THF at  $-78\text{ }^{\circ}\text{C}$ . After 15 min, 1 equiv of *p*-toluenesulfonyl azide was added, the reaction was stirred for 30 min, and then 2 equiv of the indicated electrophile were added. The reactions were stirred at  $-78\text{ }^{\circ}\text{C}$ . <sup>b</sup> Isolated yield. <sup>c</sup> After addition of MeI the reaction was warmed to ambient temperature. Ts = *p*-toluenesulfonyl; Tris = 2,4,6-triisopropylphenylsulfonyl.

were stored as solutions in CDCl<sub>3</sub> no observable isomerization occurred (see Supporting Information for more details). These data lend evidence for the isomerization



**Figure 1.** Adjacent triazoles in crystal structure of **3a**. Carbon atoms are colored gray, nitrogen atoms are blue, oxygen atoms are red, and sulfur atoms are yellow.

taking place even in the crystalline form. The crystal structure of **3a** further supports this hypothesis since the nitrogen atom in the 3-position is in relatively close proximity to the sulfur atom of an adjacent molecule (Figure 1). Other isomerization mechanisms are also possible, including radical reactions, but these were not explored since the isomerization did not occur to an appreciable amount in solution.

Importantly, the reactions described herein combine two readily available starting materials, terminal alkynes and sulfonyl azides, to efficiently and selectively assemble 1,5-substituted sulfonyl triazoles. This reaction displays the substituents of the triazole much closer together than the complementary 1,4-triazoles and thus provides for new structural opportunities. Additionally, the putative triazole anion was trapped with various electrophiles to

regioselectively form 1,4,5-trisubstituted triazoles. Further explorations into this reaction, including the isomerization of 1,5-triazoles to 1,4-triazoles, are ongoing.

**Acknowledgment.** Funding for this project from The University of North Carolina at Greensboro is gratefully acknowledged. The authors thank Dr. Franklin J. Moy (UNCG), Dr. Brandie Ehrmann (UNCG), and Dr. Cynthia S. Day (Wake Forest University) for assisting with the analysis of NMR, mass spectrometry, and X-ray crystallography data, respectively.

**Supporting Information Available.** Experimental data and spectroscopic characterization of all products and crystallographic data for **3a**. This material is available free of charge via the Internet at <http://pubs.acs.org>.