

SmI₂-Mediated Cross-Coupling of Nitrones with β -Silyl Acrylates: Synthesis of (+)-Australine


Pierre Gilles and Sandrine Py*

Département de Chimie Moléculaire (SERCO) UMR-5250, ICMG FR-2607, CNRS, Université Joseph Fourier, BP 53, 38041 Grenoble Cedex 09, France

sandrine.py@ujf-grenoble.fr

Received December 20, 2011

ABSTRACT



The SmI₂-mediated cross-coupling of nitrones with β -silyl- α,β -unsaturated esters, followed by zinc reduction, allows an efficient and highly diastereoselective preparation of β -silyl lactams, which are precursors of β -hydroxy lactams through Tamao–Fleming oxidation. By applying the method to a cyclic, carbohydrate-derived nitronone, a new synthesis of (+)-australine has been realized in only 11 steps and in 21% overall yield from L-xylose.

Since its introduction in 1977, samarium diiodide has become a useful reagent for the reduction of a wide range of organic functions and for stereoselective formation of carbon–carbon bonds.¹ A few years ago, the SmI₂-mediated cross-coupling of nitrones with α,β -unsaturated esters was reported² and has since proved useful for elaboration of the bicyclic skeleton of pyrrolizidine iminosugars, such as (+)-hyacinthacine A₂.³ Recently, we have become interested in designing syntheses of other naturally occurring pyrrolizidines, especially those bearing hydroxyl groups in position 6 and/or 7 (Figure 1).

The introduction of a hydroxyl group in position 7 of pyrrolizidines was first attempted through reductive cross-coupling of nitronone **1** with α,β -unsaturated esters bearing an oxygen atom in the β position. However, treatment of




Figure 1. Natural polyhydroxylated pyrrolizidines.

nitronone **1** and ethyl (*E*)-3-trimethylsilyloxypropenoate⁴ (**2**) with SmI₂ in the presence of water^{3a} did not provide the expected cross-coupled product, but *N*-hydroxy-pyrrolizidine **3**, resulting from the reduction of the starting nitronone (Scheme 1).

Although alkoxy-substituted captodative olefins have been used as Michael acceptors for the synthesis of β -hydroxy- γ -lactones through SmI₂-mediated cross-coupling with ketones⁵ and aldehydes,⁶ the yields are often


(1) (a) Namy, J. L.; Girard, P.; Kagan, H. B. *Nouv. J. Chim.* **1977**, *1*, 5. (b) Girard, P.; Namy, J. L.; Kagan, H. B. *J. Am. Chem. Soc.* **1980**, *102*, 2693. (c) Procter, D. J.; Flowers, R. A., II; Skrydstrup, T., Eds.; *Organic Synthesis Using Samarium Diiodide: A Practical Guide*; Royal Society of Chemistry Publishing: 2010.

(2) (a) Masson, G.; Cividino, P.; Py, S.; Vallée, Y. *Angew. Chem., Int. Ed.* **2003**, *42*, 2265. (b) Riber, D.; Skrydstrup, T. *Org. Lett.* **2003**, *5*, 229. (c) Cividino, P.; Py, S.; Delair, P.; Greene, A. E. *J. Org. Chem.* **2007**, *72*, 485.

(3) (a) Desvergnès, S.; Py, S.; Vallée, Y. *J. Org. Chem.* **2005**, *70*, 1459. (b) Desvergnès, S.; Desvergnès, V.; Martin, O. R.; Itoh, K.; Liu, H.-w.; Py, S. *Bioorg. Med. Chem.* **2007**, *15*, 6443. (c) Reháč, J.; Fišera, L.; Kožíšek, J.; Bellovičová, L. *Tetrahedron* **2011**, *67*, 5762.

(4) Foehlich, B.; Giering, W. *Synthesis* **1980**, 231.

Scheme 1. Attempted Cross-Coupling of Nitronne **1** and Acrylate **2**




moderate in the case of *intermolecular* reactions. Probably due to electronic effects, β -alkoxy-esters are poor acceptors for nitronne conjugate addition reactions. In contrast, the SmI₂-mediated cross-coupling of nitrones with alkyl β -substituted α,β -unsaturated esters, such as methyl crotonate, afforded the expected coupling product in good yields and with excellent stereoselectivities.^{2a,c} Carbon and silicon, being in the same column of the periodic table, present similar electronic properties.⁷ We thus thought that the SmI₂-mediated cross-coupling of nitrones with β -silyl-substituted α,β -unsaturated esters could be more successful than with the alkoxy esters. The silyl group would then serve as a masked hydroxyl group, to be later revealed by Tamao–Fleming oxidation.⁸ β -Silyl acrylates or acrylamides have already been used in conjugate addition reactions⁹ but, to the best of our knowledge, never in a SmI₂-mediated cross-coupling reaction.¹⁰ Here, we report the SmI₂-mediated cross-coupling of nitrones with a β -silyl acrylate and its application to the synthesis of (+)-australine.

In a preliminary assay, nitronne **4**¹¹ and ethyl (*E*)-3-(dimethylphenylsilyl)propenoate (**5**)¹² in the presence of water (8 equiv) at -78 °C were treated with SmI₂. Through coordination to SmI₂, water forms a more reactive species, often allowing faster reactions and better

yields.¹³ Under these conditions, the desired *N*-hydroxylamines **6a,b** were obtained, but proved to be relatively unstable in air. Fortunately, a direct, single-pot reduction of the reaction product using Zn/AcOH¹⁴ afforded the lactams **7a,b** in 51% yield and in a 90/10 *cis/trans*¹⁵ ratio (Table 1, entry 1), accompanied by the β -silylpropanoate **8**,¹⁶ arising from reduction of **5** (35%).

Table 1. SmI₂-Mediated Cross-Coupling of Nitronne **4** and Acrylate **5**



entry	SmI ₂ (equiv)	additives (equiv)	temp (°C)	time (h)	7a,b (yield %)	<i>cis/trans</i> ^a
1	3	H ₂ O (8)	-78	28	51	90/10
2	4	H ₂ O (8)	-78 to -30	0.1	53	90/10
3	4	H ₂ O/LiBr (8/12)	-78 to -30	0.1	84	>98/2

^a Ratio determined by ¹H analysis of the crude reaction mixture.

Next, the same reagents were mixed at -78 °C and the cooling bath was removed at the end of SmI₂ addition. At a temperature of -30 °C (reached after 5 min), the starting nitronne was completely consumed. However, after zinc reduction of the intermediate *N*-hydroxylamines **6a,b**, lactams **7a,b** (90/10 *cis/trans* ratio) were isolated in a similar 53% yield (Table 1, entry 2). The introduction into the reaction mixture of LiBr¹⁷ proved to be more rewarding. Indeed, with a combination of SmI₂ and H₂O/LiBr additives, only the *cis* lactam **7a** was formed and in an improved yield of 84% (Table 1, entry 3). Flowers has demonstrated that the addition of 12 equiv of LiBr to solutions of SmI₂ in THF provides a system with an oxidation potential of -1.98 V as compared with -1.33 V for SmI₂ alone vs the Ag/AgNO₃ reference electrode.¹⁸ The increase in the reducing ability of the system may well account for the improvement in the reaction efficiency. Furthermore, introduction of the lithium cation seems to favor the formation of the *cis* lactam.

(5) Kerrigan, N. J.; Upadhyay, T.; Procter, D. J. *Tetrahedron Lett.* **2004**, *45*, 9087.

(6) Donadel, O. J.; Martín, T.; Padrón, J. M. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 18.

(7) As a consequence, silicon has been considered as an isostere of carbon in drug discovery: (a) Showell, G. A.; Mills, J. S. *Drug Discovery Today* **2003**, *8*, 551. (b) Bains, W.; Tacke, R. *Curr. Opin. Drug Discovery Dev.* **2003**, *6*, 526. (c) Gately, S.; West, R. *Drug Dev. Res.* **2007**, *68*, 156.

(8) (a) Tamao, K.; Ishida, N.; Tanaka, T.; Kumada, M. *Organometallics* **1983**, *2*, 1694. (b) Fleming, I.; Henning, R.; Plaut, H. *J. Chem. Soc., Chem. Commun.* **1984**, 29.

(9) (a) Klumpp, G. W.; Mierop, A. J. C.; Vrieling, J. J.; Brugman, A.; Schakel, M. *J. Am. Chem. Soc.* **1985**, *107*, 6740. (b) Fleming, I.; Lawrence, N. J. *J. Chem. Soc., Perkin Trans. 1* **1998**, 2679. (c) Clayden, J.; Watson, D. W.; Helliwell, M.; Chambers, M. *Chem. Commun.* **2003**, 2582. (d) Stork, G.; Tang, P. C.; Casey, M.; Goodman, B.; Toyota, M. *J. Am. Chem. Soc.* **2005**, *127*, 16255. (e) Shintani, R.; Okamoto, K.; Hayashi, T. *Org. Lett.* **2005**, *7*, 4757. (f) Lipshutz, B. H.; Tanaka, N.; Taft, B. R.; Lee, C.-T. *Org. Lett.* **2006**, *8*, 1963. (g) Balskus, E. P.; Jacobsen, E. N. *J. Am. Chem. Soc.* **2006**, *128*, 6810.

(10) For the SmI₂-mediated hydrodimerization of silyl acrylates, see: (a) Fleming, I.; Ghosh, S. K. *J. Chem. Soc., Chem. Commun.* **1992**, 1775. (b) Fleming, I.; Ghosh, S. K. *J. Chem. Soc., Perkin Trans. 1* **1998**, 2711.

(11) Dondoni, A.; Franco, S.; Junquera, F.; Merchan, F.; Merino, P.; Tejero, T. *Synth. Commun.* **1994**, *24*, 2537.

(12) (a) Takeshita, K.; Seki, Y.; Kawamoto, K.; Murai, S.; Sonoda, N. *J. Org. Chem.* **1987**, *52*, 4864. (b) Sieburth, S. M.; Lang, J. *J. Org. Chem.* **1999**, *64*, 1780.

(13) Szostak, M.; Spain, M.; Parmar, D.; Procter, D. J. *Chem. Commun.* **2012**, *8*, 330.

(14) See Supporting Information.

(15) Relative configuration was determined by NOESY experiments. See Supporting Information.


(16) (a) Sommer, L. H.; Marans, N. S. *J. Am. Chem. Soc.* **1950**, *72*, 1935. (b) Nakashima, T.; Fujiyama, R.; Fujio, M.; Tsuno, Y. *Bull. Chem. Soc. Jpn.* **1999**, *72*, 741.

(17) Peltier, H. M.; McMahon, J. P.; Patterson, A. W.; Ellman, J. A. *J. Am. Chem. Soc.* **2006**, *128*, 16018.

(18) Fuchs, J. R.; Mitchell, M. L.; Shabangi, M.; Flowers, R. A., II. *Tetrahedron Lett.* **1997**, *38*, 8157.

The cross-coupling of nitron **4** was also considered with ethyl (*Z*)-3-(dimethylphenylsilyl)propenoate (**9**).¹⁹ Surprisingly, at $-78\text{ }^{\circ}\text{C}$ and in the presence of water, no trace of *N*-hydroxyamino esters **6a,b** was detected (Scheme 2). The starting nitron was mostly recovered (74%), while acrylate **9** was reduced to propanoate **8** (28%) and dimerized adipic acid derivatives **10a,b** (31%). These results suggest that acrylate **9** is reduced by SmI_2 faster than nitron **4** (and than its *E*-isomer, **5**). This reduction competing with the formation of a (ketyl-type) radical anion from **4**²⁰ would thus hamper conjugate addition onto the unsaturated ester.²¹

Scheme 2. Attempted Reaction of Nitron **4** with *Z*-Acrylate **9**



Having in hand suitable reaction conditions for the coupling of nitron **4** with **5**, we next turned to studying the reaction of this β -silyl acrylate with nitron **1**,^{3a} which exhibits an appropriate substitution pattern for the synthesis of the different alkaloids represented in Figure 1. When nitron **1** was treated with acrylate **5** under the same conditions (Table 1, entry 3), lactam **11** was isolated in 49% yield as a single diastereomer (Table 2, entry 1). The intermediate *N*-hydroxypyrrolidine **12** could also be isolated in 49% yield and as a single diastereomer when the zinc reduction step was omitted.

With the aim of improving the efficiency of the reductive coupling step, various alternative proton sources were investigated, with a particular focus on the effect of their acidity. Indeed, the rate of SmI_2 -mediated reduction of acetophenone has been shown to be proportional to the acidity of the proton source used as an additive.²² The groups of Procter²³ and Nicolaou²⁴ have also demonstrated the role of the acidity of additives on the efficiency of C–C bond formation induced by SmI_2 . In the presence of phenol ($\text{p}K_{\text{a}}$ 9.9) or 2,4,6-trichlorophenol ($\text{p}K_{\text{a}}$ 6.2), the yields of the expected lactam **11** were lower (or similar) than with water (Table 2, entries 2 and 3). On the other hand, use of the noncoordinating hexafluoroisopropanol (HFIP, $\text{p}K_{\text{a}}$ 9.3) as the additive afforded improved efficiency in

(19) Cardona, F.; Parmeggiani, C.; Faggi, E.; Bonaccini, C.; Gratterer, P.; Sim, L.; Gloster, T. M.; Roberts, S.; Davies, G. J.; Rose, D. R.; Goti, A. *Chem.—Eur. J.* **2009**, *15*, 1627.

(20) Masson, G.; Py, S.; Vallée, Y. *Angew. Chem., Int. Ed.* **2002**, *41*, 1772.


(21) For previous work showing easier reductive dimerization of *Z*- β -silyl acrylates in comparison to their *E*-isomers see, ref 10.

(22) Chopade, P. R.; Prasad, E.; Flowers, R. A. *J. Am. Chem. Soc.* **2004**, *126*, 44.

(23) Edmonds, D. J.; Muir, K. W.; Procter, D. J. *J. Org. Chem.* **2003**, *68*, 3190.

(24) Nicolaou, K. C.; Li, A.; Edmonds, D. J.; Tria, G. S.; Ellery, S. P. *J. Am. Chem. Soc.* **2009**, *131*, 16905.

Table 2. Direct Preparation of Lactam **11**



entry	additives (equiv)	temp ($^{\circ}\text{C}$)	time (h)	11 (%)	dr ^a
1	$\text{H}_2\text{O}/\text{LiBr}$ (8/12)	-78 to -30	0.1	49	>98/2
2	PhOH/LiBr (8/12)	-78 to -30	0.1	32	>98/2
3	2,4,6-triCl-PhOH/LiBr (8/12)	-78 to -30	0.1	50	>98/2
4	HFIP/LiBr (8/12)	-78 to -30	0.1	64	>98/2

^a Only one diastereomer was detected by ^1H NMR analysis of the crude reaction mixture.

the coupling, with **11** being isolated in 64% yield (Table 2, entry 4). It is not clear how to rationalize the effect of HFIP, its $\text{p}K_{\text{a}}$ being intermediate between those of the two phenols tested. Although the role of proton sources in such reactions has been extensively investigated, it is still not fully understood and acidity is certainly not the only parameter to consider.²⁵


The configuration of the new stereocenters in **11** was determined to be *S* at both C-7 and C-7a from NOESY correlations.¹⁴ The approach of the β -silyl acrylate by the *Re* face of nitron **1** is directed by the substituents of the nitron (at C-1 and C-3 in the product).^{3a} Remarkably, the configuration of the silylated stereocenter is also completely controlled in this SmI_2 -mediated cross-coupling reaction. Such control may be explained by a favored chelated transition state **A**, similar to those proposed in Heathcock's model for conjugate addition of *E*-enolates on β -substituted acrylic esters (Scheme 3).²⁶

Pyrrolizidinone **11** is thus a possible precursor of (+)-australine, a polyhydroxylated pyrrolizidine isolated from

(25) (a) Hutton, T. K.; Muir, K. W.; Procter, D. J. *Org. Lett.* **2003**, *5*, 4811. (b) Amiel-Levy, M.; Hoz, S. *J. Am. Chem. Soc.* **2009**, *131*, 8280. (c) Sadasivam, D. V.; Teprovich, J. A.; Procter, D. J.; Flowers, I. I. R. A. *Org. Lett.* **2010**, *12*, 4140.

(26) Oare, D. A.; Heathcock, C. H. Stereochemistry of the Base-Promoted Michael Addition Reaction. In *Topics in Stereochemistry*; Eliel, E. L., Wilen, S. H., Eds.; Wiley: New York, 1989; Vol. 19, pp 227–408. For a recent discussion and computational investigation on intermolecular Michael reactions, see: Kwan, E. E.; Evans, D. A. *Org. Lett.* **2010**, *12*, 5124. The authors thank one of the referees for bringing this paper to their attention.

Scheme 3. Possible Transition State for the SmI₂-Promoted Reductive Coupling of Nitron 1 and Acrylate 5




the seeds of *Castanospermum australe*.²⁷ This natural product has attracted the interest of chemists because it not only alters amyloglucosidase activity²⁸ but also possesses a structure exhibiting five contiguous stereogenic centers, which represents a synthetic challenge.²⁹

When the Tamao–Fleming oxidation of pyrrolizidinone 11 was first carried out under acidic (HBF₄•OEt₂ in DCM, then H₂O₂, KF in DMF) or electrophilic (Br₂ and AcO₂H) conditions, decomposition of the starting material was observed. The protocol reported by Woerpel for silyl group oxidation under basic conditions was proved more successful:³⁰ pyrrolizidinone 11 was treated with potassium hydride, TBAF, and cumene hydroperoxide. Pyrrolizidinone 13, obtained in moderate yield (57%), was reduced by a BH₃•DMS complex to give pyrrolizidine 14 (Scheme 4).³¹ The use of cumene hydroperoxide, instead of *tert*-butyl hydroperoxide, reportedly avoids desilylation.³² Finally, it was found that reduction of pyrrolizidinone 11 to pyrrolizidine 15 prior to Tamao–Fleming oxidation gave a better overall yield (65% vs 50% for the first sequence). Deprotection of the hydroxyl groups was then accomplished by hydrogenolysis to afford quantitatively (+)-australine. The spectroscopic data and optical rotation of our synthetic sample of (+)-australine were in accordance with those previously reported for the

natural substance ($[\alpha]_D^{20} = 17.1$ (*c* 2.9, MeOH); lit.: $[\alpha]_D^{20} = 18.6$ (*c* 2.5, MeOH)).^{29b}

Scheme 4. Synthesis of (+)-Australine



In conclusion, we have shown that the β -silyl acrylate 5 is an excellent partner for the SmI₂-mediated *intermolecular* cross-coupling of nitrones, one that is capable of yielding β -silyl γ -amino acid derivatives or β -silyl γ -lactams with nearly complete diastereoselection. Through the use of this new cross-coupling, (+)-australine could be obtained in 11 steps and with an overall yield of 21% from L-xylose. This methodology should be useful for the synthesis of other polyhydroxylated pyrrolizidines.

Acknowledgment. P.G. is grateful to the Ministère de l'Enseignement Supérieur et de la Recherche for a doctoral fellowship.

Supporting Information Available. Characterization data, full experimental procedures, and copies of NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

(30) Smitrovich, J. H.; Woerpel, K. A. *J. Org. Chem.* **1996**, *61*, 6044.

(31) Pearson, W. H.; Hines, J. V. *J. Org. Chem.* **2000**, *65*, 5785.

(32) Peng, Z.-H.; Woerpel, K. A. *Org. Lett.* **2002**, *4*, 2945.

The authors declare no competing financial interest.

(27) Molyneux, R. J.; Benson, M.; Wong, R. Y.; Tropea, J. E.; Elbein, A. D. *J. Nat. Prod.* **1988**, *51*, 1198.

(28) Tropea, J. E.; Molyneux, R. J.; Kaushal, G. P.; Pan, Y. T.; Mitchell, M.; Elbein, A. D. *Biochemistry* **1989**, *28*, 2027.

(29) Selected syntheses of (+)-australine: (a) White, J. D.; Hrnčiar, P.; Yokochi, A. F. T. *J. Am. Chem. Soc.* **1998**, *120*, 7359. (b) Denmark, S. E.; Martinborough, E. A. *J. Am. Chem. Soc.* **1999**, *121*, 3046. (c) Romero, A.; Wong, C.-H. *J. Org. Chem.* **2000**, *65*, 8264. (d) Lauritsen, A.; Madsen, R. *Org. Biomol. Chem.* **2006**, *4*, 2898. (e) Ribes, C.; Falomir, E.; Carda, M.; Marco, J. A. *Org. Lett.* **2007**, *9*, 77. (f) Trost, B. M.; Aponick, A.; Stanzl, B. N. *Chem.—Eur. J.* **2007**, *13*, 9547. (g) Ritthiwigrom, T.; Willis, A. C.; Pyne, S. G. *J. Org. Chem.* **2010**, *75*, 815.