

Mimicking Nature: Synthetic Nicotinamide Cofactors for C=C Bio-reduction Using Enoate Reductases


Caroline E. Paul,^{†,§} Serena Gargiulo,^{‡,§} Diederik J. Opperman,^{||} Iván Lavandera,[†]
Vicente Gotor-Fernández,[†] Vicente Gotor,[†] Andreas Taglieber,[⊥]
Isabel W. C. E. Arends,[‡] and Frank Hollmann^{*,‡}

Department of Biotechnology, Delft University of Technology, Julianalaan 136,
2628BL Delft, The Netherlands, Department of Biotechnology, University of the Free State,
Bloemfontein 9300, South Africa, Department of Organic and Inorganic Chemistry,
University of Oviedo, Calle Julián Clavería 8, 33006 Oviedo, Spain, and Firmenich SA,
Corporate R&D, Route des Jeunes 1, CH-1211 Geneva, Switzerland

f.hollmann@tudelft.nl

Received November 25, 2012

ABSTRACT



A series of synthetic nicotinamide cofactors were synthesized to replace natural nicotinamide cofactors and promote enoate reductase (ER) catalyzed reactions without compromising the activity or stereoselectivity of the bioreduction process. Conversions and enantioselectivities of >99% were obtained for C=C bioreductions, and the process was successfully upscaled. Furthermore, high chemoselectivity was observed when employing these nicotinamide cofactor mimics (mNADs) with crude extracts in ER-catalyzed reactions.

The asymmetric reduction of conjugated C=C double bonds using enoate reductases (ERs, EC 1.3.1.31) is receiving great interest in preparative organic chemistry.¹ The rapidly expanding scope of ER-catalyzed stereospecific reductions makes this method a viable alternative to

transition-metal-catalyzed reductions. New ERs are constantly added to the toolbox from natural sources^{1d,2} and obtained *via* protein engineering.³ These enzymes are en route to becoming truly practical catalysts, although one

[‡] Delft University of Technology.

^{||} University of the Free State.

[†] University of Oviedo.

[⊥] Firmenich SA.

[§] These authors contributed equally.

(1) (a) *Enzyme Catalysis in Organic Synthesis*, 3rd ed.; Drauz, K., Gröger, H., May, O., Eds.; Wiley-VCH: Weinheim, 2012. (b) Hollmann, F.; Arends, I. W. C. E.; Holtmann, D. *Green Chem.* **2011**, *13*, 2285–2313. (c) Faber, K. *Biotransformations in Organic Chemistry*, 6th ed.; Springer: Berlin, 2011. (d) Toogood, H. S.; Gardiner, J. M.; Scrutton, N. S. *ChemCatChem* **2010**, *2*, 892–914. (e) Stürmer, R.; Hauer, B.; Hall, M.; Faber, K. *Curr. Opin. Chem. Biol.* **2007**, *11*, 203–213.

(2) For recent articles, see: (a) Liu, Y.-J.; Pei, X.-Q.; Lin, H.; Gai, P.; Liu, Y.-C.; Wu, Z.-L. *Appl. Microbiol. Biotechnol.* **2012**, *95*, 635–645. (b) Iqbal, N.; Rudroff, F.; Brige, A.; Van Beeumen, J.; Mihovilovic, M. D. *Tetrahedron* **2012**, *68*, 7619–7623. (c) Richter, N.; Gröger, H.; Hummel, W. *Appl. Microbiol. Biotechnol.* **2011**, *89*, 79–89. (d) Oberdorfer, G.; Steinkellner, G.; Stueckler, C.; Faber, K.; Gruber, K. *ChemCatChem* **2011**, 1562–1566.

(3) (a) Reetz, M. T. *Angew. Chem., Int. Ed.* **2011**, *50*, 138–174. (b) Toogood, H. S.; Fryszkowska, A.; Hulley, M.; Sakuma, M.; Mansell, D.; Stephens, G. M.; Gardiner, J. M.; Scrutton, N. S. *ChemBioChem* **2011**, *12*, 738–749. (c) Hulley, M. E.; Toogood, H. S.; Fryszkowska, A.; Mansell, D.; Stephens, G. M.; Gardiner, J. M.; Scrutton, N. S. *ChemBioChem* **2010**, *11*, 2433–2447. (d) Bougioukou, D. J.; Kille, S.; Taglieber, A.; Reetz, M. T. *Adv. Synth. Catal.* **2009**, *351*, 3287–3305.


remaining challenge involves their dependency on reduced nicotinamide cofactors [NAD(P)H] providing the reducing equivalents needed for the alkene bioreduction. NAD-(P)H is rather expensive,⁴ forbidding its stoichiometric use on a large scale. In principle, this can be overcome by using an (enzymatic) cofactor regeneration system.⁵

Another challenge resulting from the NAD(P)H-dependency of ERs arises with conjugated aldehydes and ketones as starting materials. Here, frequently unsatisfactory chemoselectivity is observed unless highly purified, alcohol dehydrogenase (ADH)-free, enzyme preparations are used. The reason thereof lies in the overlap of the substrate scope for both enzyme classes. As a result, both substrates and products of the ER-catalyzed transformation can also be converted by ‘contaminating’ ADHs leading to complex product mixtures, impairing the overall chemoselectivity of the reactions.^{1b,6}

Substitution of NAD(P)H as a reducing agent by other reductants appears to be straightforward with ERs and may be the method of choice to circumvent the above-mentioned challenges. Indeed, some promising approaches for NAD(P)H-independent regeneration have been reported recently.^{5a,7}

We became interested in synthetic, functional mimics of the natural nicotinamide cofactors (mNAD, Scheme 1) as stoichiometric reductants to promote ER-catalyzed reduction reactions.


Scheme 1. Asymmetric Bioreduction of Conjugated C=C Double Bonds Using Synthetic Nicotinamide Mimics (1-5)a and 6



These mNADs are simple and cheap to synthesize, starting from commercially available pyridine derivatives

(1,3-5)c; thus, the nitrogen was alkylated with benzyl or *n*-butyl bromide under reflux to obtain the bromide salts (1-5)b in high yields (81–92%), and the pyridinium ring was reduced into the corresponding dihydropyridine (1-5)a in moderate to high yields (35–81%) with sodium dithionite and sodium bicarbonate (Scheme 2).

Scheme 2. Straightforward Two-Step Synthesis of the Reduced Nicotinamide Mimics mNADs (1-5)a



Synthetic mNADs have received considerable attention as cost-efficient alternatives to the natural NAD(P)H cofactors.⁸ Unfortunately, the catalytic efficiencies of the wild-type alcohol dehydrogenases with mNADs generally fall back by orders of magnitude below their activity with the natural cofactors.⁹ In that respect, ERs represent an exception as they exhibit significant ‘cofactor promiscuity’.^{7a-c} In a first set of experiments, we evaluated the scope of enzymes accepting the mNAD **1a** as a replacement for NAD(P)H. As a model reaction, we chose the reduction of ketoisophorone (**7a**) to the corresponding levodione product (**7b**) to assess the conversion as well as the enantioselectivity of the ER-catalyzed reaction (Table 1). We were pleased to find that **1a** could replace the natural cofactors with a range of ERs without impairing the final yield or enantioselectivity of the reaction (entries 1–9). It is worth mentioning here that in the absence of either cofactor or enzyme no conversion was detectable within the time frame of the experiments.

For further investigations the enoate reductase homologue from *Thermus scotoductus* (TsER) was used.¹¹ As shown in Table 2, a broad range of different enones (entries 1–8), enals (entries 9–12), and maleimides (entries 13–16) could be converted in excellent yield and enantiospecificity demonstrating the preparative broadness of the ‘mimic approach’. Currently, we are bringing the proposed bioreduction scheme to a preparative scale. A first gram-scale reduction of *N*-phenyl-2-methyl maleimide (500 mM) gave excellent conversion and optical purities of the product (>99%) and acceptable isolated yields

(4) For more detailed information, see the Supporting Information.

(5) (a) Hollmann, F.; Arends, I. W. C. E.; Bühler, K. *ChemCatChem* **2010**, *2*, 762–782. (b) Weckbecker, A.; Gröger, H.; Hummel, W. *Biosystems Engineering I: Creating Superior Biocatalysts*, Vol. 120; Springer-Verlag: Berlin, 2010; pp 195–242. (c) Wichmann, R.; Vasic-Racki, D. *Technology Transfer in Biotechnology: from Lab to Industry to Production*, Vol. 92; Springer-Verlag: Berlin, 2005; pp 225–260. (d) Zhao, H.; van der Donk, W. A. *Curr. Opin. Biotechnol.* **2003**, *14*, 583–589.

(6) (a) Gargiulo, S.; Hanefeld, U.; Opperman, D. J.; Arends, I. W. C. E.; Hollmann, F. *Chem. Commun.* **2012**, *48*, 6630–6632. (b) Hall, M.; Stückler, C.; Ehammer, H.; Pointner, E.; Oberdorfer, G.; Gruber, K.; Hauer, B.; Stürmer, R.; Kroutil, W.; Macheroux, P.; Faber, K. *Adv. Synth. Catal.* **2008**, *350*, 411–418. (c) Hall, M.; Stückler, C.; Kroutil, W.; Macheroux, P.; Faber, K. *Angew. Chem., Int. Ed.* **2007**, *46*, 3934–3937. (d) Hall, M.; Hauer, B.; Stürmer, R.; Kroutil, W.; Faber, K. *Tetrahedron: Asymmetry* **2006**, *17*, 3058–3062.

(7) (a) Bernard, J.; van Heerden, E.; Arends, I. W. C. E.; Opperman, D. J.; Hollmann, F. *ChemCatChem* **2012**, *4*, 196–199. (b) Stückler, C.; Reiter, T. C.; Baudendistel, N.; Faber, K. *Tetrahedron* **2010**, *66*, 663–667. (c) Mifsud Grau, M.; van der Toorn, J. C.; Otten, L. G.; Macheroux, P.; Taglieber, A.; Zilly, F. E.; Arends, I. W. C. E.; Hollmann, F. *Adv. Synth. Catal.* **2009**, *351*, 3279–3286. (d) Taglieber, A.; Schulz, F.; Hollmann, F.; Rusek, M.; Reetz, M. T. *ChemBioChem* **2008**, *9*, 565–572.

(8) (a) Du, W. M.; Yu, Z. K. *Synlett* **2012**, 1300–1304. (b) Campbell, E.; Meredith, M.; Minter, S. D.; Banta, S. *Chem. Commun.* **2012**, *48*, 1898–1900. (c) Zhang, Y. H. P. *ACS Catal.* **2011**, *1*, 998–1009. (d) Ryan, J. D.; Fish, R. H.; Clark, D. S. *ChemBioChem* **2008**, *9*, 2579–2582. (e) Lutz, J.; Hollmann, F.; Ho, T. V.; Schnyder, A.; Fish, R. H.; Schmid, A. *J. Organomet. Chem.* **2004**, *689*, 4783–4790. (f) Lo, H. C.; Fish, R. H. *Angew. Chem., Int. Ed.* **2002**, *41*, 478–481. (g) Ansell, R. J.; Lowe, C. R. *Appl. Microbiol. Biotechnol.* **1999**, *51*, 703–710. (h) Ansell, R. J.; Small, D. A. P.; Lowe, C. R. *J. Mol. Catal. B: Enzym.* **1999**, *6*, 111–123. (i) Mauzerall, D.; Westheimer, F. H. *J. Am. Chem. Soc.* **1955**, *77*, 2261–2264.

(9) This adds up in a biochemical sense, as the high specificity of enzymes for either phosphorylated or nonphosphorylated nicotinamide cofactors is essential for controlling cellular metabolism.

Table 1. Performance of **1a** as a Replacement for NADH or NADPH in the Asymmetric Bioreduction of Ketoisophorone (**7a**) with Different ERs^a

entry	ER ^b	cofactor	conversion (%) ^c	ee (%) ^c
1	YqjM	NADH	92	84 (R)
2		NADPH	90	87 (R)
3		1a	96	85 (R)
4	TsER	NADH	90	>95 (R)
5		NADPH	>99	>95 (R)
6		1a	>99	>95 (R)
7	RmER	NADH	76	95 (R)
8		NADPH	73	95 (R)
9		1a	72	96 (R)

^a Conditions: [substrate]₀ = [cofactor]₀ = 10 mM, [ER] = 90–200 µg/mL, T = 30 °C, reaction time: 4 h. ^b YqjM: ER from *Bacillus subtilis*,¹⁰ TsER: ER homologue from *Thermus scotoductus*,¹¹ RmER: ER from *Ralstonia metallidurans* CH34. ^c Determined by GC analysis.

(> 70%), showing the synthetic usefulness of this C=C reduction process to obtain enantioenriched compounds on a large scale.

In order to obtain a more detailed insight into the reduction performance, the mNADs (**1–5a**) and **6** were evaluated and compared to the natural cofactors (NADH and NADPH).

Figure 1 shows the time courses of these bioreduction reactions. With the exception of the nitrile analogue (**5a**) and the Hantzsch ester (**6**), all synthetic nicotinamides exhibited equal or better activity with TsER. For example, initial rates obtained with **2a** were 1.5-fold higher than those with NADPH and almost double those with NADH. It is worth mentioning that the enantioselectivity in all cases was exclusive. The observed high activity of mimics (**1–4a**), even exceeding the activity of the natural cofactors, was somewhat unexpected, as with other enzyme classes investigated so far a significant decrease in activity had been observed. Molecular docking simulations of TsER with the mimics (Figure 2) confirmed the existence of productive binding modes for mimics **1a** to **5a**, leading to the reduced catalytically active FMNH₂ species.¹² Interestingly, for the Hantzsch ester (**6**) only unproductive binding was observed, supporting the experimental observations. Possibly, the increased activity with mimics can be explained by an increased FMN-reduction rate.

(10) (a) Kitzing, K.; Fitzpatrick, T. B.; Wilken, C.; Sawa, J.; Bourenkov, G. P.; Macheroux, P.; Clausen, T. *J. Biol. Chem.* **2005**, *280*, 27904–27913. (b) Fitzpatrick, T. B.; Amrhein, N.; Macheroux, P. *J. Biol. Chem.* **2003**, *278*, 19891–19897.

(11) (a) Opperman, D. J.; Sewell, B. T.; Litthauer, D.; Isupov, M. N.; Littlechild, J. A.; van Heerden, E. *Biochem. Biophys. Res. Commun.* **2010**, *393*, 426–431. (b) Opperman, D. J.; Piater, L. A.; van Heerden, E. *J. Bacteriol.* **2008**, *190*, 3076–3082.

(12) Although at this stage we cannot entirely exclude a mechanism wherein FMN is reduced in solution and then binds to the apo-ER.

Table 2. Substrate Scope of **1a**-Driven Chemoselective Reductions of α,β-Unsaturated Carbonyl Compounds (**7–14a**)^a

entry	product	cofactor	conversion (%) ^b	ee (%) ^b
1		NADH	>99	>95 (R)
2		1a	>99	>95 (R)
3		NADH	93	>99 (S)
4		1a	82	>99 (S)
5		NADH	>99	>99 (R)
6		1a	>99	>99 (R)
7		NADH	>99	>99 (R)
8		1a	>99	>99 (R)
9		NADH	>99	>99 (R)
10		1a	>99	>99 (R)
11		NADH	21	n.d. ^c
12		1a	18	n.d. ^c
13		NADH	>99	n.a. ^d
14		1a	>99	n.a. ^d
15		NADH	>99	>99 (R)
16		1a	>99	>99 (R)

^a Conditions: [substrate]₀ = [cofactor]₀ = 10 mM, [TsER] = 90 µg/mL, T = 30 °C, reaction time: 4 h. ^b Determined by GC analysis. ^c n.d. = not determined. ^d n.a. = not applicable.




Figure 1. Comparison of enzymatic conversions for the reduction of ketoisophorone (**7a**) to (6R)-levodione (**7b**) catalyzed by TsER using different cofactors. Conditions: [ketoisophorone]₀ = [cofactor]₀ = 10 mM, [TsER] = 100 µg/mL = 2.2 µM in MOPS buffer (50 mM, pH 7 with 5 mM CaCl₂, containing 2% v/v MeOH in the case of (**1–5a**) and **6**. **1a** (◇), **2a** (●), **3a** (Δ), **4a** (○), **5a** (□), **6** (×), NADH (▲), NADPH (■).

Finally, we investigated the presumed chemoselectivity advantage of using NADH mimics over natural cofactors. For this, the product distribution in the reduction of citral (**15**) was compared using crude preparations of YqjM recombinantly expressed in *Escherichia coli*. As shown in




Figure 2. Exemplary result of docking **1a** into the active site of TsER.

Table 3, using **1a** as a reductant, the desired C=C double bond reduction product (citronellal, **16**) was obtained as the sole product, while with NADH significant amounts of carbonyl reduction products (geraniol **17** and citronellol **18**) were formed.

Overall, we have demonstrated that cheap synthetic analogues of the natural nicotinamide cofactors (mNADs) represent a true alternative to the established regeneration systems to promote ER-catalyzed reduction reactions and apply them to the preparative scale. Already under non-optimized conditions, equal or higher conversions have been obtained compared to those with the natural cofactors, without altering the selectivity.

Another very interesting opportunity with mNADs lies in the bioorthogonality¹³ of these reduction schemes. The very poor activity of most enzyme classes tested so far with mNADs (particularly alcohol dehydrogenases and monooxygenases) enables the use of poorly purified (and hence cheap) ER preparations without impairing the selectivity of the desired reactions. Admittedly, issues such as the *in situ* regeneration of catalytic amounts of mNADs

(13) Ji, D.; Wang, L.; Hou, S.; Liu, W.; Wang, J.; Wang, Q.; Zhao, Z. K. *J. Am. Chem. Soc.* **2011**, *133*, 20857–20862.

Table 3. Chemoselectivity of YqjM-Crude Extract-Catalyzed Reduction of Citral (**15**) Using NADH and **1a** as Stoichiometric Reductant

entry	cofactor	16 (%) ^a	17 (%) ^a	18 (%) ^a	selectivity (%) ^b
1	NADH	21	20	20	34
2	1a	18	<1	<1	>90

^aConversion determined by GC analysis. ^bC=C double bond reduction over C=O bond reduction.

and/or recycling will have to be addressed in order to fully exploit their catalytic potential. Also, more mechanistic and kinetic studies as well as further modeling will be necessary to understand the catalytic mechanism. These studies are currently underway in our laboratories, eventually leading to a truly practical approach for white biotechnology.

Acknowledgment. Financial support from the Marie Curie ITN ‘BIOTRAINS’ (Grant Agreement No. 238531) is gratefully acknowledged. We would like to thank R. H. Fish (University of California Berkeley) for helpful discussions and Maarten Gorseling and Remco van Oosten (both Delft University of Technology) for technical assistance. I.L. acknowledges MICINN for his research contract under the Ramón y Cajal Program.

Supporting Information Available. Experimental procedures, full characterization and spectroscopic data for compounds (**1–5**)**a–b**, molecular docking and analytical methods. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.