

Synthesis of 3-Alkyl- or 3-Alkenyl-2-amidobenzofurans via Electrophilic Cyclization of *o*-Anisole-Substituted Ynamides with Carbocations


Yulong Kong,[†] Kezhi Jiang,[†] Jian Cao,* Lingzi Fu, Lian Yu, Guoqiao Lai, Yuming Cui, Ziqiang Hu, and Guanhai Wang

Key Laboratory of Organosilicon Chemistry and Material Technology of Ministry of Education, Hangzhou Normal University, Wenyi Road 222, Hangzhou 310012, People's Republic of China

caojianhznu@yahoo.cn

Received December 19, 2012

ABSTRACT



A facile carbocation-induced electrophilic cyclization reaction has been developed for the synthesis of 3-alkyl- or 3-alkenyl-2-amidobenzofurans from *o*-anisole-substituted ynamides and diarylmethanol or 1,1-diarylprop-2-yn-1-ol.

Since new and more efficient methods for accessing ynamides were established,¹ ynamides have emerged as important synthons in modern organic synthesis.² A variety of important functional groups and cyclic systems such

as enamides,³ amidines,⁴ indoles,⁵ and pyridines⁶ were successfully synthesized from ynamides. Benzofurans are the fundamental motif found in various natural products and the key structural units for pharmaceutical compounds.⁷ As a special class of functionalized benzofurans, 2-amidobenzofurans are of considerable interest^{8–11} and many methods have been established for their synthesis, such as

[†] These authors contributed equally.

(1) (a) Frederick, M. O.; Mulder, J. A.; Tracey, M. R.; Hsung, R. P.; Huang, J.; Kurtz, K. C. M.; Shen, L.; Douglas, C. J. *J. Am. Chem. Soc.* **2003**, *125*, 2368. (b) Zhang, Y.; Hsung, R. P.; Tracey, M. R.; Kurtz, K. C. M.; Vera, E. L. *Org. Lett.* **2004**, *6*, 1151. (c) Zhang, X.; Zhang, Y.; Huang, J.; Hsung, R. P.; Kurtz, K. C. M.; Oppenheimer, J.; Petersen, M. E.; Sagamanova, I. K.; Shen, L.; Tracey, M. R. *J. Org. Chem.* **2006**, *71*, 4170. (d) Coste, A.; Karthikeyan, G.; Couty, F.; Evano, G. *Angew. Chem., Int. Ed.* **2009**, *48*, 4381. (e) Coste, A.; Couty, F.; Evano, G. *Org. Lett.* **2009**, *11*, 4454. (f) Hamada, T.; Ye, X.; Stahl, S. S. *J. Am. Chem. Soc.* **2008**, *130*, 833. (g) Jia, W.; Jiao, N. *Org. Lett.* **2010**, *12*, 2000. (h) Jouvin, K.; Couty, F.; Evano, G. *Org. Lett.* **2010**, *12*, 3272. (i) Sueda, T.; Oshima, A.; Teno, N. *Org. Lett.* **2011**, *13*, 3996. (j) Laouiti, A.; Rammah, M. M.; Rammah, M. B.; Marrot, J.; Couty, F.; Evano, G. *Org. Lett.* **2012**, *14*, 6. (k) Jouvin, K.; Heimbürger, J.; Evano, G. *Chem. Sci.* **2012**, *3*, 756. (l) Laouiti, A.; Jouvin, K.; Bourdreux, F.; Rammah, M. M.; Rammah, M. B.; Evano, G. *Synthesis* **2012**, *44*, 1491.

(2) (a) DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, *110*, 5064. (b) Evano, G.; Coste, A.; Jouvin, K. *Angew. Chem., Int. Ed.* **2010**, *49*, 2840. (c) Evano, G.; Jouvin, K.; Coste, A. *Synthesis* **2013**, *45*, 17.

(3) (a) Mulder, J. A.; Kurtz, K. C. M.; Hsung, R. P.; Coverdale, H.; Frederick, M. O.; Shen, L.; Zificsak, C. A. *Org. Lett.* **2003**, *5*, 1547. (b) Zhang, X.; Zhang, Y.; Huang, J.; Hsung, R. P.; Kurtz, K. C. M.; Oppenheimer, J.; Petersen, M. E.; Sagamanova, I. K.; Shen, L.; Tracey, M. R. *J. Org. Chem.* **2006**, *71*, 4170. (c) Cao, J.; Kong, Y.; Deng, Y.; Lai, G.; Cui, Y.; Hu, Z.; Wang, G. *Org. Biomol. Chem.* **2012**, *10*, 9556.

(4) (a) Kramer, S.; Dooleweerd, K.; Lindhardt, A. T.; Rottländer, M.; Skrydstrup, T. *Org. Lett.* **2009**, *11*, 4208. (b) Shindoh, N.; Takemoto, Y.; Takasu, K. *Chem.—Eur. J.* **2009**, *15*, 7026.


(5) (a) Dunetz, J. R.; Danheiser, R. L. *J. Am. Chem. Soc.* **2005**, *127*, 5776. (b) Witulski, B.; Stengel, T. *Angew. Chem., Int. Ed.* **1999**, *38*, 2426. (c) Zhang, Y.; Hsung, R. P.; Zhang, X.; Huang, J.; Slater, B. W.; Davis, A. *Org. Lett.* **2005**, *7*, 1047. (d) Yao, P.-Y.; Zhang, Y.; Hsung, R. P.; Zhao, K. *Org. Lett.* **2008**, *10*, 4275. (e) Witulski, B.; Alayrac, C.; Tevzadze-Saefel, L. *Angew. Chem., Int. Ed.* **2003**, *42*, 4257. (f) Kramer, S.; Dooleweerd, K.; Lindhardt, A. T.; Rottländer, M.; Skrydstrup, T. *Org. Lett.* **2009**, *11*, 4208. (g) Cao, J.; Xu, Y.; Kong, Y.; Cui, Y.; Hu, Z.; Wang, G.; Deng, Y.; Lai, G. *Org. Lett.* **2012**, *14*, 38.

(6) (a) Movassaghi, M.; Hill, M. D.; Ahmad, O. *J. Am. Chem. Soc.* **2007**, *129*, 10096. (b) Garcia, P.; Moulin, S.; Miclo, Y.; Leboeuf, D.; Gandon, V.; Aubert, C.; Malacria, M. *Chem.—Eur. J.* **2009**, *15*, 2129. (c) Tanaka, R.; Yuza, A.; Watai, Y.; Suzuki, D.; Takayama, Y.; Sato, F.; Urabe, H. *J. Am. Chem. Soc.* **2005**, *127*, 7774. (d) Garcia, P.; Evanno, Y.; George, P.; Sevrin, M.; Ricci, G.; Malacria, M.; Aubert, C.; Gandon, V. *Org. Lett.* **2011**, *13*, 2030. (e) Gati, W.; Rammah, M. M.; Rammah, M. B.; Couty, F.; Evano, G. *J. Am. Chem. Soc.* **2012**, *134*, 9078.

(7) (a) Ziegert, R. E.; Toraeng, J.; Knepper, K.; Braese, S. *J. Comb. Chem.* **2005**, *7*, 147. (b) Hou, X.-L.; Yang, Z.; Yeung, K.-S.; Wong, H. N. C. *Prog. Heterocycl. Chem.* **2005**, *17*, 142.

Beckmann rearrangement of aryl benzofuranyl oximes,⁹ Curtius rearrangement of benzofuran-2-carboxylic acid,¹⁰ and nitration/reduction/*N*-acylation of 2-trimethylstannyl benzofurans.¹¹ Recently, Hsung and co-workers developed an elegant synthesis of 2-amidobenzofurans from *O*-anisyl ynamides via a Rh(I)-catalyzed demethylation–cyclization reaction,¹² and the Skrydstrup group reported a Pd-catalyzed, one-pot, two-step synthesis of 2-amidobenzofurans from ynamides and *o*-iodophenol.¹³ However, these methods could only afford 3-unsubstituted 2-amidobenzofurans. Electrophilic cyclization chemistry developed by Larock and co-workers provided access to various heterocyclic compounds including benzofurans.¹⁴ Many electrophilic reagents (I, Br, PhSe, PhS, etc.) could induce the electrophilic cyclization of a carbon–carbon triple bond (Scheme 1a). To the best of our knowledge, carbocation-induced electrophilic cyclization of alkynes is still unreported. Considering that ynamides are more electron-rich than ordinary alkynes because of the donating ability of the nitrogen lone pair toward the alkynyl motif, we anticipate that ynamides would undergo electrophilic cyclization reactions with carbocations to form a C–C and C–O bond (Scheme 1, b). Herein, we wish to report the novel synthesis of 3-substituted 2-amidobenzofurans via electrophilic cyclization of *o*-anisole-substituted ynamides with carbocations.

Scheme 1



(8) (a) Makovetskii, V. P.; Dzvinchuk, I. B.; Yaremko, V. V.; Svishchuk, A. A. *Ukr. Khim. Zh.* **1979**, *45*, 652. (b) King, T. J.; Newall, C. E. *J. Chem. Soc.* **1965**, 974.

(9) Demerseman, P.; Guillaumel, J.; Cheutin, A.; Royer, R. *Bull. Soc. Chim. Fr.* **1970**, 2253.

(10) Carrër, A.; Florent, J.; Auvrouin, E.; Rousselle, P.; Bertounesque, E. *J. Org. Chem.* **2011**, *76*, 2502.

(11) Einhorn, J. F.; Demerseman, P.; Royer, R. *Synthesis* **1984**, 978.


(12) Oppenheimer, J.; Johnson, W. L.; Tracey, M. R.; Hsung, R. P.; Yao, P.-Y.; Liu, R.; Zhao, K. *Org. Lett.* **2007**, *9*, 2361.

(13) Dooleweerd, K.; Ruhland, T.; Skrydstrup, T. *Org. Lett.* **2009**, *11*, 221.

(14) For electrophilic cyclization chemistry, see: (a) Yue, D.; Yao, T.; Larock, R. C. *J. Org. Chem.* **2005**, *70*, 10292. (b) Cho, C.-H.; Neuenswander, B.; Lushington, G.; Larock, R. C. *J. Comb. Chem.* **2008**, *10*, 941. (c) Manarin, F.; Roehrs, J. A.; Mozzaquatro Gay, R.; Brandão, R.; Menezes, P. H.; Nogueira, C. W.; Zeni, G. *J. Org. Chem.* **2009**, *74*, 2153. (d) Yao, T.; Yue, D.; Larock, R. C. *J. Org. Chem.* **2005**, *70*, 9985. (e) Mehta, S.; Waldo, J. P.; Larock, R. C. *J. Org. Chem.* **2009**, *74*, 1141. (f) Mehta, S.; Larock, R. C. *J. Org. Chem.* **2010**, *75*, 1652.

Initial studies focused on the reaction of *O*-anisyl ynamide **1a** with diphenylmethanol **2a**. ZnCl₂ was first chosen as the Lewis acid to generate the carbocation with diphenylmethanol in CH₂Cl₂. Fortunately, the expected 2-amidobenzofuran **3a** was obtained in 37% yield. Subsequent screening of Lewis acids showed that ZnBr₂ gave a better result than ZnCl₂ or ZnI₂ did (Table 1, entries 1–3), while other Lewis acids such as AlCl₃, FeCl₃, and BF₃·OEt₂ were not effective (entries 4–6). Solvents were also important. Other common solvents such as toluene or THF did not give any expected product **3a** (entries 7–8). It should be noted that when a catalytic amount of ZnBr₂ was used, the yield was reduced dramatically (entry 9). Thus the following reaction conditions were chosen as optimum for all subsequent reactions: 0.5 mmol of **1**, 1.0 mmol of **2**, and 1.0 mmol of ZnBr₂ in CH₂Cl₂ were stirred at rt under N₂.

Table 1. Optimization of the Reaction Conditions^a



entry	Lewis acid	solvent	time (h)	yield of 3a (%)
1	ZnCl ₂	CH ₂ Cl ₂	4	37
2	ZnBr ₂	CH ₂ Cl ₂	3	72
3	ZnI ₂	CH ₂ Cl ₂	3	66
4	AlCl ₃	CH ₂ Cl ₂	14	0
5	FeCl ₃	CH ₂ Cl ₂	14	0
6	BF ₃ ·OEt ₂	CH ₂ Cl ₂	1	trace
7	ZnBr ₂	toluene	1	0
8	ZnBr ₂	THF	12	NR
9 ^b	ZnBr ₂	CH ₂ Cl ₂	12	7

^a Unless specified otherwise, the reaction was carried out using **1a** (0.5 mmol), **2a** (1.0 mmol), and a Lewis acid (1.0 mmol) in solvent (4 mL) at rt under N₂. ^b 0.05 mmol of ZnBr₂ was added.

Under the optimized conditions, the scope of this carbocation-induced electrophilic cyclization reaction was further investigated. The reaction was successful for various *O*-anisyl ynamides (Table 2, entries 1–5). It should be noted that when the amido group in **1** was replaced by an aryl or alkyl group, the corresponding 3-alkylbenzofuran was not detected. Substituted diarylmethanol, 1-arylprop-2-en-1-ol, and 1-arylprop-2-yn-1-ol worked well to afford the corresponding 3-alkyl, allyl, and propargyl 2-amidobenzofuran respectively (Table 2, entries 6–9), while simple phenylmethanol could not undergo this electrophilic cyclization reaction.

Interestingly, 3-allenyl-2-amidobenzofurans **5** were formed when 1,1-diarylprop-2-yn-1-ols **4** were used as the precursors of carbocations, and the corresponding 3-propargyl-2-amidobenzofurans were not found. As shown in Table 3, the reaction was successful for various *O*-anisyl ynamides (entries 1–4). The R³ and R⁴ groups of **4** can be a substituted

Table 2. Synthesis of 3-Alkyl-2-amidobenzofurans **3**^a

entry	1	2	yield of 3
1			 3a 72%
2		2a	 3b 84%
3		2a	 3c 54%
4		2a	 3d 69%
5		2a	 3e 76%
6	1c		 3f 60%
7	1b		 3g 54%
8	1a		 3h 71%
9	1c		 3i 58%


^a The reaction was carried out using **1** (0.5 mmol), **2** (1.0 mmol), and ZnBr₂ (1.0 mmol) in CH₂Cl₂ (4 mL) at rt under N₂.

Table 3. Synthesis of 3-Alkenyl-2-amidobenzofurans **5**^a

entry	1	4	yield of 5
1	1a		 5a 65%
2	1b	4a	 5b 88%
3	1c	4a	 5c 75%
4			 5d 85%
5	1f		 5e 62%
6	1b		 5f 82%
7	1b		 5g 89%
8	1b		 5h 70%

^a The reaction was carried out using **1** (0.5 mmol), **4** (1.0 mmol), and ZnBr₂ (1.0 mmol) in CH₂Cl₂ (4 mL) at rt under N₂.

Scheme 2



phenyl group with either an electron-donating or -withdrawing group (entries 6–7), while R^5 can be an aryl or alkyl group (entries 4, 5, 8). Thus this reaction provided access to benzofuran-substituted allenes which are important synthons in modern organic synthesis.¹⁵

The ESI-TOF MS analysis of the reaction mixture of **1b** and **2a** showed the existence of a diphenylmethyl cation,

(15) (a) Krause, N.; Winter, C. *Chem. Rev.* **2011**, *111*, 1994. (b) Aubert, C.; Fensterbank, L.; Garcia, P.; Malacria, M.; Simonneau, A. *Chem. Rev.* **2011**, *111*, 1954. (c) Yu, S.; Ma, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 3074.

(16) For the generation of keteniminium ions from ynamides, see: (a) Zhang, Y.; Hsung, R. P.; Zhang, X.; Huang, J.; Slater, B. W.; Davis, A. *Org. Lett.* **2005**, *7*, 1047. (b) Mulder, J. A.; Hsung, R. P.; Frederick, M. O.; Tracey, M. R.; Zifcsak, C. A. *Org. Lett.* **2002**, *4*, 1383.

while the byproduct MeBr was obviously observed by GC-MS analysis of the gas composition of the reaction (see the Supporting Information for details). On the basis of the above experimental observations, a possible mechanism is given in Scheme 2. Initially, carbocation **A** would be formed from alcohol **2** with $ZnBr_2$. Electrophilic addition of the in situ generated carbocation with *O*-anisyl ynamide **1** gives the intermediate **B**.¹⁶ Subsequent cyclization occurred to afford *O*-methyl benzofuran **C**, which is demethylated by a bromide anion to produce 2-amidobenzofuran **3**. In the case of alcohol **4**, propargylic carbocation/allenyl carbocation **D** formed from **4** and $ZnBr_2$ reacts with ynamide **1** to afford 3-allenyl-2-amidobenzofuran **5** selectively probably because of the steric hindrance of the two aryl groups.

In conclusion, we have described a novel carbocation-induced electrophilic cyclization reaction of *o*-anisole-substituted ynamides. 3-Alkyl, allyl, and propargyl 2-amidobenzofuran were provided respectively from various disubstituted alcohols. By utilization of 1,1-diarylprop-2-yn-1-ol, 3-allenyl-2-amidobenzofurans were afforded in high yield. Further investigation for the construction of other heterocyclic and carbocyclic systems via this carbocation-induced electrophilic cyclization reaction is ongoing in our laboratory.

Acknowledgment. We are grateful to the National Natural Science Foundation of China (Project No. 21102029) for financial support.

Supporting Information Available. Spectroscopic data for all new compounds. Detailed experimental procedures. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.