

# Catalytic Asymmetric $\alpha$ -Aldol Reaction of Vinylogous *N*-Heterocyclic Carbene Enolates: Formation of Quaternary and Labile Tertiary Stereocenters

Xiuqin Dong and Jianwei Sun\*

Department of Chemistry, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR, China

**S** Supporting Information


**ABSTRACT:** Simple *N*-heterocyclic carbene (NHC)-enolates are widely studied versatile species. However, their vinylogous siblings (i.e., vinylogous NHC-enolates) have been much less studied. Here we disclose the first catalytic asymmetric  $\alpha$ -aldol reaction of vinylogous NHC-enolates. With trifluoropyruvate as the carbon electrophile, the efficient C–C bond formation process displays not only complete  $\alpha$ -regioselectivity but also excellent stereocontrol over the two newly established challenging stereocenters (one quaternary and the other labile tertiary), furnishing a range of highly enantioenriched  $\beta,\gamma$ -unsaturated  $\alpha$ -fluoroalkylated esters.



Asymmetric C–C bond formation is fundamentally important in organic chemistry.<sup>1</sup> In the past decade, organocatalytic C–C bond formation has seen exponential development, providing an attractive complement to the previously dominant metal- and biocatalytic approaches.<sup>2</sup> In particular, *N*-heterocyclic carbene (NHC) catalysis has emerged as a versatile platform for the discovery of new C–C bond-forming reactions.<sup>3</sup> Notably, these processes are typically highly enantioselective because of the excellent chiral transfer ability of the NHC-bound “umpolung” reagents, such as acyl anion equivalents, enolates, homoenolates, etc. Among them, NHC-enolates have received tremendous attention due to their versatility in reacting with various electrophiles to form  $\alpha$ -chiral carboxylic acid derivatives, thus representing an attractive substitute to the traditional metal- and chiral auxiliary-based enolates.<sup>4–9</sup> However, in terms of C–C bond formation, the majority of these reactions are cycloaddition reactions (e.g., [4 + 2], [2 + 2], [2 + 2 + 2]).<sup>10</sup> Intermolecular asymmetric  $\alpha$ -aldol-type reactions of NHC-enolates that do not involve ring formation are scarce.<sup>11</sup>

Furthermore, in contrast to the intensive studies of simple NHC-enolates, the reactivity of their vinylogous siblings (i.e., vinylogous NHC-enolates) have been much less studied,<sup>12,13</sup> despite the widely accepted importance of vinylogous enolates in remote C–C bond formation.<sup>14</sup> Ye and Chi pioneered the catalytic asymmetric C–C bond formation reactions of vinylogous NHC-enolates, but these reactions are again limited to cycloaddition at the  $\gamma$  position (Scheme 1).<sup>12</sup> To the best of our knowledge, C–C bond formation at the  $\alpha$  position of vinylogous NHC-enolates is unknown. In continuation of our effort in NHC catalysis,<sup>13a,15</sup> here we report an efficient C–C bond formation that takes place not only selectively at the  $\alpha$  position but also without cycloaddition. Specifically, considering the importance of fluoro- and trifluoromethyl-substituted molecules in medicinal chemistry and materials science,<sup>16</sup> we

## Scheme 1. C–C Bond Formation of Vinylogous NHC-Enolates




employed trifluoropyruvate as the carbon electrophile. Thus, the new C–C bond is formed between quaternary and labile tertiary (both allylic and  $\alpha$ -carbonyl) stereocenters,<sup>17</sup> which is a challenge in terms of stereocontrol, particularly for acyclic products (Scheme 1).

We started the study with racemic  $\gamma$ -reducible enal **1a** as the model substrate to generate the vinylogous NHC enolate and methyl trifluoropyruvate **2a** as the carbon electrophile (Table 1). Due to the presence of a labile (racemizable) tertiary stereocenter in the desired product, the weak base  $K_2HPO_4$  was employed. Initial evaluation of various NHC precatalysts with DCM as the solvent and MeOH as the nucleophile indicated that the desired product **3a** could be formed in low to moderate yield, but with good diastereoselectivity and excellent enantioselectivity (entries 1–6). Among them, triazolium salt

Received: March 19, 2014

Published: April 11, 2014

Table 1. Reaction Condition Optimization<sup>a</sup>

entry	precat.	base	solvent	yield ( <b>3a</b> ) <sup>b</sup>	dr ( <b>3a</b> ) <sup>c</sup>	ee ( <b>3a</b> ) <sup>d</sup>	<b>3a/4</b> <sup>e</sup>
1	<b>A</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	18%	15.0:1	96%	86:14
2	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	57%	10.8:1	94%	84:16
3	<b>C</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	trace	—	—	—
4	<b>D</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	trace	—	—	—
5	<b>E</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	trace	—	—	—
6	<b>F</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM	26%	6.0:1	90%	67:33
7	<b>B</b>	K <sub>3</sub> PO <sub>4</sub>	DCM	44%	8.3:1	93%	79:21
8	<b>B</b>	NaOAc	DCM	47%	8.5:1	92%	64:36
9	<b>B</b>	Cs <sub>2</sub> CO <sub>3</sub>	DCM	37%	1:1.2	83%	86:14
10	<b>B</b>	TEA	DCM	46%	5.5:1	77%	52:48
11	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	<i>n</i> -hexane	29%	1.1:1	88%	53:47
12	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	Et <sub>2</sub> O	26%	5.0:1	87%	67:33
13	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	MeCN	44%	3.3:1	93%	100:0
14	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	EtOAc	43%	5.3:1	93%	84:16
15	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	MeNO <sub>2</sub>	18%	3.7:1	91%	58:42
16	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM/MeCN <sup>g</sup>	66%	9.5:1	96%	100:0
17 <sup>f</sup>	<b>B</b>	K <sub>2</sub> HPO <sub>4</sub>	DCM/MeCN <sup>g</sup>	75%	9.1:1	96%	100:0

<sup>a</sup>The reaction was run with **1a** (0.05 mmol), **2a** (0.06 mmol), precat. (0.01 mmol), K<sub>2</sub>HPO<sub>4</sub> (0.1 mmol), MeOH (0.25 mmol), and solvent (0.5 mL). <sup>b</sup>Combined yield of the diastereomers determined by <sup>1</sup>HNMR analysis of the crude reaction mixture with CH<sub>2</sub>Br<sub>2</sub> as an internal standard. <sup>c</sup>The ratio (dr and **3a/4**) was determined by <sup>1</sup>HNMR analysis of the crude reaction mixture. <sup>d</sup>The ee value of the major diastereomer determined by HPLC analysis. <sup>e</sup>DCM/MeCN = 1:2 (v/v), 1.0 mL. <sup>f</sup>2.0 equiv of **2a**.

**B** gave the most promising results (entry 2).<sup>18</sup> Byproduct **4** was also observed, as a result of  $\alpha$ -protonation instead of C–C bond formation of the corresponding vinylogous enolate. Next, various bases were screened. Other weak bases K<sub>3</sub>PO<sub>4</sub> and NaOAc did not improve the product yield and diastereoselectivity. Relatively strong bases, such as Cs<sub>2</sub>CO<sub>3</sub> and Et<sub>3</sub>N, resulted in decreased stereoselectivity, presumably due to the base-labile  $\alpha$  tertiary stereocenter. Subsequent solvent screening did not improve the results either (entries 11–15), but interestingly we found that the formation of byproduct **4** was suppressed with MeCN as solvent (entry 13). Inspired by this observation, we next resorted to mixed solvent using DCM and MeCN (v/v = 1:2), which slightly improved the product yield (entry 16). Finally, increasing the loading of **2a** resulted in the further enhanced yield of **4a** (75%) together with good diastereoselectivity and excellent enantioselectivity (entry 17).

With the optimized standard reaction conditions, next we carried out a scope study. As shown in Table 2, a wide range of  $\gamma$  aryl- and alkyl-substituted enals **1** could participate smoothly in the C–C bond formation process, and desired  $\beta,\gamma$ -unsaturated  $\alpha$ -alkylated esters **3** were formed with good to excellent stereoselectivity. Ethyl trifluoropyruvate also provided similar levels of efficiency and selectivity (entries 14–19). Different alcohols, such as BnOH, allyl alcohol, and propargyl alcohol, could also serve as effective nucleophiles to form the corresponding ester products (entries 21–24). It is noteworthy

Table 2. Scope Study<sup>a</sup>

entry	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	time (h)	<b>3</b>	yield <sup>b</sup>	dr <sup>c</sup>	ee <sup>d</sup>
1	Ph	Me	Me	12	<b>3a</b>	82%	7.1:1	96%
2	2-MeC <sub>6</sub> H <sub>4</sub>	Me	Me	12	<b>3b</b>	74%	7.5:1	96%
3	2-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Me	Me	16	<b>3c</b>	46%	6.2:1	92%
4 <sup>g</sup>	2-(vinyl)C <sub>6</sub> H <sub>4</sub>	Me	Me	90	<b>3d</b>	69%	10.3:1	97%
5	2-(TIPSOCH <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	Me	Me	16	<b>3e</b>	78%	10.7:1	96%
6	3-(MeO)C <sub>6</sub> H <sub>4</sub>	Me	Me	12	<b>3f</b>	65%	6.2:1	96%
7 <sup>g</sup>	3-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Me	Me	90	<b>3g</b>	56%	6.8:1	98%
8	3,5-(CH <sub>3</sub> ) <sub>2</sub> C <sub>6</sub> H <sub>3</sub>	Me	Me	16	<b>3h</b>	73%	9.1:1	96%
9	4-MeC <sub>6</sub> H <sub>4</sub>	Me	Me	16	<b>3i</b>	71%	6.5:1	96%
10	4-(MeO)C <sub>6</sub> H <sub>4</sub>	Me	Me	12	<b>3j</b>	79%	6.6:1	96%
11	4-FC <sub>6</sub> H <sub>4</sub>	Me	Me	12	<b>3k</b>	64%	7.7:1	96%
12 <sup>g</sup>	4-(vinyl)C <sub>6</sub> H <sub>4</sub>	Me	Me	90	<b>3l</b>	81%	9.5:1	98%
13		Me	Me	21	<b>3m</b>	63%	11.8:1	97%
14		Et	Me	22	<b>3n</b>	63%	11.8:1	95%
15	Ph	Et	Me	12	<b>3o</b>	80%	9.2:1	96%
16	2-MeC <sub>6</sub> H <sub>4</sub>	Et	Me	45	<b>3p</b>	72%	8.0:1	88%
17 <sup>g</sup>	2-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Et	Me	90	<b>3q</b>	70%	4.5:1	94%
18 <sup>g</sup>	3-(MeO)C <sub>6</sub> H <sub>4</sub>	Et	Me	90	<b>3r</b>	69%	8.0:1	98%
19 <sup>g</sup>	4-FC <sub>6</sub> H <sub>4</sub>	Et	Me	90	<b>3s</b>	74%	7.0:1	98%
20	<sup>t</sup> Bu	Me	Me	51	<b>3t</b>	55%	5.0:1	95%
21	Bn	Et	Bn	90	<b>3u</b>	41%	4.3:1	99%
22 <sup>g</sup>	Ph	Me	Bn	40	<b>3v</b>	80%	4.7:1	99%
23 <sup>g</sup>	Ph	Me	allyl	40	<b>3w</b>	79%	6.3:1	98%
24 <sup>g</sup>	Ph	Me	propargyl	40	<b>3x</b>	81%	5.3:1	96%

<sup>a</sup>The reaction was run with **1** (0.35 mmol), **2** (0.70 mmol), **B** (0.07 mmol), K<sub>2</sub>HPO<sub>4</sub> (0.7 mmol), R<sup>3</sup>OH (1.75 mmol), and solvent (DCM/MeCN = 1:2, v/v, 7.0 mL). <sup>b</sup>Isolated combined yield of the diastereomers. <sup>c</sup>The dr value was determined by <sup>1</sup>H NMR of the crude mixture. <sup>d</sup>The ee value was determined by HPLC. <sup>e</sup>Run at 0 °C.

that in all these examples the C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bonds between quaternary and labile tertiary stereocenters were all formed under mild conditions with excellent stereocontrol, which is particularly noteworthy for the formation of acyclic products.

In order to rationalize the absolute and relative stereochemistry of the two new stereocenters in the products, we have proposed conformation **I** for the vinylogous NHC-enolate and transition state **II** for the C–C formation step (Figure 1).





Figure 1. Plausible vinylogous NHC-enolate conformation and C–C bond formation transition state.

In conformation **I**, the chiral NHC backbone of the *Z*-enolate blocks the *Si* face and the electrophile approaches from the *Re* face (bottom face) for new bond formation, thus determining the absolute stereochemistry. In the C–C formation step, the orientation of the trifluoropyruvate determines the relative stereochemistry. We have proposed a six-membered chairlike transition state (**II**), in which a molecule of MeOH activates the electrophile carbonyl group through hydrogen bonding and simultaneously approaches the developing acyl carbon. In this

transition state, the large CF<sub>3</sub> group adopts the equatorial position (*A*-value of CF<sub>3</sub> 2.4–2.5 > *A*-value of CO<sub>2</sub>Me 1.2–1.3).<sup>19</sup> Thus, the enolate nucleophilic carbon attacks the *Re* face of the carbonyl, thereby setting the quaternary stereocenter. These rationalizations are in complete agreement with the observed product stereochemistry.

The highly enantioenriched products obtained via our catalytic intermolecular C–C bond-forming process can be transformed into other useful molecules. For example, the C=C bond in product **3a** can be efficiently hydrogenated to form saturated diester **5** without erosion in stereoselectivity (eq 1).



Moreover, due to the presence of a homoallylic alcohol subunit, product **3a** could undergo efficient iodoetherification in the presence of I<sub>2</sub> and NaHCO<sub>3</sub> to form densely functionalized tetrahydrofuran **6** as essentially a single diastereomer (eq 2). The relative stereochemistry was established by NOESY. It is worth noting that trifluoromethyl-substituted tetrahydrofuran subunits are of great interest in medicinal chemistry.<sup>20</sup>

In summary, we have disclosed an unprecedented intermolecular asymmetric  $\alpha$ -aldol reaction of vinyllogous NHC-enolates, a type of versatile but less explored species relative to simple NHC-enolates. In contrast to the known C–C bond formation at the  $\gamma$  position of vinyllogous NHC-enolates, our reaction exhibits complete  $\alpha$  selectivity. Unlike most cycloaddition reactions of NHC-enolates with external carbon electrophiles, our reaction does not involve a cycloaddition step. Notably, two challenging stereocenters, one quaternary and the other labile tertiary (both allylic and  $\alpha$ -carbonyl), are also established in an acyclic product with excellent absolute and relative stereocontrol. A range of highly enantioenriched  $\beta,\gamma$ -unsaturated  $\alpha$ -fluoroalkylated esters have been synthesized with high efficiency under mild conditions. These products can be easily transformed into other useful molecules, such as densely functionalized tetrahydrofurans. Further studies on vinyllogous NHC-enolates are underway.

## ■ ASSOCIATED CONTENT

### Supporting Information

Experimental procedures, full spectroscopic data, and NMR spectra for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: [sunjw@ust.hk](mailto:sunjw@ust.hk)

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

Financial support was provided by HKUST and Hong Kong RGC (ECS605812 and M-HKUST607/12).

## ■ REFERENCES

- (1) (a) Ojima, I. *Catalytic Asymmetric Synthesis*, 3rd ed.; John Wiley: Hoboken, NJ; 2010. (b) Christmann, M.; Bräse, S. *Asymmetric Synthesis II: More Methods and Applications*; Wiley-VCH: Hoboken, NJ, 2012.
- (2) For selected reviews, see: (a) Enders, D.; Narine, A. A. *J. Org. Chem.* **2008**, *73*, 7857. (b) Scheffler, U.; Mahrwald, R. *Chem.—Eur. J.* **2013**, *19*, 14346. (c) Mahrwald, R. *Drug Discovery Today: Technologies* **2013**, *10*, e29. (d) Volla, C. M. R.; Atodiresi, I.; Rueping, M. *Chem. Rev.* **2014**, *114*, 2390.
- (3) For recent reviews and highlights on NHC catalysis, see: (a) Moore, J. L.; Rovis, T. *Top. Curr. Chem.* **2011**, *291*, 77. (b) Vora, H. U.; Rovis, T. *Aldrichimica Acta* **2011**, *44*, 3. (c) Nair, V.; Menon, R. S.; Biju, A. T.; Sinu, C. R.; Paul, R. R.; Jose, A.; Sreekumar, V. *Chem. Soc. Rev.* **2011**, *40*, 5336. (d) Rong, Z.-Q.; Zhang, W.; Yang, G.-Q.; You, S.-L. *Curr. Org. Chem.* **2011**, *15*, 3077. (e) Biju, A. T.; Kuhl, N.; Glorius, F. *Acc. Chem. Res.* **2011**, *44*, 1182. (f) Chiang, P.-C.; Bode, J. W. *TCI MAIL* **2011**, *149*, 2. (g) Vora, H. U.; Wheeler, P.; Rovis, T. *Adv. Synth. Catal.* **2012**, *354*, 1617. (h) Bugaut, X.; Glorius, F. *Chem. Soc. Rev.* **2012**, *41*, 3511. (i) Izquierdo, J.; Hutson, G. E.; Cohen, D. T.; Scheidt, K. A. *Angew. Chem., Int. Ed.* **2012**, *51*, 11686. (j) Cohen, D. T.; Scheidt, K. A. *Chem. Sci.* **2012**, *3*, 53. (k) Grossmann, A.; Enders, D. *Angew. Chem., Int. Ed.* **2012**, *51*, 314. (l) Bode, J. W. *Nat. Chem.* **2013**, *5*, 813. (m) Sarkar, S. D.; Biswas, A.; Samanta, R. C.; Studer, A. *Chem.—Eur. J.* **2013**, *19*, 4664. (n) Ryan, S. J.; Candish, L.; Lupton, D. W. *Chem. Soc. Rev.* **2013**, *42*, 4906. (o) Chauhan, P.; Ender, D. *Angew. Chem., Int. Ed.* **2014**, *53*, 1485. (p) Mahatthanachai, J.; Bode, J. W. *Acc. Chem. Res.* **2014**, *47*, 696.
- (4) For reviews/articles on azolium enolate generation and reactivity, see: (a) Douglas, J.; Churchill, G.; Smith, A. D. *Synthesis* **2012**, *44*, 2295. (b) Chen, X.-Y.; Ye, S. *Synlett* **2013**, *24*, 1614. (c) Maji, B.; Mayr, H. *Angew. Chem., Int. Ed.* **2013**, *52*, 11163.
- (5) For pioneering work involving NHC-enol/enolates, see: (a) Reynolds, N. T.; De Alaniz, J. R.; Rovis, T. *J. Am. Chem. Soc.* **2004**, *126*, 9518. (b) Reynolds, N. T.; Rovis, T. *J. Am. Chem. Soc.* **2005**, *127*, 16406.
- (6) For examples of intermolecular C–C bond formation of NHC-enolates: (a) He, M.; Struble, J. R.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 8418. (b) He, M.; Uc, G. J.; Bode, J. W. *J. Am. Chem. Soc.* **2006**, *128*, 15088. (c) He, M.; Beahm, B. J.; Bode, J. W. *Org. Lett.* **2008**, *10*, 3817. (d) Zhang, Y.-R.; He, L.; Wu, X.; Shao, P.-L.; Ye, S. *Org. Lett.* **2008**, *10*, 277. (e) Duguet, N.; Campbell, C. D.; Slawin, A. M. Z.; Smith, A. D. *Org. Biomol. Chem.* **2008**, *6*, 1108. (f) Zhang, Y. R.; Lv, H.; Zhou, D.; Ye, S. *Chem.—Eur. J.* **2008**, *14*, 8473. (g) Wang, X.-N.; Shao, P.-L.; Lv, H.; Ye, S. *Org. Lett.* **2009**, *11*, 4029. (h) Lv, H.; Chen, X.-Y.; Sun, L.-H.; Ye, S. *J. Org. Chem.* **2010**, *75*, 6973. (i) Kaeobamrung, J.; Kozlowski, M. C.; Bode, J. W. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 20661. (j) Jian, T.-Y.; Shao, P.-L.; Ye, S. *Chem. Commun.* **2011**, *47*, 2381. (k) Wang, X.-N.; Shen, L.-T.; Ye, S. *Org. Lett.* **2011**, *13*, 6382. (l) Rong, Z.-Q.; Jia, M.-Q.; You, S.-L. *Tetrahedron* **2011**, *67*, 9329. (m) Fang, X.; Chen, X.; Chi, Y. R. *Org. Lett.* **2011**, *13*, 4708. (n) Lv, H.; Mo, J.; Fang, X.; Chi, Y. R. *Org. Lett.* **2011**, *13*, 5366. (o) Hao, L.; Du, Y.; Lv, H.; Chen, X.; Jiang, H.; Shao, Y.; Chi, Y. R. *Org. Lett.* **2012**, *14*, 2154. (p) Yang, L.; Wang, F.; Chua, P. J.; Lv, Y.; Zhong, L.-J.; Zhong, G. *Org. Lett.* **2012**, *14*, 2894. (q) Zhao, X.; Ruhl, K. E.; Rovis, T. *Angew. Chem., Int. Ed.* **2012**, *51*, 12330. (r) Fu, Z.; Sun, H.; Chen, S.; Tiwari, B.; Li, G.; Chi, Y. R. *Chem. Commun.* **2013**, *49*, 261. (s) Leckie, S. M.; Brown, T. B.; Pryde, D.; Lebl, T.; Slawin, A. M. Z.; Smith, A. D. *Org. Biomol. Chem.* **2013**, *11*, 3230. (t) Davies, A. T.; Taylor, J. E.; Douglas, J.; Collett, C. J.; Morrill, L. C.; Fallan, C.; Slawin, A. M. Z.; Churchill, G.; Smith, A. D. *J. Org. Chem.* **2013**, *78*, 9243. (u) Douglas, J.; Taylor, J. E.; Churchill, G.; Slawin, A. M. Z.; Smith, A. D. *J. Org. Chem.* **2013**, *78*, 3925. (v) McCusker, E. O.; Scheidt, K. A. *Angew. Chem., Int. Ed.* **2013**, *52*, 13616. (w) Ni, Q.; Zhang, H.; Grossmann, A.; Loh, C. C. J.; Merckens, C.; Enders, D. *Angew. Chem., Int. Ed.* **2013**, *52*, 13562.
- (7) For examples involving C–C bond formation between an NHC-enolate and an intramolecular carbon electrophile, see: (a) Chiang, P.-C.; Kaeobamrung, J.; Bode, J. W. *J. Am. Chem. Soc.* **2007**, *129*, 3520.



(b) Wadamoto, M.; Phillips, E. M.; Reynolds, T. E.; Scheidt, K. A. *J. Am. Chem. Soc.* **2007**, *129*, 10098. (c) Phillips, E. M.; Wadamoto, M.; Chan, A.; Scheidt, K. A. *Angew. Chem., Int. Ed.* **2007**, *46*, 3107. (d) He, M.; Bode, J. W. *J. Am. Chem. Soc.* **2008**, *130*, 418. (e) Li, Y.; Wang, X.-Q.; Zheng, C.; You, S.-L. *Chem. Commun.* **2009**, 5823. (f) Cardinal-David, B.; Raup, D. E. A.; Scheidt, K. A. *J. Am. Chem. Soc.* **2010**, *132*, 5345. (g) Cohen, D. T.; Cardinal-David, B.; Scheidt, K. A. *Angew. Chem., Int. Ed.* **2011**, *50*, 1678. (h) Fang, X.; Jiang, K.; Xing, C.; Hao, L.; Chi, Y. R. *Angew. Chem., Int. Ed.* **2011**, *50*, 1910. (i) Biswas, A.; De Sarkar, S.; Tebben, L.; Studer, A. *Chem. Commun.* **2012**, *48*, 5190. (j) Cohen, D. T.; Eichman, C. C.; Phillips, E. M.; Zarefsky, E. R.; Scheidt, K. A. *Angew. Chem., Int. Ed.* **2012**, *51*, 7309. (k) Fu, Z.; Xu, J.; Zhu, T.; Leong, W. W. Y.; Chi, Y. R. *Nat. Chem.* **2013**, *5*, 835.

(8) For asymmetric  $\alpha$  protonation of NHC-enolates, see ref 5b and: (a) Wang, X.-N.; Lv, H.; Huang, X.-L.; Ye, S. *Org. Biomol. Chem.* **2009**, *7*, 346. (b) Vora, H. U.; Rovis, T. *J. Am. Chem. Soc.* **2010**, *132*, 2860. (c) Concellón, C.; Duguet, N.; Smith, A. D. *Adv. Synth. Catal.* **2009**, *351*, 3001.

(9) Asymmetric  $\alpha$  carbon–heteroatom bond formation of NHC-enolates: (a) Huang, X.-L.; He, L.; Shao, P.-L.; Ye, S. *Angew. Chem., Int. Ed.* **2009**, *48*, 192. (b) Douglas, J.; K. Ling, B.; Concellón, C.; Churchill, G.; Slawin, A. M. Z.; Smith, A. D. *Eur. J. Org. Chem.* **2010**, 5863. (c) Shao, P.-L.; Chen, X.-Y.; Ye, S. *Angew. Chem., Int. Ed.* **2010**, *49*, 8412. (d) Jian, T.-Y.; He, L.; Tang, C.; Ye, S. *Angew. Chem., Int. Ed.* **2011**, *50*, 9104. (e) Taylor, J. E.; Daniels, D. S. B.; Smith, A. D. *Org. Lett.* **2013**, *15*, 6058.

(10) It is presumably because of the favorable thermodynamics in the ring-closing step to facilitate catalyst turnover. A concerted pathway for a [4 + 2] example was proposed based on calculations: Allen, S. E.; Mahhatthananchai, J.; Bode, J. W.; Kozłowski, M. C. *J. Am. Chem. Soc.* **2012**, *134*, 12098.

(11) For an example, which generates  $\beta$ - instead of  $\alpha$ -chirality, see: Kawanaka, Y.; Phillips, E. M.; Scheidt, K. A. *J. Am. Chem. Soc.* **2009**, *131*, 18028.

(12) Asymmetric C–C bond formation of vinylogous NHC enolates: (a) Shen, L.-T.; Shao, P.-L.; Ye, S. *Adv. Synth. Catal.* **2011**, *353*, 1943. (b) Mo, J.; Chen, X.; Chi, Y. R. *J. Am. Chem. Soc.* **2012**, *134*, 8810. (c) Chen, X.; Yang, S.; Song, B.-A.; Chi, Y. R. *Angew. Chem., Int. Ed.* **2013**, *52*, 11134. (d) Xu, J.; Jin, Z.; Chi, Y. R. *Org. Lett.* **2013**, *15*, 5028. (e) Wang, M.; Huang, Z.; Xu, J.; Chi, Y. R. *J. Am. Chem. Soc.* **2014**, *136*, 1214.

(13) Asymmetric carbon–heteroatom bond formation of vinylogous NHC enolates: (a) Zhao, Y.-M.; Cheung, M. S.; Lin, Z.; Sun, J. *Angew. Chem., Int. Ed.* **2012**, *51*, 10359. (b) Chen, X.-Y.; Xia, F.; Cheng, J.-T.; Ye, S. *Angew. Chem., Int. Ed.* **2013**, *52*, 10644.

(14) For reviews of vinylogous enolates (mostly for aldol reactions), see: (a) Casiraghi, G.; Zanardi, F. *Chem. Rev.* **2000**, *100*, 1929. (b) Denmark, S. E.; Heemstra, J. R.; Beutner, G. L. *Angew. Chem., Int. Ed.* **2005**, *44*, 4682. (c) Xu, B.; Hammond, G. B. *Synlett* **2010**, 1442. (d) Casiraghi, G.; Battistini, L.; Curti, C.; Rassa, G.; Zanardi, F. *Chem. Rev.* **2011**, *111*, 3076.

(15) (a) Zhao, Y.-M.; Tam, Y.; Wang, Y.-J.; Li, Z.; Sun, J. *Org. Lett.* **2012**, *14*, 1398. (b) Dong, X.; Zhao, Y.-M.; Sun, J. *Synlett* **2013**, *24*, 1221.

(16) (a) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881. (b) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320. (c) Jeschke, P. *ChemBioChem* **2004**, *5*, 570. (d) Hung, M. H.; Farnham, W. B.; Feiring, A. E.; Rozen, S. In *Fluoropolymers: Synthesis*; Hougham, G., Cassidy, P. E., Johns, K., Davidson, T., Eds.; Kluwer Academic/Plenum Publishers: New York, 1999; Vol. 1, p 51.

(17) The  $\alpha$  tertiary chirality of  $\beta,\gamma$ -unsaturated carbonyl compounds is potentially labile due to easy racemization, particularly for reactions under basic conditions. For examples of catalytic enantioselective formation of such stereocenters, see: (a) Davies, H. W.; Jin, Q. *J. Am. Chem. Soc.* **2004**, *126*, 10862. (b) Lou, S.; Fu, G. C. *J. Am. Chem. Soc.* **2010**, *132*, 5010. (c) Skucas, E.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2012**, *134*, 9090.

(18) Precatalyst **B** was first disclosed by Bode and co-workers. See ref 6a and: Struble, J. R.; Bode, J. W.; Lian, Y.; Davies, H. M. L. *Org. Synth.* **2010**, *87*, 362.

(19) Eliel, E. L.; Wilen, S. H. *Stereochemistry of Organic Compounds*; Wiley: New York, 1994; p 696.

(20) For selected examples, see: (a) Frezza, M.; Balestrino, D.; Souler, L.; Reverchon, S.; Queneau, Y.; Forestier, C.; Doutheau, A. *Eur. J. Org. Chem.* **2006**, 4731. (b) Hall, A.; Farthing, C. N.; Castro Pineiro, J. L. Patent WO 098213A1, 2012. (c) Alvaro, G.; Décor, A.; Hamprecht, D.; Marasco, A. Patent WO 168710A1, 2012.