

Enantioselective Total Synthesis of (–)-Nardoaristolone B via a Gold(I)-Catalyzed Oxidative Cyclization


Anna Homs,^{†,§} Michael E. Muratore,^{†,§} and Antonio M. Echavarren^{*,†,‡}

[†]Institute of Chemical Research of Catalonia (ICIQ), Av. Països Catalans 16, 43007 Tarragona, Spain

[‡]Departament de Química Analítica i Química Orgànica, Universitat Rovira i Virgili, C/Marcel·lí Domingo s/n, 43007 Tarragona, Spain

S Supporting Information

ABSTRACT: The first enantioselective total synthesis of (–)-nardoaristolone B is accomplished by the implementation of an enantio- and diastereoselective copper(I)-catalyzed conjugate addition/enolate trapping sequence and a gold(I)-catalyzed oxidative cyclization (intermolecular oxidant), employed for the first time in total synthesis.



Nardoaristolone B (**1**) was isolated in 2013 from *Nardostachys chinensis* Batal, a plant of the genus *Nardostachys* endemic of the Himalayan mountains.¹ The synthesis of the racemic mixture has been recently reported.² Closely related sesquiterpene (–)-aristolone (**2**) was isolated much earlier, in 1955, from the roots of *Aristolochia debilis*³ and has been synthesized in racemic form by various research groups.⁴ Nardoaristolone B (**1**) exhibits protective activity on the injury of neonatal rat cardiomyocytes.


We were intrigued by the possibility of accessing **1** and other members of the aristolone family by combining the highly efficient Cu(I)-catalyzed asymmetric conjugate addition/ α -alkylation cascade of α,β -unsaturated cyclic ketones developed by the groups of Alexakis and Cramer^{5–8} with the Au(I)-catalyzed oxidative cyclization of enynes recently discovered by Liu (Scheme 1).⁹ This last method, based on the gold(I)-catalyzed oxidative functionalization of alkynes pioneered by Toste¹⁰ and Zhang^{11,12} could offer direct access to this family of compounds from cyclohexanone **4** as the common inter-

mediate. Here we report the first enantioselective total synthesis of nardoaristolone B (**1**) and additional studies on the scope of the Au(I)-catalyzed oxidative cyclization of 1,6-enynes.

Although the conjugate methylation of 2-methyl-2-cyclohexenone proceeded satisfactorily at $-35\text{ }^{\circ}\text{C}$,^{5b} the subsequent α -alkylation proved to be challenging using methyl iodide. However, employing methyl iodide under high concentration and using 1:1 mixture of HMPA/THF before addition of MeI led to trisubstituted cyclohexanone **3** in 45–55% yield (Scheme 2). Employing the optimal chiral phosphoramidite ligand as reported by Alexakis,^{5a} the reaction proceeded with 91–92% enantiomeric excess and 3:1 dr. Careful purification by standard column chromatography allowed us to isolate **3** in essentially pure form (>30:1 dr). The isomerization of *exo*-olefin **3** into the corresponding trisubstituted *endo*-alkene **4** was not trivial, and a range of conditions was screened.¹³ Fortunately, the use of $\text{RhCl}_3 \cdot x\text{H}_2\text{O}$ (5 mol %) in ethanol at $75\text{ }^{\circ}\text{C}$ led to the desired *endo*-olefin **4** in 74% yield. The conversion of cyclohexanone **4** into enol triflate **5** was performed under standard conditions (82% yield). Sonogashira cross-coupling of **5** with trimethylsilyl acetylene employing $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (2 mol %) and CuI (5 mol %) in a DMF/ Et_3N mixture, followed by methanolysis of the TMS group led to 1,5-enyne **6** in satisfactory yield (74% over two steps) (Scheme 2).

Pleasingly, 1,5-enyne underwent the desired gold(I)-catalyzed oxidative cyclization using 8-methylquinoline *N*-oxide (PNOS) and $\text{I}^{\text{Pr}}\text{AuNTf}_2$ as catalyst in 1,2-dichloroethane ($(\text{CH}_2\text{Cl})_2$) at $80\text{ }^{\circ}\text{C}$,⁹ albeit with low isolated yield (20%, along with 25% of the simply cycloisomerized enyne **9**). Careful scrutiny of conditions revealed that the choice of the oxidant was crucial in order to favor the desired oxidative cyclization over the cycloisomerization. Thus, 3,5-dichloropyridine *N*-oxide (PNO3) proved to be superior to all the other *N*-oxides


Scheme 1. Synthetic Plan toward Enantioenriched (–)-Nardoaristolone B and (–)-Aristolone




Received: December 7, 2014

Published: January 7, 2015

Scheme 2. Synthesis of 1,5-Enyne 6




screened (Table 1, entries 3 and 11).^{13,14} Interestingly, using isomeric 2,6-dichloropyridine *N*-oxide (**PNO4**) led exclusively to diene **9** (Table 1, entry 4), whereas a complex mixture was

Table 1. Screening of Conditions for the Gold(I)-Catalyzed Oxidative Cyclization of **6**

entry	[Au]	oxidant	yield of 7/9 ^{a,b} (%)
1	IPrAuNTf ₂	PNO1	31/5
2	IPrAuNTf ₂	PNO2	20/36
3	IPrAuNTf ₂	PNO3	74/15
4	IPrAuNTf ₂	PNO4	0/55
5	IPrAuNTf ₂	PNO5	20/25
6	IPrAuNTf ₂	none	complex mixture
7	(JohnPhos)AuCl/AgNTf ₂	PNO3	43/15
8 ^c	<i>t</i> BuXPhosNTf ₂	PNO3	18/15
9	[(ArO) ₃ P]AuCl/AgNTf ₂	PNO3	55/2
10	IMesAuNTf ₂	PNO3	55/2
11	IPrAuNTf ₂ ^d	PNO3	74 (74) ^e /15



^aYields determined by ¹H NMR analysis of the crude mixture using diphenylmethane as internal standard. ^bFull conversion of starting material was observed unless otherwise stated. ^c13% unreacted starting material were also visible. ^d5 mol % of catalyst; Ar = 2,4-(*t*Bu)₂C₆H₃. ^eIsolated yield.



obtained under standard cycloisomerization conditions in the absence of any oxidant (Table 1, entry 6). The competitive formation of cycloisomerization product **9**, along with **7**, in these reactions suggests that both products result from a common cyclopropyl gold(I) intermediate.¹⁵ However, the alternative mechanism involving an earlier oxidation of the terminal alkyne to form an α -oxo gold(I) carbene intermediate that leads to **7** by intramolecular cyclopropanation, in parallel with a simple gold(I)-catalyzed cycloisomerization, cannot be excluded.⁹


With the optimal conditions in hand, we performed an oxidative cyclization in the presence of only 5 mol % of IPrAuNTf₂ and 3,5-dichloropyridine *N*-oxide. The desired product **7** was isolated in good yield (74%) along with 15% of cycloisomerized product **9**. The last step consisted in the allylic oxidation, which was accomplished in high yield (93%) using a Pd-catalyzed radical oxidation in the presence of Pearlman's catalyst (Pd(OH)₂/C) and *t*-BuOOH¹⁶ (Scheme 3). The

Scheme 3. Last Step in the Synthesis of (–)-Nardoaristolone B



spectroscopic data are in excellent agreement with the ones reported for the isolated compound and further support for the structure was obtained by X-ray diffraction analysis.¹⁷

Having accomplished the first total synthesis of nardoaristolone B, we were very keen on applying our strategy to a higher enyne in order to gain access to the core of the aristolone family of natural products. Our synthetic effort first involved the conversion of key intermediate enol triflate **5** into the corresponding 1,6-enyne **8**. Although the direct Kumada cross-coupling of **5** with propargylmagnesium bromide in the presence of various Pd- or Ni-based catalysts did not take place, we uncovered an unprecedented Kumada cross-coupling of TMS-protected propargylmagnesium bromide with enol triflates. This coupling proceeded smoothly on our model system (4-*tert*-butylcyclohexanone-derived enol triflate) employing only 2 mol % of Pd(PPh₃)₄ and 2 equiv of freshly prepared Grignard reagent.¹³ However, the cross-coupling was significantly slower on **5** and 20 mol % of Pd complex as well as 4 equiv of Grignard reagent were necessary to obtain full conversion at 23 °C. Under these conditions, the cross-coupling of **5** proceeded smoothly to afford **8** in 77% yield after methanolysis of the TMS group (Scheme 4). This substrate was then treated under a variety of conditions in order to prepare aristolone; however all attempts resulted in the formation of 6a-formyl-6-deoxonardoaristolone **10** in 65% yield when employing IPrAuNTf₂ as catalyst.¹⁵ Although (–)-aristolone (**2**) was

Scheme 4. Synthesis and Fate of 1,6-Enyne **8**

not prepared via our original strategy, the preparation of enantioenriched cyclohexanone **4** constitutes a formal synthesis of (–)-**2**, since racemic **2** has already been prepared from (±)-**4** in five steps.^{4b}

Under all the conditions examined, the gold(I)-catalyzed reaction of 1,6-enyne **8** proceeded exclusively by the 6-*exo-dig* mode. It is interesting that this result is in contrast to that observed in the reaction 1-ethynyl-2-allylbenzene, which yielded a 6-membered ring ketone by a 6-*endo-dig* oxidative cyclization.⁹ This different behavior can be ascribed to the different substitution pattern of the alkene, which usually controls the outcome in gold(I)-catalyzed cycloisomerizations of 1,6-enynes.¹⁸

In conclusion, we have developed the first enantioselective synthesis of (–)-nardoaristolone B (**1**) in seven steps and 14–17% overall yield. Our expedient strategy, by implementation of an enantio- and diastereoselective conjugate addition/enolate alkylation and the first example of a gold(I)-catalyzed oxidative cyclization of enynes in total synthesis, is perfectly suited for the rapid preparation of analogues of this natural product.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and characterization data for compounds **1** and **3–10** as well as the X-ray crystal structure of **1**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: aechavarren@icq.es.

Author Contributions

[§]These authors contributed equally.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank MINECO (Severo Ochoa Excellence Accreditation 2014–2018 (SEV-2013-0319), project CTQ2013-42106-P, and FPI predoctoral Fellowship to A.H.), the European Research Council (Advanced Grant No. 321066), the AGAUR (2014 SGR 818), and the ICIQ Foundation. M.E.M. acknowledges the receipt of a COFUND postdoctoral fellowship (Marie Curie program).

■ REFERENCES

- (1) Liu, M. A.; Duan, Y.-H.; Hou, Y.-L.; Li, C.; Gao, H.; Dai, Y.; Yao, X.-S. *Org. Lett.* **2013**, *15*, 1000–1003.
- (2) Handore, K. L.; Reddy, S. *Org. Lett.* **2014**, *16*, 4252–4255.
- (3) Kariyone, T.; Naito, S. *J. Pharm. Soc. Jpn.* **1955**, *75*, 1511–1516.
- (4) (a) Berger, C.; Franck-Neumann, M.; Ourisson, G. *Tetrahedron Lett.* **1968**, *9*, 3451–3452. (b) Piers, E.; Britton, R. W.; (de) Waal, W. *Can. J. Chem.* **1969**, *47*, 831–840. (c) Prasad, C. V. C.; Chan, T. H. *J. Org. Chem.* **1987**, *52*, 120–124.
- (5) (a) Vuagnoux-d'Augustin, M.; Alexakis, A. *Chem.–Eur. J.* **2007**, *13*, 9647–9662. (b) Ngoc, D. T.; Albicker, M.; Schneider, L.; Cramer, N. *Org. Biomol. Chem.* **2010**, *8*, 1781–1784.
- (6) For key pioneering studies on Cu(I)-catalyzed enantioselective conjugate additions of various organometallic reagents, see: (a) Alexakis, A.; Mutti, S.; Normant, J. F. *J. Am. Chem. Soc.* **1991**, *113*, 6332–6334. (b) Alexakis, A.; Frutos, J.; Mangeney, P. *Tetrahedron: Asymmetry* **1993**, *4*, 2427–2430. (c) Feringa, B. L.; Badorrey, R.; Peña, D.; Harutyunyan, S. R.; Minnaard, A. J. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5834–5838. (d) d'Augustin, M.; Palais, L.; Alexakis, A. *Angew. Chem., Int. Ed.* **2005**, *44*, 1376–1378.
- (7) For selected reviews on asymmetric conjugate additions, see: (a) Alexakis, A.; Benhaim, C. *Eur. J. Org. Chem.* **2002**, 3221–3236. (b) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. *Chem. Rev.* **2008**, *108*, 2796–2823. (c) Haerutyunyan, S. R.; den Hartog, T.; Geurts, K.; Minnaard, A. J.; Feringa, B. L. *Chem. Rev.* **2008**, *108*, 2824–2852.
- (8) For other selected asymmetric conjugate addition/enolate trapping methods, see: (a) Li, K.; Alexakis, A. *Tetrahedron Lett.* **2005**, *46*, 5823–5826. (b) Rathgeb, X.; March, S.; Alexakis, A. *J. Org. Chem.* **2006**, *71*, 5737–5742. (c) Germain, N.; Guéneé, L.; Mauduit, M.; Alexakis, A. *Org. Lett.* **2014**, *16*, 118–121. (d) Germain, N.; Schlaefli, D.; Chellat, M.; Rosset, S.; Alexakis, A. *Org. Lett.* **2014**, *16*, 2006–2009.
- (9) Vasu, D.; Hung, H.-H.; Bhunia, S.; Gawade, S. A.; Das, A.; Liu, R.-S. *Angew. Chem., Int. Ed.* **2011**, *50*, 6911–6914.
- (10) Witham, C. A.; Mauleón, P.; Shapiro, N. D.; Sherry, N. D.; Toste, F. D. *J. Am. Chem. Soc.* **2007**, *129*, 5838–5839.
- (11) (a) Qian, D.; Zhang, J. *Chem. Commun.* **2011**, *47*, 11152–11154. (b) Qian, D.; Hu, H.; Liu, F.; Tang, B.; Ye, W.; Wang, Y.; Zhang, J. *Angew. Chem., Int. Ed.* **2014**, *53*, 13751–13755.
- (12) Ji, K.; Zheng, Z.; Wang, Z.; Zhang, L. *Angew. Chem., Int. Ed.* **2014**, DOI: 10.1002/anie201409300.
- (13) See the Supporting Information for a comprehensive survey of conditions.
- (14) It is interesting to note that when 8-methylquinoline *N*-oxide was replaced by 3,5-dichloropyridine *N*-oxide, cyclopentadienyl aldehydes were obtained from 1,5-enynes: Hung, H.-H.; Liao, Y.-C.; Liu, R.-S. *J. Org. Chem.* **2013**, *78*, 7970–7976.
- (15) An analogous oxidation of a cyclopropyl gold(I) carbene was demonstrated to occur in the cyclization of 1,6-enynes with Ph₂SO as the oxidant.¹⁰
- (16) Yu, J.-Q.; Corey, E. J. *J. Am. Chem. Soc.* **2003**, *125*, 3232–3233.
- (17) CCDC 1037494 (**1**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- (18) (a) Jiménez-Núñez, E.; Echavarren, A. M. *Chem. Rev.* **2008**, *108*, 3326–3350. (b) Obradors, C.; Echavarren, A. M. *Acc. Chem. Res.* **2014**, *47*, 902–912.