

# Heck Coupling Using a Vinylido-MIDA Boronate: An Efficient and Modular Access to Polyene Frameworks

Johan Cornil,<sup>†</sup> Pierre-Georges Echeverria,<sup>‡</sup> Phannarath Phansavath,<sup>‡</sup> Virginie Ratovelomanana-Vidal,<sup>‡</sup> Amandine Guérinot,<sup>\*,†</sup> and Janine Cossy<sup>\*,†</sup>

<sup>†</sup>Laboratoire de Chimie Organique, Institute of Chemistry, Biology and Innovation (CBI)-UMR 8231, ESPCI ParisTech, CNRS, 10 rue Vauquelin 75231 Paris Cedex 05, France

<sup>‡</sup>Chimie ParisTech, CNRS, Institut de Recherche de Chimie Paris, PSL Research University, 11 rue Pierre et Marie Curie, 75005 Paris, France

## Supporting Information




**ABSTRACT:** A simple Heck coupling between an alkenyl iodo-boronate and a variety of terminal olefins is disclosed. This method gives access to a wide range of dienic moieties including valuable bis-functionalized dienes. The synthetic potential of the coupling reaction is demonstrated by a short and modular preparation of several tetraenic compounds.

Myriad natural products incorporating a conjugated polyenic motif have been isolated.<sup>1</sup> They generally display attractive properties, and some of them have been used as drugs, pigments, or fluorescent probes. Thus, the synthesis of polyenic moieties still represents a significant challenge for organic chemists. Traditional olefination methods such as the Wittig,<sup>2</sup> the Horner–Wadsworth–Emmons,<sup>3</sup> and the Julia<sup>4</sup> reactions have been widely applied to the preparation of conjugated polyenes, but the lack of stereocontrol as well as troublesome purifications remain a major obstacle to their use in total synthesis of complex molecules. Metal-catalyzed  $\text{sp}^2\text{--}\text{sp}^2$  cross-couplings overcome this difficulty, allowing the construction of polyenic fragments in a stereospecific fashion.<sup>5</sup> Hence, Suzuki, Negishi, Stille, or Heck cross-couplings recently emerged as privileged strategies for the assembly of polyenic moieties.<sup>6</sup> However, the preparation, the potential toxicity, as well as the instability of the alkenyl metal reagents prevent these reactions from becoming the universal method for the stereoselective construction of polyenes. Burke et al. partly solved this problem by focusing on *N*-methyliminoacetic acid (MIDA) boronates which have several advantages over boronic acids or other surrogates: they are bench-stable, compatible with silica gel chromatography, crystalline, inert toward a range of reaction conditions, and in addition, easy to cleave.<sup>7,8</sup> Thus, bis-functionalized alkenyl building-blocks incorporating both an halide atom (I, Br) and a MIDA boronate have been synthesized and successfully involved in several Pd-catalyzed cross-couplings including Suzuki, Stille, or Negishi reactions, allowing the modular synthesis of a large variety of polyenic natural products.<sup>9,10</sup> However, the preparation of alkenyl metal reagents is still necessary to perform such cross-couplings, and the Heck reaction, which requires simple olefins, could appear as a very attractive alternative.<sup>11</sup> Surprisingly, only one example of Heck

coupling involving the protected haloboronic acid **1a** and the reactive methyl acrylate has been reported.<sup>9a,c,12</sup> Herein, we disclose a Pd-catalyzed Heck reaction between the iodoalkenyl boronate **1b** and a broad range of olefins including functionalized alkenes (Scheme 1). The great potential of this method is highlighted by the short synthesis of several polyenic units.

## Scheme 1. Heck Reaction with Halogenoalkenyl Boronates



We first tried to generalize the reported Burke et al. conditions [ $\text{Pd(OAc)}_2$ ,  $\text{PPh}_3$ , and  $\text{Et}_3\text{N}$ ] for the Heck coupling between **1a** and several alkenes (Scheme 2). However, when the bromo boronate **1a** was treated with *N*-tosylallylamine, no reaction

## Scheme 2. Limitations of the Heck Coupling under Burke et al.'s Reported Conditions




Received: January 6, 2015

Published: February 2, 2015

occurred. In addition, when vinylpinacolborane was used as the olefinic partner, a complex mixture of products was formed.

In order to overcome these limitations, we switched to the more reactive iodoboronate **1b**, and after the reaction conditions were tuned, Pd(OAc)<sub>2</sub> (5 mol %) and AgOAc (1.5 equiv) were found to afford the coupling products with both ethyl acrylate and *N*-tosylallylamine in good yields (76% and 71%, respectively) in a regioselective fashion (100:0 and 93:7 linear/branched ratio) (Scheme 3).<sup>13</sup>

Scheme 3. Heck Coupling Involving Iodoboronate **1b**



With these conditions in hand, the scope and limitations of the Heck coupling involving iodo boronate **1b** were explored. Pleasingly, the reaction appeared quite general, and several functional groups such as an acetal were well tolerated when present in the olefinic partner (Table 1, entry 1). Allyl alcohol derivatives were also suitable under the reaction conditions, and

Table 1. Scope and Limitations of the Heck Coupling

entry	3	product	L/B or L/I <sup>[a]</sup>	yield <sup>[b]</sup>
1	<b>3c</b>		L/I = 85:15	75%
2	<b>3d</b>		L/B = 88:12	58%
3	<b>3e</b>		L/B = 87:13	63%
4	<b>3f</b>		L/B = 90:10	74%
5	<b>3g</b>		L/I = 85:15	65%
6	<b>3h</b>		-	0%
7	<b>3i</b> <sup>[c]</sup>		80:10:10 <sup>[d]</sup>	76%
8	<b>3j</b>		L/I = 95:5	77%
9	<b>3k</b>		L/I = 95:5	68%
10	<b>3l</b>		L/I = 95:5	75%
11	<b>3m</b>		L/I = 95:5	74%

<sup>a</sup>L/B: linear/branched. In some cases, the impurity (I) could not be identified undoubtedly as the branched isomer. The isomers were not separable. <sup>b</sup>The yield was calculated on the L/B or L/I mixture. <sup>c</sup>Purchased as a 95:5 mixture of (*E*)/(*Z*) isomers. <sup>d</sup>Mixture of isomers.

the protection of the hydroxyl group as a silyl ether proved to be beneficial (Table 1, entries 2–4). In all cases, a satisfactory regioselectivity of about 90:10 in favor of the linear isomer was observed. An enantiopure alcohol was successfully used to produce **3g** (65% yield, 85% purity) without epimerization of the stereocenters, thus highlighting the synthetic value of the method (Table 1, entry 5). Unfortunately, the coupling did not proceed with disubstituted olefins such as cyclohexenone (Table 1, entry 6). In contrast, a triene was formed when **1b** was reacted with methylpenta-2,4-dienoate **2i** (76%) albeit in a 80:10:10 mixture of isomers (Table 1, entry 7).<sup>14</sup> Styrenic partners exhibited good reactivity under the coupling conditions whatever the electronic nature of the substituents on the aryl moiety, and the expected products were isolated in good yields and high regioselectivities (Table 1, entries 8 and 9). Interestingly, the presence of heteroaromatics such as pyridine or indole was not detrimental to the reaction as **3l** and **3m** were isolated in good yields (Table 1, entries 10 and 11).

These promising results prompted us to examine the reactivity of functionalized alkene partners in order to access bifunctional dienes which are highly valuable building blocks in polyene synthesis.<sup>15,16</sup> Bis-functionalized Si–B dienes **5a** and **5b** were efficiently prepared from iodo-boronate **1b** (60% and 89% yield, respectively) (Table 2, entries 1 and 2). Vinyl and allyl phosphonates **4c** and **4d** were reacted with iodoboronate **1b** delivering the bifunctional dienes **5c** and **5d** in good yields (66% and 60%, respectively) (Table 2, entries 3 and 4). To the best of our knowledge, these dienes are unprecedented in the literature and could be involved subsequent transformations such as metal-catalyzed cross-couplings or HWE reaction. Gratifyingly, the bis-borylated diene **5e** was formed with an excellent yield of 90% (Table 2, entry 5). Thus, it is worth noting that this attractive building block was obtained in a one-step procedure from commercially available compounds, using a Heck coupling.<sup>17,18</sup>

Table 2. Access to Bifunctionalized Dienes


entry	5	product	L/B or L/I <sup>[a]</sup>	yield
1	<b>5a</b>		L/I = 95:5	60%
2	<b>5b</b>		L/B = 80:20	89%
3	<b>5c</b>		L/I = 95:5	66%
4	<b>5d</b>		L/B = 85:15	60%
5	<b>5e</b>		L/I = 90:10	90%

<sup>a</sup>L/B: linear/branched. In some cases, the impurity (I) could not be identified undoubtedly as the branched isomer.

An iodo desilylation was performed on **5b**, thus achieving a short synthesis of the dienic iodo boronate **6** which could be involved in an iterative Heck coupling (Scheme 4).<sup>19</sup> The synthetic utility of **5e** was well illustrated by performing two consecutive Suzuki cross-couplings that allowed the efficient and stereospecific formation of the (*E,E,E,Z*)-tetraene **8**.<sup>20</sup>


In order to demonstrate the high synthetic potential of the method, the MIDA boronate **3b** was transformed into the corresponding alkenyl iodide **9**, which was then submitted to a Suzuki cross-coupling with a range of MIDA boronates (Table

Scheme 4. Transformations of Dienes 5b and 5e



3). It is worth noting that these boronate partners were previously obtained by a Heck coupling. Only a few combinations among all possible are depicted here showing that this short sequence of reactions provides an easy and modular access to a large variety of polyenic frameworks.

Table 3. Synthesis of Tetraenes 10a–c




entry	product	purity	10 (yield)
1		92%	10a (74%)
2		84%	10b (77%)
3		93%	10c (75%)

<sup>a</sup>The isomeric purity was evaluated on the <sup>1</sup>H NMR spectrum after purification.

During the course of our studies toward the synthesis of the macrolactam mirabalin, we were interested in the preparation of tetraene 13 (Scheme 5).<sup>21</sup> Starting from the Heck product 3g, the targeted compound could be prepared in 3 steps with a satisfying global yield of 46%.<sup>22</sup> This straightforward synthesis advantageously competes with those already reported in the literature to access the same tetraenic motif.<sup>23,24</sup>

Scheme 5. Straightforward Synthesis of Tetraene 13



In conclusion, a simple, general and chemoselective Heck coupling involving an alkenyl iodoboronate is reported. A wide range of alkenes was compatible with the conditions, providing a library of dienic compounds. The stereoselective method gave a straightforward access to bis-functionalized building blocks, which could be easily transformed into polyenic frameworks. The use of simple olefins as the coupling partner makes the reaction easy-to-run even on highly advanced synthetic intermediates as no preparation of alkenyl metal is required. As such, this modular strategy should rapidly enter into the chemist toolbox for the construction of polyenic natural and non-natural products.

## ■ ASSOCIATED CONTENT

### Supporting Information

Experimental procedures and spectral data for all new compounds are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: [amandine.guerinot@espci.fr](mailto:amandine.guerinot@espci.fr).

\*E-mail: [janine.cossy@espci.fr](mailto:janine.cossy@espci.fr).

### Notes

The authors declare no competing financial interest.

## ■ REFERENCES

- (1) (b) Madden, K. S.; Mosa, F.; Whiting, A. *Org. Biomol. Chem.* **2014**, *12*, 7877. (a) Whiting, A.; Thirsk, C. *J. Chem. Soc., Perkin Trans. 1* **2002**, *8*, 999.
- (2) Maryanoff, B. E.; Reitz, B. *Chem. Rev.* **1989**, *89*, 863.
- (3) Wadsworth, W. S. *Org. React.* **1977**, *25*, 73.
- (4) (a) Blakemore, P. R. *J. Chem. Soc., Perkin Trans. 1* **2002**, 2563. (b) Aissa, C. *Eur. J. Org. Chem.* **2009**, 1831. (c) Markó, I. E.; Pospíšil, J. In *Science of Synthesis, Vol 47a: Alkenes*; de Meijere, A., Ed.; George Thieme Verlag: Stuttgart, 2009, 105.
- (5) (a) Diederich, F.; Stang, P. *Metal-Catalyzed Cross-Coupling Reactions*; Wiley-VCH, Weinheim, 1998. (b) Negishi, E. I. *Handbook of Organopalladium Chemistry for Organic Synthesis*; John Wiley & Sons: New York, 2002; Vol. 1. (c) *Metal-Catalyzed Cross-Coupling Reactions and More*; de Meijere, A., Bräse, S., Oestreich, M., Eds.; Wiley-VCH: Weinheim, 2014. (d) Carboni, B.; Carreaux, F.; Molander, G. A.; Wolfe, J. P.; Larhed, M. *Science of Synthesis, Cross-Coupling and Heck-Type Reactions*, 1 George Thieme Verlag: Stuttgart, 2013; pp 265–321. (e) Negishi, E. I.; Wang, G.; Rao, H.; Xu, Z. *J. Org. Chem.* **2010**, *75*, 3151.
- (6) Cross-metathesis has been used for diene synthesis; see: Wojtkielewicz, A. *Curr. Org. Synth.* **2013**, *10*, 43.
- (7) (a) Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2007**, *129*, 6716. (b) Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2008**, *130*, 14084. (c) Knapp, D. M.; Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2009**, *131*, 6961. (d) Close, A. J.; Kemmitt, P.; Emmerson, M. K.; Spencer, J. *Tetrahedron* **2014**, *70*, 9125.
- (8) For other surrogates, see: (a) Molander, G. A.; Felix, L. A. *J. Org. Chem.* **2005**, *70*, 3950. (b) Molander, G. A.; Sandrock, D. L. *J. Am. Chem. Soc.* **2008**, *130*, 15792. (c) Nogushi, H.; Hojo, K.; Suginoe, M. *J. Am. Chem. Soc.* **2007**, *129*, 758. (d) Nogushi, H.; Shioda, T.; Chou, C.-M.; Suginoe, M. *Org. Lett.* **2008**, *10*, 377. (e) Li, J.; Burke, M. D. *J. Am. Chem. Soc.* **2011**, *133*, 13774.
- (9) (a) Lee, S. J.; Gray, K. C.; Paek, J. S.; Burke, M. D. *J. Am. Chem. Soc.* **2008**, *130*, 466. (b) Lee, S. J.; Anderson, T. M.; Burke, M. D. *Angew. Chem., Int. Ed.* **2010**, *49*, 8860. (c) Woerly, E. M.; Struble, J. R.; Palyam, N.; O'Hara, S. P.; Burke, M. D. *Tetrahedron* **2011**, *67*, 4333. (d) Woerly, E. M.; Miller, J. E.; Burke, M. D. *Tetrahedron* **2013**, *69*, 7732. (e) Woerly, E. M.; Roy, J.; Burke, M. D. *Nature Chem.* **2014**, *6*, 484. (f) Weber, A.; Dehn, R.; Schläger, N.; Dieter, B.; Kirschning, A. *Org. Lett.* **2014**, *16*, 568. (g) McLaughlin, M. G.; McAdam, C. A.; Cook, M. J. *Org. Lett.* **2015**, *17*, 10.
- (10) For recent reviews on iterative cross-coupling, see: (a) Wang, C.; Glorius, F. *Angew. Chem., Int. Ed.* **2009**, *48*, 5240. (b) Tobisu, M.; Chatani, N. *Angew. Chem., Int. Ed.* **2009**, *48*, 3565.
- (11) For reviews on Heck reaction, see: (a) Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009. (b) Alonso, F.; Beletskaya, I. P.; Yus, M. *Tetrahedron* **2005**, *61*, 11771.
- (12) Alternatively, Heck coupling reactions involving vinyl boronates have been reported for the construction of dienes. For selected examples, see: (a) Stewart, S. K.; Whiting, A. *Tetrahedron Lett.* **1995**, *36*, 3925. (b) Hénaff, N.; Whiting, A. *Org. Lett.* **1999**, *1*, 1137. (c) Hénaff, N.; Whiting, A. *Tetrahedron* **2000**, *56*, 5193. (d) Lightfoot, A. P.; Maw, G.; Thirsk, C.; Twiddle, S. J. R.; Whiting, A. *Tetrahedron Lett.* **2003**, *44*,



7645. (e) Lightfoot, A. P.; Twiddle, S. J. R.; Whiting, A. *Org. Biomol. Chem.* **2005**, *3*, 3167. (f) Batsanov, A. S.; Knowles, J. P.; Whiting, A. *J. Org. Chem.* **2007**, *72*, 2525. (g) Sakai, M.; Guan, T.; Hara, S. *J. Fluor. Chem.* **2007**, *128*, 1444. (h) Batsanov, A. S.; Knowles, J. P.; Samsam, B.; Whiting, A. *Adv. Synth. Catal.* **2008**, *350*, 227. (i) Uno, B. E.; Gillis, E. P.; Burke, M. D. *Tetrahedron* **2009**, *65*, 3130. (j) Delcamp, J. H.; Gormisky, P. E.; White, M. C. *J. Am. Chem. Soc.* **2013**, *135*, 8460. (k) Szudkowska-Frątczak, J.; Ryba, A.; Franczyk, A.; Walkowiak, J.; Kubicki, M.; Pawluć, P. *Appl. Organomet. Chem.* **2014**, *28*, 137.

(13) See the Supporting Information for details.

(14) The (*E,E*)-diene was the major isomer, and the other isomers could not be identified without a doubt.

(15) For other syntheses of similar bis-functionalized dienes, see: (a) Badubri, F.; Farinola, G. M.; Flandanese, V.; Mazzone, L.; Naso, F. *Tetrahedron* **1998**, *54*, 1085. (b) Walkowiak, J.; Jankowska-Wadja, M.; Marciniak, B. *Chem.—Eur. J.* **2008**, *14*, 6679. (c) Maxal, C.; Vaultier, M. *Tetrahedron Lett.* **1994**, *35*, 3089. (d) Daini, M.; Suginome, M. *Chem. Commun.* **2008**, 5224. (e) Suginome, M.; Matsuda, T.; Ito, Y. *Organometallics* **1998**, *17*, 5233. (f) Fujii, S.; Chang, S. Y.; Burke, M. D. *Angew. Chem., Int. Ed.* **2011**, *50*, 7862. (g) Desurmont, G.; Klein, R.; Uhlenbrock, S.; Laloë, E.; Deloux, L.; Giolando, D. M.; Kim, Y. W.; Pereira, S.; Srebnik, M. *Organometallics* **1996**, *15*, 3323. (h) Albrecht, K.; Kaufmann, D. E. In *Science of Synthesis*; George Thieme Verlag: Stuttgart, 2005; Vol. 6, p 697.

(16) For selected examples of other bis-functionalized dienes and trienes, see: (a) Lipshutz, B. H.; Lindsley, C. *J. Am. Chem. Soc.* **1997**, *119*, 4555. (b) Denmark, S. E.; Tymonko, S. A. *J. Am. Chem. Soc.* **2005**, *127*, 8004. (c) Coleman, R. S.; Lu, X.; Modolo, I. *J. Am. Chem. Soc.* **2007**, *129*, 3826. (d) Coleman, R. S.; Walczak, M. C. *Org. Lett.* **2005**, *7*, 2289. (e) Badubri, F.; Fiandanese, V.; Marchese, G.; Naso, F. *J. Chem. Soc., Chem. Commun.* **1991**, 237. (f) Badubri, F.; Fiandanese, V.; Naso, F.; Punzi, A. *Tetrahedron Lett.* **1994**, *35*, 2067. (g) Singidi, R. R.; RajanBabu, T. V. *Org. Lett.* **2010**, *12*, 2622.

(17) As a comparison, Burke et al. prepared the same building block in five steps; see ref 15f.

(18) The Heck cross-coupling was performed on gram scale, and a batch of 2 g of **5e** was prepared without erosion of the yield.

(19) For other methods to access dienic halogeno-boronates, see: (a) Woerly, E. M.; Cherney, A. H.; Davis, E. K.; Burke, M. D. *J. Am. Chem. Soc.* **2010**, *132*, 6941. (b) Singidi, R. R.; Kutney, A. M.; Galluci, J. C.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2010**, *132*, 13078. (c) Gillis, E. P.; Lee, S. J.; Gray, K.; Burke, M. D.; Knapp, D. M. WO US84156, 2007. (d) Sakai, M.; Guan, T.; Hara, S. *J. Fluorine Chem.* **2007**, *178*, 1444.

(20) Knapp, D. M.; Gillis, E. P.; Burke, M. D. *J. Am. Chem. Soc.* **2009**, *131*, 6961.

(21) (a) Plaza, A.; Baker, H. L.; Bewley, C. A. *J. Nat. Prod.* **2008**, *71*, 473. (b) Plaza, A.; Baker, H. L.; Bewley, C. A. *J. Nat. Prod.* **2009**, *72*, 324 (Correction). (c) Echeverria, P.-G.; Prévost, S.; Cornil, J.; Férard, C.; Raymond, S.; Guérinot, A.; Cossy, J.; Ratovelomanana-Vidal, V.; Phansavath, P. *Org. Lett.* **2014**, *16*, 2390.

(22) In **12**, the major isomer was the (*E,E,E,E*)-tetraene. After a purification by flash chromatography on silica gel, **13** was isolated as a single (*E,E,E,E*)-isomer.

(23) (a) Denmark, S. E.; Fujimori, S. *J. Am. Chem. Soc.* **2005**, *127*, 8971. (b) Mori, Y.; Assai, M.; Kawade, J.-I.; Furukawa, H. *Tetrahedron* **1995**, *51*, 5315. (c) Amans, D.; Bellostta, V.; Dacquet, C.; Ktorza, A.; Hennuyer, N.; Staels, B.; Caignard, D.-H.; Cossy, J. *Org. Biomol. Chem.* **2012**, *10*, 6169. (d) Evans, D. A.; Connell, B. T. *J. Am. Chem. Soc.* **2003**, *125*, 10899.

(24) A 200 mg portion of the tetraenic fragment **13** could be obtained through this sequence of reactions.