


First Total Synthesis of Pandamarine

Dimitris Kalaitzakis, Dimitris Noutsias, and Georgios Vassilikogiannakis*

Department of Chemistry, University of Crete, Vasilika Vouton, 71003 Iraklion, Crete, Greece


S Supporting Information



ABSTRACT: The first total synthesis of pandamarine, an alkaloid isolated from *Pandanus amaryllifolius* is reported. The key step of this extremely short (six steps in total) and protecting group-free synthesis is a highly efficient cascade reaction sequence initiated by the photooxidation of an easily accessible and symmetric difuran precursor.

Pandamarine (**1**, Scheme 1) is an alkaloid that was isolated in 1992 by Byrne's group from the leaves of the tropical plant *Pandanus amaryllifolius*.¹ Pandamarilactone-1 (**2**, Scheme 1), was also isolated from the same plant one year later.² Biogenetically, the isolation team have proposed that pandamarine (**1**) might be the spiro-*N,N*-ketalization product of a symmetric *bis*-lactam (**3**, Scheme 1).^{1,3} This proposal is

Scheme 1. Proposed Biosynthetic Origin of Pandamarine^{1,3}



consistent with the fact that pandamarine (**1**) occurs as a racemate. It was suggested that this possible biogenetic intermediate **3** might in turn be obtained from (2S)-4-hydroxy-4-methylglutamic acid (**4**, Scheme 1) since this amino acid was found in a related species, *P. veitchii*.⁴ *Pandanus amaryllifolius* is one species out of 600–700 *Pandanus* trees, or shrub-like plants, that are widely distributed throughout the tropics. Many medicinal applications for extracts from the genus *Pandanus* have been reported.^{5,6} Among these reports, it

was recently shown that a *Pandanus amaryllifolius* leaf extract has antihyperglycemic effects.⁶


Recently, Robertson's group reported the first total synthesis of pandamarilactone-1 (**2**).⁷ Within the same study, it was also proven that pandamarine (**1**) could not be produced by subjecting pandamarilactone-1 (**2**) to acid and then base sequentially (H₂SO₄/NH₃), under conditions that mimicked those used in the isolation in order to separate the alkaloid from nonalkaloidal constituents. Therefore, it was proposed that pandamarine (**1**) is a genuine natural product rather than an artifact of the isolation process.

Pandamarine's structure was determined by X-ray diffraction¹ and consists of both a diazaspino[4.5] and a 5-ylidenepyrrol-2(*SH*)-one unit, connected by a carbon chain (C7–C9, **1**, Scheme 1). The diazaspino[4.5] motif is found in a small number of other natural products such as haplophytine,⁸ isoschizogamine,⁹ and the related alkaloids leuconoxine,¹⁰ the leuconodines,¹¹ and melodinine E.¹² In contrast, the 5-ylidenepyrrol-2(*SH*)-one unit is ubiquitous appearing in many natural products and pharmaceuticals.^{13,14} Unsurprisingly, reports dealing with the synthesis of diazaspino[4.5] scaffolds are limited,¹⁵ while many methods have been developed for the synthesis of 5-ylidenepyrrol-2(*SH*)-one motifs.¹³

We recently developed a methodology for the synthesis of 5-hydroxy-1*H*-pyrrol-2(*SH*)-ones of type **C** and their dehydrated counterparts 5-ylidenepyrrol-2(*SH*)-ones **D** (Scheme 2) starting from furans and using a variety of primary amines, or ammonia, as the nitrogen source.¹⁶ More specifically, a complex cascade reaction sequence begins with the oxidation of substituted furans by singlet oxygen, generated using methylene blue (MB), or rose bengal (RB), and visible light. The initial products of this sequence are 2-pyrrolidinones of type **B** (Scheme 2).^{17,18} The reaction sequence can easily be extended to the synthesis of α,β -unsaturated γ -hydroxy lactams of type **C** through the catalytic oxidation of the 2-pyrrolidinones **B** by MB


Received: June 10, 2015

Published: June 30, 2015

Scheme 2. Synthesis of 5-Ylidenepyrrrol-2(*SH*)-ones **D** from Furans


in the presence of molecular dioxygen.¹⁶ We found that this second oxidation step is accelerated by the basic environment of the reaction mixture at this stage. The dehydrated counterparts (**D**, Scheme 2) can be the final product of the sequence if next a Brønsted acid (HCOOH, TFA, or PTSA) is used.¹⁶ Now, we wanted to investigate if this methodology could be applied to a difuran precursor (**7**, Scheme 3) and in the presence of an unprotected secondary amine appended as part of the alkyl side chain.

Scheme 3. Retrosynthetic Analysis




Our retrosynthetic analysis for pandamarine (**1**) is presented in Scheme 3. Pandamarine **1** might be derived by a ketalization/dehydration reaction of the symmetrical *bis*-5-hydroxy-1*H*-pyrrol-2(*SH*)-one **5** which it was hoped might be obtained as the MB-catalyzed oxidation product of the intermediate *bis*-2-pyrrolidinone **6**.¹⁶ This pyrrolidinone might in turn be accessed via photooxygenation of the symmetric difuran **7** using ammonia as a nitrogen source.^{16,18}

The synthesis of photooxygenation substrate **7** starting from the commercially available 5-hexynitrile (**8**) is described in Scheme 4. The alkynyl lithium anion of **8** was added to

Scheme 4. Synthesis of Difuran **7**

unprotected hydroxyacetone to afford diol **9**. In the next couple of steps, we proceeded as Robertson⁷ had in his synthesis of pandamarilactone-1 (**2**) starting initially with the cyclodehydration of alkynyl diol **9** by silver nitrate to furnish furan **10**. However, we performed this reaction in methanol instead of CH_2Cl_2 because of the low solubility of diol **9** in CH_2Cl_2 . Nitrile **10** was separated into two portions; the first was reduced to the corresponding aldehyde **11** and the second to amine **12**. Difuran **7** was then obtained via the reductive amination of aldehyde **11** using sodium borohydride and amine **12**.


The photooxygenation reaction of difuran **7**, which, it should be noted, contains an unprotected amine, was performed in methanol (0.3 mmol, final concentration 40 mM) using MB (0.1 mM, 0.25 mol % compared to the furan substrate) as the photosensitizer (Scheme 5). After irradiation of the solution with visible light in the presence of molecular oxygen (bubbling through the solution) the double oxidation of the substrate was complete in just 2.5 min (the reaction was monitored by 1H NMR). More MB (6 mol % final ratio, 2.4 mM final

Scheme 5. Synthesis of Pandamarine from Difuran **7**

concentration) was then added, and the photooxidized product was reduced with Me₂S (7 equiv, 75 min). Intriguingly, the subsequent addition of aqueous ammonia (4 equiv) afforded within 5 h intermediate **5**, as evidenced by the crude ¹H NMR spectrum. The combination of MB (6 mol %) with ammonia (4 equiv) was proven to be very effective for the formation of the intermediate *bis*-2-pyrrolidinone **6** and its subsequent one-pot oxidation to the desired *bis*-5-hydroxy-1*H*-pyrrol-2(*5H*)-one **5** (Scheme 5). The final step of the reaction sequence was to effect the ketalization/dehydration of **5**. This transformation turned out to be quite tricky as the direct addition of TFA in CHCl₃ did not yield pandamarine (**1**). However, the complete removal of the volatile contaminants from the reaction mixture before the addition of TFA was the key to success. More precisely, after the bulk of the solvent had been removed using a rotary evaporator, the residue was subjected to high vacuum at 50 °C for 5 h prior to the addition of the acid. To our delight, in this case, the addition of TFA (2 equiv) to a solution of **5** in CHCl₃ containing molecular sieves (4 Å) afforded pandamarine (30% overall isolated yield starting from **7**, Scheme 5) in 2 h. The ¹H and ¹³C NMR data are in full agreement with those reported in the isolation paper.¹

To find out if lactam **3** was indeed the precursor of the natural product (**1**), under the conditions we had employed, compound **3** was prepared independently starting from the Boc-protected di(furylalkyl)amine **13** (Scheme 6). Subsequent

Scheme 6. Synthesis of *bis*-Lactam **3**



application of the early stages of the protocol we had developed, led to the transformation of **13** into the Boc-protected *bis*-5-hydroxy-1*H*-pyrrol-2(*5H*)-one **14** which was then dehydrated and deprotected to furnish the desired *bis*-lactam **3** when the reaction mixture was treated with HCO₂H at 40 °C (41% overall isolated yield starting from **13**). The *Z,Z* geometry of **3** was proven by NOE studies. Treatment of **3** under the conditions applied for the transformation of **5** to **1** (Scheme 5), or under even harsher conditions (6 equiv TFA, 18 h) did not give the natural product (recovery of starting

material). This experiment would suggest that under our conditions, the spiro-*N,N*-ketalization of **5** precedes the dehydration. In other words, **3** is not an intermediate in the transformation of **5** to **1**.

In conclusion, we have presented the first total synthesis of pandamarine (**1**), as well as, its proposed biogenetic intermediate (**3**). The developed synthetic strategy is very practical and extremely fast (just six steps from **8** to **1**). No protecting groups are used in the pandamarine (**1**) synthesis, oxygen is the terminal oxidant and the step-economy is very good. The key step of this approach is a cascade reaction sequence initiated by singlet oxygen which leads to the transformation of a very simple linear difuran precursor **7** into the natural product.

■ ASSOCIATED CONTENT

§ Supporting Information

Experimental procedures, full spectroscopic data, and copies of ¹H and ¹³C NMR spectra for all new compounds. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b01693.

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: vasil@chemistry.uoc.gr.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 277588.

■ REFERENCES

- (1) Byrne, L. T.; Guevara, B. Q.; Patalinghug, W. C.; Regio, B. V.; Ualat, C. R.; White, A. H. *Aust. J. Chem.* **1992**, *45*, 1903.
- (2) Nonato, M. G.; Garson, M. J.; Truscott, R. J. W.; Carver, J. A. *Phytochemistry* **1993**, *34*, 1159.
- (3) Nonato, M. G.; Takayama, H.; Garson, M. J. *Alkaloids: Chemistry and Biology*; Elsevier: 2008; Vol. 66, p 215.
- (4) Bell, E. A.; Meier, L. K.; Sorensen, H. *Phytochemistry* **1981**, *20*, 2213.
- (5) (a) Santos, A. C.; Santos, G. A.; Obligacion, M. B. S.; Olay, I. P.; Fojas, F. R. *Philippine Plants and their contained Natural Products and Pharmaceutical literature survey*; National Research Council of the Philippines: Bicutan, Taguig, Metro Manila, 1981. (b) Tan, M. *Philippine Medicinal Plants in Common Use: Their Phytochemistry and Pharmacology*; AKAP: Quezon City, Philippines, 1980.
- (6) Chiabchalar, A.; Nooron, N. *Pharmacogn. Mag.* **2015**, *11*, 117.
- (7) Seah, K. Y.; Macnaughton, S. J.; Dallimore, J. W. P.; Robertson, J. *Org. Lett.* **2014**, *16*, 884.
- (8) (a) Rogers, E. F.; Snyder, H. R.; Fischer, R. F. *J. Am. Chem. Soc.* **1952**, *74*, 1987. (b) Snyder, H. R.; Fischer, R. F.; Walker, J. F.; Els, H. E.; Nussberger, G. A. *J. Am. Chem. Soc.* **1954**, *76*, 2819. (c) Snyder, H. R.; Fischer, R. F.; Walker, J. F.; Els, H. E.; Nussberger, G. A. *J. Am. Chem. Soc.* **1954**, *76*, 4601. (d) Snyder, H. R.; Strohmayer, H. F.; Mooney, R. A. *J. Am. Chem. Soc.* **1958**, *80*, 3708.
- (9) (a) Renner, U.; Kernweisz, P. *Experientia* **1963**, *19*, 244. (b) Renner, U. *Lloydia* **1964**, *27*, 406. (c) Hájiček, J.; Taimr, J.; Buděšínský, M. *Tetrahedron Lett.* **1998**, *39*, 505.
- (10) Abe, F.; Yamauchi, T. *Phytochemistry* **1994**, *35*, 169.
- (11) Gan, C.-Y.; Low, Y.-Y.; Thomas, N. F.; Kam, T.-S. *J. Nat. Prod.* **2013**, *76*, 957.

(12) Feng, T.; Cai, X.-H.; Liu, Y.-P.; Li, Y.; Wang, Y.-Y.; Luo, X.-D. *J. Nat. Prod.* **2010**, *73*, 22.

(13) Nay, B.; Riache, N.; Evanno, L. *Nat. Prod. Rep.* **2009**, *26*, 1044.

(14) For recent examples, see: (a) Chatzimpaloglou, A.; Kolosov, M.; Eckols, T. K.; Tweardy, D. J.; Sarli, V. *J. Org. Chem.* **2014**, *79*, 4043. (b) Kumar, M. M. K.; Naik, J. D.; Satyavathi, K.; Ramana, H.; Varma, P. R.; Nagasree, K. P.; Smitha, D.; Rao, D. V. *Nat. Prod. Res.* **2014**, *28*, 888. (c) Miyazaki, H.; Miyake, T.; Terakawa, Y.; Ohmizu, H.; Ogiku, T.; Ohtani, A. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 546.

(15) (a) Büchel, K. H.; Bocz, A. K.; Korte, F. *Chem. Ber.* **1966**, *99*, 724. (b) Tsuge, O.; Watanabe, H.; Masuda, K.; Yousif, M. M. *J. Org. Chem.* **1979**, *44*, 4543. (c) Martín-López, M. J.; Bermejo, F. *Tetrahedron* **1998**, *54*, 12379. (d) Matsumoto, K.; Tokuyama, H.; Fukuyama, T. *Synlett* **2007**, 3137. (e) Kalaitzakis, D.; Antonatou, E.; Vassilikogiannakis, G. *Chem. Commun.* **2014**, *50*, 400.

(16) Kalaitzakis, D.; Kouridaki, A.; Noutsias, D.; Montagnon, T.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2015**, *54*, 6283.

(17) For a recent review concerning the photooxygenation of furans, see: Montagnon, T.; Kalaitzakis, D.; Triantafyllakis, M.; Stratakis, M.; Vassilikogiannakis, G. *Chem. Commun.* **2014**, *50*, 15480.

(18) (a) Kalaitzakis, D.; Montagnon, T.; Alexopoulou, I.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2012**, *51*, 8868. (b) Kalaitzakis, D.; Montagnon, T.; Antonatou, E.; Bardaji, N.; Vassilikogiannakis, G. *Chem. - Eur. J.* **2013**, *19*, 10119. (c) Kalaitzakis, D.; Montagnon, T.; Antonatou, E.; Vassilikogiannakis, G. *Org. Lett.* **2013**, *15*, 3714.