


Base-Promoted N-Pyridylation of Heteroarenes Using *N*-Propargyl Enaminones as Equivalents of Pyridine Scaffolds

Guolin Cheng,* Yunxiang Weng, Xifa Yang, and Xiuling Cui*

Engineering Research Center of Molecular Medicine, Ministry of Education, Key Laboratory of Xiamen Marine and Gene Drugs, Institutes of Molecular Medicine and School of Biomedical Sciences, Huaqiao University, Xiamen 361021, China

S Supporting Information

ABSTRACT: N-Pyridylation of various N-heteroarenes, including N-heteroarene-containing peptides, was achieved using *N*-propargyl enaminones (isolated or generated in situ from propargyl amine and propynones) as masked polysubstituted pyridine cores. This metal-free procedure proceeds under mild reaction conditions and generates 1 equiv of H₂O as the sole byproduct.




Pyridines represent one of the most important classes of heterocyclic compounds.¹ Among them, heteroaryl-containing pyridines have attracted much research interest due to their prevalence in agrochemicals,² pharmaceuticals,³ and advanced materials.⁴ Over the past decade, various synthetic methods,⁵ including transition-metal-catalyzed reactions,⁶ organocatalyzed reactions,⁷ and the metal-free reactions,⁸ have been developed to selectively synthesize highly functionalized pyridine derivatives. Despite these advances, the Ullman-type *N*-pyridylation of heteroarenes is still the most commonly used method for constructing *N*-pyridyl heteroarenes, which requires functionalized pyridine cores as the starting materials.⁹ Novel approaches to these motifs are rare and remain to be explored.¹⁰

In organic synthesis, the cascade reactions are used to reduce waste, avoid isolating intermediates, and improve the reaction efficiency. We recently have described a cascade reaction for the synthesis of 1,5-disubstituted 1,2,3-triazoles from enaminones.^{11a} In the exploration of this reaction, an unexpected seven-membered ring 1,4-oxazepine **2** was observed from *N*-propargyl enaminones **1**.¹² It was considered that an equilibrium might exist between 1,4-oxazepines **2** and the epoxide intermediates **B** through a 6 π -electrocyclization/walk rearrangement cascade reaction.¹³ Intermediate **B** is proposed to be trapped by a suitable nucleophile to form 2,3-dihydropyridines **4**, which would be further aromatized to give substituted pyridines **5** (Scheme 1).

In our initial survey, indole **3a** and 1,3-diphenyl-3-(propargylamino)-2-propen-1-one **1a** were used as substrates to verify the above hypothesis (Table 1). To our delight, the desired product **5aa** was isolated in 35% yield when the mixture of **3a** and **1a** in DMSO was treated with LiOtBu for 5 min (entry 1).¹⁴ Its molecular structure was confirmed by single crystal X-ray diffraction. Acceptable yields were achieved in the presence of other strong bases, such as NaOtBu, KOtBu, NaOEt, KOH, and NaOH (entries 2–6). NaOH gave the best result, affording **5aa** in 90% yield (entry 6). In contrast, no reactions occurred when weak bases, such as K₂CO₃ or Et₃N, were used (entries 7, 8). Further investigation showed that

Scheme 1. A Proposed Route to Pyridine Derivatives from *N*-Propargyl Enaminones




polar aprotic solvents favored this transformation (entries 9–13). However, no better result was obtained than DMSO. Remarkably, a respectable 76% yield of **5aa** was achieved using 0.3 equiv of NaOH (entry 14).

With the optimal reaction conditions established, we started our investigation by reacting enaminone **1a** with different indoles (Scheme 2). A diverse set of functional groups, such as alkoxy, halogen, cyano, formyl, and nitro, were tolerated in this procedure owing to the mild reaction conditions. Electron-donating groups in the indole ring **3ab**, **3ac** led to higher yields than stronger electron-withdrawing groups **3ah–aj**, probably because electron-donating substituents enhanced the nucleophilicity of N atom of indole. It was obviously observed that the steric hindrance on the C2- or C7-position of indole lowered the nucleophilicity of the N atom and affected the reaction efficiency **3al–ao**. Notably, double *N*-pyridylation of bis-(indole) could afford the desired product **5ap** in 55% yield. This heteroaryl-containing pyridine with C₂ symmetry is expected to be developed as a new functional material or ligand in an organic reaction. Next, we focused our attention on

Received: June 19, 2015

Published: July 22, 2015


Table 1. Optimizing the Reaction Conditions for the Cascade Reaction of Indole 3a with 1,3-Diphenyl-3-(propargylamino)-2-propen-1-one 1a^a



| entry | solvent | base | time (min) | yield (%) ^b |
|-----------------|---------|--------------------------------|------------|------------------------|
| 1 | DMSO | LiOtBu | 5 | 35 |
| 2 | DMSO | NaOtBu | 5 | 86 |
| 3 | DMSO | KOtBu | 5 | 78 |
| 4 | DMSO | NaOEt | 60 | 60 |
| 5 | DMSO | KOH | 5 | 87 |
| 6 | DMSO | NaOH | 5 | 90 |
| 7 | DMSO | K ₂ CO ₃ | 60 | 0 |
| 8 | DMSO | Et ₃ N | 60 | 0 |
| 9 | DMF | NaOH | 5 | 88 |
| 10 | NMP | NaOH | 5 | 89 |
| 11 | MeOH | NaOH | 60 | 0 |
| 12 | THF | NaOH | 60 | 0 |
| 13 | toluene | NaOH | 60 | 0 |
| 14 ^c | DMSO | NaOH | 60 | 76 |

^aReaction conditions: 3a (0.5 mmol), 1a (0.5 mmol), base (0.5 mmol), in 1 mL solvent, rt. ^bIsolated yields. ^c0.15 mmol base. DMSO = dimethyl sulfoxide. DMF = *N,N*-dimethylformamide. NMP = *N*-methyl-2-pyrrolidone. THF = tetrahydrofuran.

Scheme 2. Scope of N-Nucleophiles^a




^aReaction conditions: 3a (0.5 mmol), 1a (0.5 mmol), NaOH (0.5 mmol), DMSO (1 mL), rt.

investigating the scope of other N-heteroarenes. Delightfully, various N-heteroarenes, including carbazole, pyrrole, imidazole, and pyrazole, furnished the corresponding products 5aq-at in good yields. Benzimidazole and 7-azaindole were less reactive and gave the corresponding products in lower yields. Moreover, indazole could be employed in this process with good efficiency. A mixture of 1*H*-indazole 5aw and 2*H*-indazole 5aw' was generated in 85% total yield. In all cases, these rapid reactions were completed within 30 min. Furthermore, other nucleophiles, such as benzylamine, aniline, acetanilide, succinimide, and benzyl thiol, were also tested. No desired products were observed.

Our previous works have indicated that *N*-propargyl enamines 1 could be easily prepared from propargyl amine 6 and propynones 7.¹¹ Thus, a "one pot" two-step strategy to construct 2-(1-indolyl)pyridines 5ba-bx starting from 6 and 7a-x was examined (Scheme 3). It was worth noticing that the

Scheme 3. Scope of Propynones^a




^aReaction conditions: 6 (0.55 mmol), 7 (0.5 mmol), DMSO (1 mL), 60 °C. After the reaction mixture was cooled to rt, indole (0.5 mmol) and NaOH (0.5 mmol) were added. Isolated yields based on 7.



reaction was equally successful with both isolated and *in situ* generated 1. As illustrated in Scheme 3, the reaction was not significantly affected by the electronic effects of the substituents. Propynones 7 bearing electron-rich or electron-deficient groups on R¹- or R²- were both effective. When sterically hindered propynones were used as substrates, the corresponding products 5bh-bk, 5bx were furnished in lower yields. Remarkably, 1-naphthalenyl and 2-thienyl propynones were suitable substrates, affording 5bl, 5bm in 71% and 70% yields, respectively. However, alkyl groups on R¹- affected the reaction efficiency. No desired product was formed when R²-

was an alkyl group. 3-Phenylprop-2-yn-1-amine **8** was also used instead of **6** to further explore the substrate scope; however, the desired product was not formed.

The Ullman-type N-pyridylation of heteroarenes is often carried out at elevated temperatures (100–190 °C), in the presence of copper salts and bases.⁹ The harsh reaction conditions of Ullman reactions limited its application in the pharmaceutical industry and life science because of the metal contamination and the racemization of chiral compounds. These challenges prompted us to explore our procedure in the modification of complex molecules, especially peptides. As expected, L-tryptophane-containing dipeptide **3x** could be modified, affording **5ax** in 82% yield as a single diastereomer (eq 1).




To understand the mechanism of this cascade reaction, some experiments were carried out. When the standard reaction of **1a** with indole was quenched with water after 2 min, the 1,4-oxazepine intermediate **2a** was isolated in 35% yield, which could be further transformed into the desired product **5aa** in nearly quantitative yield (eq 2). As expected, **2a** and **2b** could



be obtained from **1a** and **1b**, respectively, under standard conditions in the absence of indole (eq 3).¹⁴ The tentative 2,3-dihydropyridine intermediate **4** was not observed during the reaction, presumably because the dehydrative aromatization rate of **4** is faster than the formation rate of **4** under basic conditions. Thus, various N-heteroarenes were tested to generate intermediate **4** in the absence of a base. To our delight, when pyrazole was used as a nucleophile, the desired intermediate **4t** was formed in 67% yield. The configurations of **4t** were assigned based on the 2D-NOESY NMR spectroscopy, where the pyrazole group is *trans* to the hydroxyl group. In this case two adjacent stereocenters were created simultaneously and no diastereoisomer was formed. Finally, **5at** could be generated in almost quantitative yield from **4t** under the standard conditions (eq 4).

Based on the experimental results obtained, the reaction mechanism was proposed. As shown in Scheme 4, an initial propargyl–allenyl isomerization/enolization cascade reaction of enaminones **1** formed iminoenolate intermediates **C**.¹⁵ The intramolecular 7-*exo-dig* cyclization of **C** provided 1,4-

Scheme 4. Proposed Reaction Mechanism



oxazepines **2**, which subsequently isomerized to give epoxide intermediates **B** via a 6 π -electrocyclization/walk rearrangement cascade reaction.¹⁶ Epoxide ring opening of **B** with N-heteroarenes proceeds via S_N2 substitution, generating *trans*-2,3-dihydropyridine intermediates **4**. At last, the dehydrative aromatization of **4** leads to the final products **5**.

In conclusion, we have described a general and operationally simple method to construct diverse 2-(1-heteroaryl) pyridines from the reaction of N-propargyl enaminones with N-heteroarenes. This procedure allowed the formation of one pyridine core and one C–N bond in “one pot”, and it showed a wide range of functional group tolerance. This protocol only required 1 equiv of NaOH as an additive, and it generated 1 equiv of H₂O as the sole byproduct, thereby making this process environmentally friendly and atom economic. Moreover, this new process could be applied to modify heteroarene-containing complex molecules, such as peptides.

■ ASSOCIATED CONTENT

Supporting Information

Copies of ¹H NMR and ¹³C NMR for all synthesized compounds. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b01733.

■ AUTHOR INFORMATION

Corresponding Authors

*E-mail: glcheng@hqu.edu.cn.

*E-mail: cuixl@hqu.edu.cn.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the NSF of China (21202048), Program for Minjiang Scholar (10BS216), Xiamen Southern Oceanographic Center (13GYY003NF16), the NSF of Fujian Province, China (2013J01050), and Promotion Program for Young and Middle-aged Teacher in Science and Technology Research of Huaqiao University (ZQN-PY120).

REFERENCES

- (1) Michael, J. P. *Nat. Prod. Rep.* **2005**, *22*, 627–646.
- (2) (a) Kang, S.; Song, B.; Wu, J.; He, M.; Hu, D.; Jin, L.; Zeng, S.; Xue, W.; Yang, S. *Eur. J. Med. Chem.* **2013**, *67*, 14–18. (b) Wang, X.; Wu, Y. *J. Econ. Entomol.* **2012**, *105*, 1019–1023.
- (3) (a) Balbi, A.; Anzaldi, M.; Maccio, C.; Aiello, C.; Mazzei, M.; Gangemi, R.; Castagnola, P.; Miele, M.; Rosano, C.; Viale, M. *Eur. J. Med. Chem.* **2011**, *46*, 5293–5309. (b) Ciayadi, R.; Potdar, M.; Walton, K. L.; Harrison, C. A.; Kelso, G. F.; Harris, S. J.; Hearn, M. T. W. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 5642–5645. (c) Balbi, A.; Anzaldi, M.; Maccio, C.; Aiello, C.; Mazzei, M.; Gangemi, R.; Castagnola, P.; Miele, M.; Rosano, C.; Viale, M. *Eur. J. Med. Chem.* **2011**, *46*, 5293–5309.
- (4) Hang, X.-C.; Fleetham, T.; Turner, E.; Brooks, J.; Li, J. *Angew. Chem., Int. Ed.* **2013**, *52*, 6753–6756.
- (5) For reviews of pyridine synthesis: (a) Bull, J. A.; Mousseau, J. J.; Pelletier, G.; Charette, A. B. *Chem. Rev.* **2012**, *112*, 2642–2713. (b) Heller, B.; Hapke, M. *Chem. Soc. Rev.* **2007**, *36*, 1085–1094.
- (6) For recent examples of transition-metal-catalyzed pyridine synthesis: (a) Prechter, A.; Henrion, G.; Faudot dit Bel, P.; Gagosz, F. *Angew. Chem., Int. Ed.* **2014**, *53*, 4959–4963. (b) Wu, Q.; Zhang, Y.; Cui, S. *Org. Lett.* **2014**, *16*, 1350–1353. (c) Neely, J. M.; Rovis, T. *J. Am. Chem. Soc.* **2014**, *136*, 2735–2738. (d) Lee, H.; Sim, Y.-K.; Park, J.-W.; Jun, C.-H. *Chem. - Eur. J.* **2014**, *20*, 323–333. (e) Jiang, Y.; Park, C.-M. *Chem. Sci.* **2014**, *5*, 2347–2351. (f) Wei, Y.; Yoshikai, N. *J. Am. Chem. Soc.* **2013**, *135*, 3756–3759. (g) Neely, J. M.; Rovis, T. *J. Am. Chem. Soc.* **2013**, *135*, 66–69. (h) Michlik, S.; Kempe, R. *Angew. Chem., Int. Ed.* **2013**, *52*, 6326–6329. (i) Loy, N. S. Y.; Singh, A.; Xu, X.; Park, C.-M. *Angew. Chem., Int. Ed.* **2013**, *52*, 2212–2216. (j) Zhao, M.-N.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Chem. Commun.* **2012**, *48*, 8105–8107. (k) Yamamoto, S.-i.; Okamoto, K.; Murakoso, M.; Kuninobu, Y.; Takai, K. *Org. Lett.* **2012**, *14*, 3182–3185. (l) Gati, W.; Rammah, M. M.; Rammah, M. B.; Couty, F.; Evano, G. *J. Am. Chem. Soc.* **2012**, *134*, 9078–9081. (m) Wang, C.; Li, X.; Wu, F.; Wan, B. *Angew. Chem., Int. Ed.* **2011**, *50*, 7162–7166. (n) Ohashi, M.; Takeda, I.; Ikawa, M.; Ogoshi, S. *J. Am. Chem. Soc.* **2011**, *133*, 18018–18021. (o) Wang, Y.-F.; Chiba, S. *J. Am. Chem. Soc.* **2009**, *131*, 12570–12572. (p) Chiba, S.; Xu, Y.-J.; Wang, Y.-F. *J. Am. Chem. Soc.* **2009**, *131*, 12886–12887. (q) Manning, J. R.; Davies, H. M. L. *J. Am. Chem. Soc.* **2008**, *130*, 8602–8603. (r) Liu, S.; Liebeskind, L. S. *J. Am. Chem. Soc.* **2008**, *130*, 6918–6919. (s) Colby, D. A.; Bergman, R. G.; Ellman, J. A. *J. Am. Chem. Soc.* **2008**, *130*, 3645–3651. (t) Barluenga, J.; Fernandez-Rodriguez, M. A.; Garcia-Garcia, P.; Aguilar, E. *J. Am. Chem. Soc.* **2008**, *130*, 2764–2765.
- (7) For recent examples of organocatalyzed pyridine synthesis: (a) Yoshida, M.; Mizuguchi, T.; Namba, K. *Angew. Chem., Int. Ed.* **2014**, *53*, 14550–14554. (b) Stark, D. G.; Morrill, L. C.; Yeh, P.-P.; Slawin, A. M. Z.; O'Riordan, T. T. J. C.; Smith, A. D. *Angew. Chem., Int. Ed.* **2013**, *52*, 11642–11646. (c) Shi, Z.; Loh, T.-P. *Angew. Chem., Int. Ed.* **2013**, *52*, 8584–8587.
- (8) For recent examples on metal-free pyridine synthesis: (a) Jiang, Y.; Park, C.-M.; Loh, T.-P. *Org. Lett.* **2014**, *16*, 3432–3435. (b) Desrosiers, J.-N.; Kelly, C. B.; Fandrick, D. R.; Nummy, L.; Campbell, S. J.; Wei, X.; Sarvestani, M.; Lee, H.; Sienkiewicz, A.; Sanyal, S.; Zeng, X.; Grinberg, N.; Ma, S.; Song, J. J.; Senanayake, C. H. *Org. Lett.* **2014**, *16*, 1724–1727. (c) Xin, X.; Wang, D.; Li, X.; Wan, B. *Tetrahedron* **2013**, *69*, 10245–10248. (d) Lei, C.-H.; Wang, D.-X.; Zhao, L.; Zhu, J.; Wang, M.-X. *J. Am. Chem. Soc.* **2013**, *135*, 4708–4711. (e) Huang, P.; Zhang, R.; Liang, Y.; Dong, D. *Org. Lett.* **2012**, *14*, 5196–5199. (f) Allais, C.; Lieby-Muller, F.; Constantieux, T.; Rodriguez, J. *Adv. Synth. Catal.* **2012**, *354*, 2537–2544. (g) Kral, K.; Hapke, M. *Angew. Chem., Int. Ed.* **2011**, *50*, 2434–2435. (h) Donohoe, T. J.; Basutto, J. A.; Bower, J. F.; Rathi, A. *Org. Lett.* **2011**, *13*, 1036–1039. (i) Sakai, T.; Danheiser, R. L. *J. Am. Chem. Soc.* **2010**, *132*, 13203–13205. (j) Sha, F.; Huang, X. *Angew. Chem., Int. Ed.* **2009**, *48*, 3458–3461.
- (9) (a) Teo, Y.-C.; Yong, F.-F.; Sim, S. *Tetrahedron* **2013**, *69*, 7279–7284. (b) Raba, A.; Anneser, M. R.; Jantke, D.; Cokoja, M.; Herrmann, W. A.; Kuehn, F. E. *Tetrahedron Lett.* **2013**, *54*, 3384–3387. (c) Liu, Z.-J.; Vors, J.-P.; Gesing, E. R. F.; Bolm, C. *Green Chem.* **2011**, *13*, 42–45. (d) Guillou, S.; Bonhomme, F. J.; Chahine, D. B.; Nesme, O.; Janin, Y. L. *Tetrahedron* **2010**, *66*, 2654–2663.
- (10) (a) Chen, J.; Pang, Q.; Sun, Y.; Li, X. *J. Org. Chem.* **2011**, *76*, 3523–3526. (b) Chu, J.-H.; Lin, P.-S.; Lee, Y.-M.; Shen, W.-T.; Wu, M.-J. *Chem. - Eur. J.* **2011**, *17*, 13613–13620. (c) Chen, J.; Song, G.; Pan, C.-L.; Li, X. *Org. Lett.* **2010**, *12*, 5426–5429.
- (11) (a) Cheng, G.; Zeng, X.; Shen, J.; Wang, X.; Cui, X. *Angew. Chem., Int. Ed.* **2013**, *52*, 13265–8. (b) Yang, X.; Cheng, G.; Shen, J.; Kuai, Cui, X. *Org. Chem. Front.* **2015**, *2*, 366–368. (c) Wang, X.; Cheng, G.; Cui, X. *Chem. Commun.* **2014**, *50*, 652–654. (d) Cheng, G.; Cui, X. *Org. Lett.* **2013**, *15*, 1480–1483. (e) Shen, J.; Cheng, G.; Cui, X. *Chem. Commun.* **2013**, *49*, 10641–10643.
- (12) (a) Martins, M. A. P.; Rossatto, M.; Frizzo, C. P.; Scapin, E.; Buriol, L.; Zanatta, N.; Bonacorso, H. G. *Tetrahedron Lett.* **2013**, *54*, 847–849. (b) Harschneck, T.; Kirsch, S. F. *J. Org. Chem.* **2011**, *76*, 2145–2156. (c) Fananas, F. J.; Arto, T.; Mendoza, A.; Rodriguez, F. *Org. Lett.* **2011**, *13*, 4184–4187. (d) Cacchi, S.; Fabrizi, G.; Filisti, E. *Org. Lett.* **2008**, *10*, 2629–2632.
- (13) (a) Kurita, J.; Iwata, K.; Tsuchiya, T. *Chem. Pharm. Bull.* **1987**, *35*, 3166–3174. (b) Kurita, J.; Iwata, K.; Tsuchiya, T. *J. Chem. Soc., Chem. Commun.* **1986**, 1188–1189.
- (14) CCDC 1029899 (5aa), 1029900 (5bf), and 1029895 (2b) can be obtained from The Cambridge Crystallographic Data Centre via <http://www.ccdc.cam.ac.uk>.
- (15) Lu, T.; Lu, Z.; Ma, Z.-X.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2013**, *113*, 4862–4904.
- (16) We cannot deny the following pathway: a conjugate addition of the nucleophile **3** to intermediate **A** directly delivers compound **4**.