


Catalytic Aerobic Dehydrogenation of Nitrogen Heterocycles Using Heterogeneous Cobalt Oxide Supported on Nitrogen-Doped Carbon

Andrei V. Iosub and Shannon S. Stahl*

Department of Chemistry, University of Wisconsin—Madison, 1101 University Avenue, Madison, Wisconsin 53706, United States

Supporting Information


ABSTRACT: Dehydrogenation of (partially) saturated heterocycles provides an important route to heteroaromatic compounds. A heterogeneous cobalt oxide catalyst, previously employed for aerobic oxidation of alcohols and amines, is shown to be effective for aerobic dehydrogenation of various 1,2,3,4-tetrahydroquinolines to the corresponding quinolines. The reactions proceed in good yields under mild conditions. Other *N*-heterocycles are also successfully oxidized to their aromatic counterparts.



(Hetero)aromatic molecules are ubiquitous in pharmaceuticals and other biologically active molecules.¹ The introduction of substituents on the aromatic framework is typically accomplished by sequential introduction of groups onto a preformed aromatic ring. Recently, our group has been exploring aerobic dehydrogenation of (partially) saturated cyclic compounds as a complementary route to prepare substituted arenes and heteroarenes.² In many cases, these methods enable synthesis of aromatic compounds with substitution patterns and/or functional groups that are difficult to access via traditional aromatic functionalization reactions. This strategy has been employed by us and others in Pd-catalyzed methods for the conversion of cyclohexanone and cyclohexenone derivatives into substituted phenols,³ aryl ethers,⁴ or anilines⁵ and for the conversion of cyclohexene derivatives into substituted arenes.⁶ More recently, we began exploring methods for aerobic dehydrogenation of partially saturated *N*-heterocycles to heteroarenes. Our initial efforts to employ Pd-based catalysts led to unsatisfactory results, and we turned our attention to bioinspired *ortho*-quinone-based catalysts that are effective for the aerobic dehydrogenative aromatization of (partially) saturated nitrogen heterocycles.^{7–9} Collectively, the aerobic dehydrogenation methods represent compelling alternatives to established methods that use stoichiometric reagents, such as DDQ or sulfur.¹⁰ In light of the challenges we encountered in the use of Pd-based catalysts for *N*-heterocycle dehydrogenation, specifically to access quinoline products (Scheme 1A), we elected to test heterogeneous Co-oxide-based catalysts for this application. Here, we present the results of this effort.

For our initial studies we chose 1,2,3,4-tetrahydroquinoline (1a) as a test substrate (Table 1). The product of this dehydrogenation, 2-methylquinoline or quinaldine (2a), resembles the core of compounds that exhibit important pharmaceutical activity, including those for the treatment of schizophrenia (PF-2545920),¹¹ HIV infection (styrylquinolines),¹² malaria (Mefloquine),¹³ asthma (Montelukast),¹⁴ and pain (JTC-801).¹⁵ Therefore, it represents a simple proof-of-concept for use in catalyst screening (Scheme 1B). Our


Scheme 1. Dehydrogenation 1,2,3,4-Tetrahydroquinolines to Quinolines



previously reported homogeneous Pd-based dehydrogenation catalysts^{3a,16} were tested, but they led to only modest yields (49–55%) with rather poor mass balance (Table 1, entries 1–3). In an effort to improve the results, we considered other catalysts, including heterogeneous variants. Heterogeneous Pd catalysts have been successfully used for the dehydrogenation of cyclohexanones and -hexenones.^{3c,4b,5e} Other heterogeneous catalysts composed of noble metals (Ru, Au, Pd₃Pb) have also been shown to be active for aerobic dehydrogenation of the parent 1,2,3,4-tetrahydroquinoline;¹⁷ however, the reactions proceed at high temperatures (>100 °C), they are accompanied by catalyst deactivation, and the results were not extended to substituted tetrahydroquinoline derivatives. During the course of these initial studies, Beller and co-workers reported a unique Co-oxide-based catalyst for aerobic alcohol oxidation (oxidative


Received: June 20, 2015

Published: September 2, 2015

Table 1. Low Temperature Dehydrogenation of 1,2,3,4-Tetrahydroquinoline

entry	catalyst (mol %)	solvent (concn [M])	temp	yield (conv) ^b
1	Pd(TFA) ₂ (5)/ 2-NMe ₂ -py (10)/ pTsOH (20)	DMSO (1)	80	52 (88)
2	Pd(TFA) ₂ (5)/ DMSO (10)	AcOH (0.2)	80	55 (100)
3	Pd(TFA) ₂ (5)/4,5- diazfluoren-9-one (5)	DMSO (0.2)	80	49 (61)
4 ^{a,c}	CoO _x -phen/AB	MeOH (0.125)	60	92 (94)
5 ^a	CoO _x -phen/AB	MeOH (0.125)	60	43 (43)
6 ^{a,c}	CoO _x -bpy/AB	MeOH (0.125)	60	0 (1)
7 ^{a,c}	CoO _x -terpy/AB	MeOH (0.125)	60	82 (100)
8 ^{a,c}	CoO _x -phd/AB	MeOH (0.125)	60	67 (74)
9 ^{a,c}	MnO _x -phen/AB	MeOH (0.125)	60	16 (32)
10 ^{a,c}	FeO _x -phen/AB	MeOH (0.125)	60	0 (52)
11 ^{a,c}	NiO _x -phen/AB	MeOH (0.125)	60	0 (13)
12 ^{a,c}	CoO _x -phen/ Vulcan XC72R	MeOH (0.125)	60	92 (100)
13 ^a	Pd/C	xylene (0.2)	60	0 (0)
14 ^a	Pd/C	MeOH (0.125)	60	13 (30)
15 ^{a,d,e}	activated C	xylene (0.2)	60	29 (66)
16 ^{a,d,f}	Co(salen)	EtOH (0.066)	60	28 (63)

^a1a (0.10 mmol), solvent, 2.5 mol % [cat.], 1 atm of O₂, orbital mixing, 60 °C, 4 h. ^b% Yields and conv determined by ¹H NMR. ^c1 equiv of K₂CO₃ added. ^dMagnetic stirring was employed. ^e100 wt % catalyst, 24 h. ^f1 mol % catalyst was used, 24 h. C = carbon; AB = Acetylene Black.



esterification of primary alcohols).^{18,19} More recently, the same researchers have applied these and related catalysts to the conversion of alcohols and amines to nitriles.²⁰ Overall, these catalysts represent appealing base-metal alternatives to heterogeneous Pd/C and other supported noble metals, and we reasoned that similar catalysts could promote aerobic dehydrogenation of heterocycles.²¹

The Co-oxide catalysts were prepared according to the protocol reported by Beller and co-workers.¹⁸ Briefly, cobalt(II) acetate was stirred with different bi- and tridentate nitrogen containing ligands in ethanol for 30 min, then acetylene black (AB) or Vulcan XC72R was added as a conductive carbon black support, and the mixture was refluxed. After drying, the black powder was pyrolyzed under an argon flow at 800 °C for 2 h. This procedure enabled the synthesis of cobalt based nanoparticles of varying sizes, encapsulated by nitrogen-rich graphene-like layers. This specific architecture has been suggested to account for the notable reactivity of these catalysts.¹⁸

Using the procedure just noted, a number of related catalysts were prepared, in which the identities of the nitrogen-containing ligand, the metal source, and/or the carbon support was varied. Our initial studies were performed with acetylene black (AB) as the carbon support because it was already available in our lab. A catalyst prepared with this material, using Co(OAc)₂ as the Co


source and 1,10-phenanthroline (phen) as the nitrogen source, provided a 92% yield of quinoline in the presence of 1 equiv of K₂CO₃ as a base. The reaction was complete within 4 h in MeOH at only 60 °C (Table 1, entry 4). Without any base, the catalyst was still active, but led to a lower final yield (43%, entry 5).²² Catalysts prepared analogously using other nitrogen-based ligands instead of phen were also investigated (entries 6–8).²² With 2,2'-bipyridine (bpy) as the nitrogen source, no product was obtained, while, with 2,6-di(2-pyridyl)pyridine (terpy) and 1,10-phenanthroline-5,6-dione (phd), 82% and 67% yields were obtained, albeit lower than when using phen. Further investigations revealed that the ratio of cobalt to phen in the catalyst preparation procedure had little impact on this dehydrogenation.²² Next, catalysts with metals other than cobalt were prepared using phen as the nitrogen source and acetylene black as the carbon source. These heterogeneous Mn, Fe, and Ni oxide materials were tested under the reaction conditions. The MnO_x-based catalyst provided low yields of product, while the rest were inactive (entries 9–11). Upon obtaining a sample of Vulcan XC72R, a Co(OAc)₂/phen-derived catalyst was prepared with this carbon source (entry 12). This catalyst led to a product yield similar to that obtained with the catalyst supported on AB, but it showed increased initial activity.²²

Control reactions were also conducted to verify the necessity of the pyrolysis, heterogeneity, and “nitrogen-doping” of the catalyst.²² Co(OAc)₂ alone was not active for the reaction, in the presence of either a nitrogenous ligand or carbon black. The carbon black support (AB) was also completely inactive for the dehydrogenation, with or without pyrolysis and/or nitrogen doping. Finally, a mixture of Co(OAc)₂/1,10-phenanthroline impregnated on acetylene black, but not pyrolyzed, was also inactive. Collectively, the results show that pyrolysis of Co(OAc)₂/phen in a 1:2 mixture on AB and Vulcan XC72R at 800 °C under Ar generates effective catalysts. These compositions align with the catalyst previously optimized and characterized by Beller and co-workers, and their nomenclature “Co₃O₄-NGr/C” is adopted below.

The Co₃O₄-NGr/C catalyst outperforms other commonly used dehydrogenation catalysts, such as Pd/C^{10a} and activated carbon. Moreover, it exhibits good activity at 60 °C, which is lower than temperatures typically required for dehydrogenation. When Pd/C was tested under comparable conditions, none of the dehydrogenation product was obtained in xylene and only a low yield was observed in methanol (Table 1, entries 13, 14). Activated carbon has been shown to promote dehydrogenation of 1,2,3,4-tetrahydroquinolines at 120 °C under O₂ with a 100 wt % loading of carbon.²³ At the lower temperatures used here, however, use of 100 wt % activated carbon affords only a 29% yield of product (entry 15). Co(salen) has been used as a catalyst previously for dehydrogenation of the parent 1,2,3,4-tetrahydroquinoline at 60 °C,²⁴ but the presence of the 2-methyl group in tetrahydroquinoline significantly attenuates the yield (28%, entry 16).

The Co₃O₄-NGr/C catalyst was then tested in the dehydrogenation of a variety of 1,2,3,4-tetrahydroquinolines under analogous reaction conditions (Scheme 2).²⁵ The parent quinoline 2b was obtained in 90% yield and 2-phenylquinoline 2c in 92% yield after 6 h. The presence of functional groups on a 2-aryl substituent influenced the dehydrogenation reactivity. Substrates containing –SMe and –CF₃ groups reacted well (2d, 2e), while those containing –NO₂ and –CN groups needed more catalyst and longer reaction times to reach completion,

Scheme 2. Substrate Scope for the Dehydrogenation of 1,2,3,4-Tetrahydroquinolines to Quinolines^a



^aConditions: **1** (0.5 mmol), 4 mL of MeOH, 25 mg of catalyst (2.5 mol %), 1 equiv of K_2CO_3 , 1 atm of O_2 , orbital stirring, 60 °C. ^b50 mg of catalyst were used. ^cReaction on 10 mmol scale of **1m**, magnetic stirring.


affording lower yields of the desired products (**2f**, **2g**). Substrates with 2-thienyl and 2-furyl groups led to high yields of the corresponding quinolines (**2h**, **2i**), showing no inhibition by the heterocyclic substituent. The results with **2c–2i** are noteworthy in demonstrating the compatibility of the catalyst with 2-arylquinolines, a motif present in pharmaceutically active compounds. The reaction forming 3-methylquinoline (**2j**) exhibited a lower mass balance and product yield; however, 4-methyl derivatives were obtained in high yields (**1k**, **1l**).

A number of tetrahydroquinolines with substitution at positions 6–8 were investigated (Scheme 2). Derivatization at these positions is usually accessible via selective electrophilic aromatic substitution reactions of tetrahydroquinolines. In contrast, the reactivity of quinolines in similar reactions is usually unselective, yielding multiple products. First, substitution at the 6-position was explored. Substrates with electron-donating substituents, such as methoxy and methyl, underwent dehydrogenation in high yields (**2m**, **2n**). The dehydrogenation of **1m** was also carried out on 10 mmol scale (1.47 g), and the product was obtained in a 94% yield. The presence of halogens F–, Cl–, and Br– at the 6-position led to good yields of dehydrogenation products **2o–2q**. These observations highlight the advantages of this Co-based catalyst over Pd/C catalysts for dehydrogenations reactions, as the latter can promote undesired reactivity with aryl halides, and the –Cl and –Br substituents are appealing functional groups for subsequent derivatization. The 6,7-benzo-fused derivative **1r** underwent dehydrogenation in good yield to the benzoquinoline **2r**. Dehydrogenation of substrates with substituents in the 8 position also led to products

in good yield (77% for the –OMe and 72% for the –Me derivatives **2s** and **2t**). The reactions involving substrates **1r–1t** were somewhat slower, presumably due to steric effects.

With these successful results in hand, we tested a number of other partially saturated *N*-heterocycles containing secondary amines for their reactivity toward dehydrogenation (Scheme 3).

Scheme 3. Dehydrogenation of Other Classes of *N*-Heterocycles^{a,b}



^aConditions: Substrate (0.1 mmol), 0.8 mL of MeOH, 5 mg of catalyst, 1 equiv of K_2CO_3 , 1 atm of O_2 , orbital stirring. ^bYields and conv determined by ^1H NMR. ^cEtOH was used as solvent. ^dCatalyst prepared with Co:phen ratio of 1:1; 10 mg of catalyst used.

The reaction of 1,2,3,4-tetrahydroquinoxaline **3** under the standard reaction conditions led to quinoxaline **4** in 97% yield. The readily accessible substrates, indoline **5** and 2-methyl indoline **7**, exhibited excellent reactivity under the reaction conditions and afforded high yields of products at room temperature in a short time. This reactivity compares favorably with the previously reported aerobic dehydrogenation of indolines with 10 mol % of a molecular Co-salen catalyst.²⁶ Hayashi and co-workers also reported dehydrogenation of the parent indoline **5** using 100 wt % of activated carbon, although a longer reaction time and higher temperature were required.^{27a} A tertiary amine substrate, *N*-methyl indoline **9** was less reactive, affording only a 33% yield of the desired indole. The oxidation of Hantzsch ester **11** was also successful, exhibiting a yield of 95% after only 2 h at room temperature, which again compares favorably with results obtained with activated carbon.^{27b} Finally, the more challenging dehydrogenation of 2-phenylimidazoline **13** led to a 43% product yield, with increased catalyst loading.

Empirical optimization efforts with this substrate showed that the best catalyst was prepared with a 1:1 ratio of Co/phen. In this case, activated carbon is a more effective reagent for dehydrogenation, affording the desired imidazole in 84% yield at 120 °C.^{27c}

In conclusion, we have shown that the dehydrogenation of 1,2,3,4-tetrahydroquinolines and other important *N*-heterocycles can be achieved using a cobalt oxide based catalyst supported on nitrogen-doped carbon under mild conditions. This heterogeneous first-row transition-metal catalyst represents a promising alternative to stoichiometric DDQ or other catalysts based on Pd for the synthesis of biologically active heteroaromatic compounds.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b01790.

Optimization data, experimental procedures, and spectroscopic characterization data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: stahl@chem.wisc.edu.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the following individuals for assistance: Dr. R. Pokhrel (synthesis of catalysts), Dr. E. J. Popczun, M. S. Ahmed, and Y. Preger (pXRD), and J. C. King (XPS) (all from UW—Madison). This work was funded by the NIH (R01-GM100143). NMR facilities were partially supported by the NSF (CHE-0342998, CHE-1048642) and NIH (S10 RR08389). MS instrumentation was partially supported by the NIH (S10 RR024601).

■ REFERENCES

- (1) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257.
- (2) For leading references, see: (a) Izawa, Y.; Pun, D.; Stahl, S. S. *Science* **2011**, *333*, 209. (b) Izawa, Y.; Zheng, C.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2013**, *52*, 3672. (c) Pun, D.; Diao, T.; Stahl, S. S. *J. Am. Chem. Soc.* **2013**, *135*, 8213. (d) Iosub, A. V.; Stahl, S. S. *J. Am. Chem. Soc.* **2015**, *137*, 3454.
- (3) (a) Imahori, T.; Tokuda, T.; Taguchi, T.; Takahata, H. *Org. Lett.* **2012**, *14*, 1172. (b) Kikushima, K.; Nishina, Y. *RSC Adv.* **2013**, *3*, 20150. (c) Zhang, J.; Jiang, Q.; Yang, D.; Zhao, X.; Dong, Y.; Liu, R. *Chem. Sci.* **2015**, *6*, 4674.
- (4) (a) Simon, M.-O.; Girard, S. A.; Li, C.-J. *Angew. Chem., Int. Ed.* **2012**, *51*, 7537. (b) Sutter, M.; Sotto, N.; Raoul, Y.; Métya, E.; Lemaire, M. *Green Chem.* **2013**, *15*, 347.
- (5) (a) Girard, S. A.; Hu, X.; Knauber, T.; Zhou, F.; Simon, M.-O.; Deng, G.-J.; Li, C.-J. *Org. Lett.* **2012**, *14*, 5606. (b) Hajra, A.; Wei, Y.; Yoshikai, N. *Org. Lett.* **2012**, *14*, 5488. (c) Xie, Y.; Liu, S.; Liu, Y.; Wen, Y.; Deng, G.-J. *Org. Lett.* **2012**, *14*, 1692. (d) Hong, W. P.; Iosub, A. V.; Stahl, S. S. *J. Am. Chem. Soc.* **2013**, *135*, 13664. (e) Sutter, M.; Duclos, M.-C.; Guichere, B.; Raoul, Y.; Métya, E.; Lemaire, M. *ACS Sustainable Chem. Eng.* **2013**, *1*, 1463.
- (6) See ref 2d and the following: Kandukuri, S. R.; Oestreich, M. J. *Org. Chem.* **2012**, *77*, 8750.
- (7) (a) Wendlandt, A. E.; Stahl, S. S. *J. Am. Chem. Soc.* **2014**, *136*, 506. (b) Wendlandt, A. E.; Stahl, S. S. *J. Am. Chem. Soc.* **2014**, *136*, 11910.
- (8) For other catalytic aerobic dehydrogenation routes to substituted heterocycles, see: (a) Flanagan, J. C. A.; Dornan, L. M.; McLaughlin, M. G.; McCreanor, N. G.; Cook, M. J.; Muldoon, M. J. *Green Chem.* **2012**, *14*, 1281. (b) Han, B.; Yang, X.-L.; Wang, C.; Bai, Y.-W.; Pan, T.-C.; Chen, X.; Yu, W. *J. Org. Chem.* **2012**, *77*, 1136. (c) Yuan, H.; Yoo, W.-J.; Miyamura, H.; Kobayashi, S. *Adv. Synth. Catal.* **2012**, *354*, 2899. (d) Dawsey, A. C.; Li, V.; Hamilton, K. C.; Wang, J.; Williams, T. J. *Dalton Trans.* **2012**, *41*, 7994. (e) Chen, S.; Hossain, M. S.; Foss, F. W., Jr. *ACS Sustainable Chem. Eng.* **2013**, *1*, 1045. (f) Chen, Z.; Chen, J.; Liu, M.; Ding, J.; Gao, W.; Huang, X.; Wu, H. *J. Org. Chem.* **2013**, *78*, 11342. (g) Yu, J.; Xu, J.; Lu, M. *Appl. Organometal. Chem.* **2013**, *27*, 606.
- (9) For other homogeneous catalytic methods for dehydrogenation of 1,2,3,4-tetrahydroquinolines, see: (a) Yamaguchi, R.; Ikeda, C.; Takahashi, Y.; Fujita, K.-i. *J. Am. Chem. Soc.* **2009**, *131*, 8410. (b) Wu, J.; Talwar, D.; Johnston, S.; Yan, M.; Xiao, J. *Angew. Chem., Int. Ed.* **2013**, *52*, 6983. (c) Chakraborty, S.; Brennessel, W. W.; Jones, W. D. *J. Am. Chem. Soc.* **2014**, *136*, 8564.
- (10) (a) Fu, P. P.; Harvey, R. G. *Chem. Rev.* **1978**, *78*, 317. (b) Buckle, D. R. *Encyclopedia of Reagents for Organic Synthesis*; John Wiley & Sons, Inc.: New York, 2010.
- (11) Verhoest, P. R.; Chapin, D. S.; Corman, M.; Fonseca, K.; Harms, J. F.; Hou, X.; Marr, E. S.; Menniti, F. S.; Nelson, F.; O'Connor, R.; Pandit, J.; Proulx-LaFrance, C.; Schmidt, A. W.; Schmidt, C. J.; Suiciak, J. A.; Liras, S. *J. Med. Chem.* **2009**, *52*, 5188.
- (12) Mouscadet, J.-F.; Desmaële, D. *Molecules* **2010**, *15*, 3048.
- (13) Croft, A.; Garner, P. *BMJ* **1997**, *315*, 1412.
- (14) Belley, M. L.; Leger, S.; Labelle, M.; Roy, P.; Xiang, Y. B.; Guay, D. U.S. Patent 5,565,473, Oct. 15, 1996.
- (15) Sestili, I.; Borioni, A.; Mustazza, C.; Rodomonte, A.; Turchetto, L.; Sbraccia, M.; Riitano, D.; Del Giudice, M. R. *Eur. J. Med. Chem.* **2004**, *39*, 1047.
- (16) (a) Diao, T.; Stahl, S. S. *J. Am. Chem. Soc.* **2011**, *133*, 14566. (b) Diao, T.; Wadzinski, T. J.; Stahl, S. S. *Chem. Sci.* **2012**, *3*, 887.
- (17) For heterogeneous catalysts based on Pd: (a) Furukawa, S.; Suga, A.; Komatsu, T. *Chem. Commun.* **2014**, *50*, 3277. Ru: (b) Yamaguchi, K.; Mizuno, N. *Angew. Chem., Int. Ed.* **2003**, *42*, 1480. (c) Li, F.; Chen, J.; Zhang, Q.; Wang, Y. *Green Chem.* **2008**, *10*, 553. (d) Yamaguchi, K.; Kim, J. W.; He, J.; Mizuno, N. *J. Catal.* **2009**, *268*, 343. Au: (e) So, M.-H.; Liu, Y.; Ho, C.-H.; Che, C.-M. *Chem. - Asian J.* **2009**, *4*, 1551.
- (18) Jagadeesh, R. V.; Junge, H.; Pohl, M.-M.; Radnik, J.; Brückner, A.; Beller, M. *J. Am. Chem. Soc.* **2013**, *135*, 10776.
- (19) See also: Zhu, J.; Kailasam, K.; Fischer, A.; Thomas, A. *ACS Catal.* **2011**, *1*, 342.
- (20) (a) Jagadeesh, R. V.; Junge, H.; Beller, M. *Nat. Commun.* **2014**, *5*, 4123. (b) Jagadeesh, R. V.; Junge, H.; Beller, M. *ChemSusChem* **2015**, *8*, 92.
- (21) For a related recent application, see: Cui, X.; Li, Y.; Bachmann, S.; Scalone, M.; Surkus, A.-E.; Junge, K.; Topf, C.; Beller, M. *J. Am. Chem. Soc.* **2015**, *137*, 10652.
- (22) See Supporting Information for details.
- (23) Tanaka, T.; Okunaga, K.-i.; Hayashi, M. *Tetrahedron Lett.* **2010**, *51*, 4633.
- (24) Nishinaga, A.; Yamazaki, S.; Matsuura, T. *Tetrahedron Lett.* **1988**, *29*, 4115.
- (25) CAUTION: Because of the flammability of methanol and O₂, large scale reactions should be conducted below the LOC of MeOH (approx. 7–8%): Osterberg, P. M.; Niemeier, J. K.; Welch, C. J.; Hawkins, J. M.; Martinelli, J. R.; Johnson, T. E.; Root, T. W.; Stahl, S. S. *Org. Process Res. Dev.* **2015**, DOI: 10.1021/ops.500328f.
- (26) Inada, A.; Nakamura, Y.; Morita, Y. *Chem. Lett.* **1980**, 1287.
- (27) (a) Nomura, Y.; Kawashita, Y.; Hayashi, M. *Heterocycles* **2007**, *74*, 629. (b) Nakamichi, N.; Kawashita, Y.; Hayashi, M. *Synthesis* **2004**, *7*, 1015. (c) Haneda, S.; Okui, A.; Ueba, C.; Hayashi, M. *Tetrahedron* **2007**, *63*, 2414.