

## Alkene *anti*-Dihydroxylation with Malonoyl Peroxides


Carla Alamillo-Ferrer,<sup>†</sup> Stuart C. Davidson,<sup>†</sup> Michael J. Rawling,<sup>†</sup> Natalie H. Theodoulou,<sup>†,‡</sup> Matthew Campbell,<sup>‡</sup> Philip G. Humphreys,<sup>‡</sup> Alan R. Kennedy,<sup>†</sup> and Nicholas C. O. Tomkinson<sup>\*,†</sup>

<sup>†</sup>WestCHEM, Department of Pure and Applied Chemistry, Thomas Graham Building, University of Strathclyde, 295 Cathedral Street, Glasgow G1 1XL, United Kingdom

<sup>‡</sup>GlaxoSmithKline Medicines Research Centre, Gunnels Wood Road, Stevenage SG1 2NY, United Kingdom

### S Supporting Information

**ABSTRACT:** Malonoyl peroxide **1**, prepared in a single step from the commercially available diacid, is an effective reagent for the *anti*-dihydroxylation of alkenes. Reaction of **1** with an alkene in the presence of acetic acid at 40 °C followed by alkaline hydrolysis leads to the corresponding diol (35–92%) with up to 13:1 *anti*-selectivity. A mechanism consistent with experimental findings is proposed that accounts for the selectivity observed.




The prevalence of 1,2-diols in natural products and biologically active molecules together with the synthetic versatility of this functionality has provided the motivation to discover new methods for their preparation. Within this toolkit, of particular note are approaches for the dihydroxylation of alkenes, which have delivered innovation and advancement in methodology development.<sup>1</sup> Although a number of procedures have been developed for the *syn*-dihydroxylation of alkenes using both metal-catalyzed<sup>2</sup> and -free procedures,<sup>3</sup> introduction of the *anti*-1,2-diol has received significantly less attention.<sup>4</sup>

The majority of methods to access *anti*-1,2-diols involve the ring-opening of epoxides.<sup>5</sup> The required epoxides can be accessed in an asymmetric manner via processes including the Sharpless,<sup>6</sup> Jacobsen<sup>7</sup> and Shi<sup>8</sup> procedures. Regio- and stereoselective reaction of the epoxides with oxygen nucleophiles then leads to the *anti*-diol.<sup>9</sup> More recently, List has reported an organocatalytic asymmetric hydrolysis of *meso*-epoxides, providing a useful alternative to these well-established methods.<sup>4b</sup> Alexanian et al. have developed both intra-<sup>10</sup> and intermolecular<sup>11</sup> methods for the stereoselective *anti*-dioxxygenation of alkenes using hydroxamic acids and oxygen as the sources of the new C–O bonds, providing the products in good yield and moderate selectivities. A more established method to access *anti*-1,2-diols from alkenes is the Prévost reaction,<sup>12</sup> which proceeds through an iodonium intermediate.<sup>13</sup>

It has been established that malonoyl peroxides can be used for the *syn*-dihydroxylation of alkenes.<sup>14</sup> For example, treatment of stilbene **2** with 1.2 equiv of peroxide **1** in the presence of water followed by basic hydrolysis leads to *syn*-diol **3** (86%; 33:1 *syn:anti*) (Scheme 1). Within this manuscript, we show that malonoyl peroxides can also be used in a complementary procedure to secure the *anti*-1,2-diol.<sup>15</sup>

The mechanistic pathway determined for the *syn*-dihydroxylation of alkenes with malonoyl peroxides is outlined in Scheme 2.<sup>16</sup> Reaction of alkene **2** with peroxide **1** leads to zwitterionic species **5**, which can cyclize to give either the 7-membered ring **6** as a single stereoisomer (minor product) or

### Scheme 1. Malonoyl Peroxides in the Dioxxygenation of Alkenes



the dioxonium species **7** (major product). Hydrolysis of **7** through the addition of water at carbon **A** leads to ester **10**. Basic hydrolysis of a crude mixture of **6** and **10** provides *syn*-diol **3** in excellent yield and outstanding selectivity (86%; 33:1 *syn:anti*).


Reactive species related to dioxonium **7** have been described as intermediates within the Prévost reaction.<sup>17</sup> Addition of a carboxylic acid to this class of compound leads to the corresponding *anti*-dioxxygenated species. We wished to determine if a similar reaction pathway could be developed whereby addition of acetic acid to **7** at carbon **B** would lead to diester **8**, hydrolysis of which would provide the *anti*-diol product **4**. Herein, we describe the development of this procedure providing a simple, peroxide-mediated metal-free method for the *anti*-dihydroxylation of alkenes.

Our investigations began by replacement of the water used in the published *syn*-dihydroxylation procedure with acetic acid (Table 1). Reaction of stilbene **2** with 1.2 equiv of peroxide **1** in

Received: September 15, 2015

Published: October 1, 2015

Scheme 2. Mechanistic Course of Dioxygenation with Malonoyl Peroxide 1

Table 1. Optimization of *anti*-Dihydroxylation<sup>a</sup>

entry	solvent	RCO <sub>2</sub> H (equiv)	R	1 (equiv)	yield (%) <sup>b</sup>	<i>anti:syn</i> <sup>c</sup>
1 <sup>d</sup>	CHCl <sub>3</sub>	1	Me	1.2	78	1:2
2 <sup>d</sup>	AcOH	35	Me	1.2	66	2:1
3	AcOH	35	Me	1.2	43	4:1
4	CH <sub>2</sub> Cl <sub>2</sub>	1	Me	1.2	78	6:1
5	PhMe	1	Me	1.2	72	4:1
6	THF	1	Me	1.2	43	1:1
7	CH <sub>2</sub> Cl <sub>2</sub>	2	Me	1.2	72	7:1
8	CH <sub>2</sub> Cl <sub>2</sub>	3	Me	1.2	77	5:1
9	CH <sub>2</sub> Cl <sub>2</sub>	5	Me	1.2	75	5:1
10 <sup>e</sup>	CH <sub>2</sub> Cl <sub>2</sub>	2	Me	1.2	77	6:1
11	CH <sub>2</sub> Cl <sub>2</sub>	2	Me	1.5	92	7:1
12	CH <sub>2</sub> Cl <sub>2</sub>	3	Me	1.5	95	6:1
13	CH <sub>2</sub> Cl <sub>2</sub>	3	Me	2.0	91	6:1
14	CH <sub>2</sub> Cl <sub>2</sub>	2	Ph	1.5	75	6:1
15	CH <sub>2</sub> Cl <sub>2</sub>	2	4-MeOC <sub>6</sub> H <sub>4</sub>	1.5	68	6:1


<sup>a</sup>All reactions performed in duplicate with stilbene (1 mmol) at 0.5 M concentration. <sup>b</sup>Isolated yield after column chromatography. <sup>c</sup>Determined by <sup>1</sup>H NMR spectroscopy on a crude reaction mixture. <sup>d</sup>Bench acetic acid was used for entries 1 and 2. In entries 3–15, the acid was dried prior to use; see Supporting Information for full details. <sup>e</sup>Ac<sub>2</sub>O added at 0.7 equiv.

the presence of one equiv of acetic acid (40 °C, 24 h) followed by basic hydrolysis gave diol 4 (entry 1; 78%, 1:2 *anti:syn*). Although these conditions favored the formation of the *syn*-diol, the diastomeric ratio was significantly different than that observed in the presence of water (1:33 *anti:syn*). This suggested our mechanistic hypothesis was correct such that the stereochemical outcome of the reaction could be reversed by changing the additive. Performing the reaction in neat acetic acid rendered the reaction *anti*-selective (entry 2; 66%, 2:1 *anti:syn*), and drying the acetic acid improved this ratio further (entry 3; 43%, 4:1 *anti:syn*). A brief examination of the reaction medium (entries 4–6) showed less polar solvents improved the *anti:syn* ratio with dichloromethane emerging as the best option

(entry 4; 78%, 6:1 *anti:syn*). Altering the equivalents of acid and peroxide (entries 7–13) led to an optimized set of reaction conditions that provided the product in excellent yield and good selectivity (entry 11; 92%, 7:1 *anti:syn*). Conducting the reaction in the presence of acetic anhydride to sequester any water present in the reaction mixture led to no significant change in the stereochemical outcome (entry 10; 77%, 6:1 *anti:syn*). It also proved possible to conduct the reaction in the presence of alternative acids such as benzoic acid (entry 14; 75%, 6:1 *anti:syn*) and 4-methoxybenzoic acid (entry 15; 68%, 6:1 *anti:syn*), neither of which had a notable influence on the reaction outcome.

Having developed effective reaction conditions for the *anti*-dihydroxylation procedure, we turned our attention to exploring the substrate scope (Scheme 3). The reaction proceeded well with alternative stilbene derivatives (entries 2, 3, and 6), delivering the products in high yield and up to 6:1 *anti:syn* selectivity. Use of *cis*-stilbene as a substrate gave product 3 in reasonable yield but with low levels of selectivity (entry 4; 79% yield, 4:3 *anti:syn*), suggesting that for this compound a more effective strategy would be to perform a *syn*-dihydroxylation on the *trans*-alkene substrate.<sup>14a</sup> We believe this erosion of selectivity is due to a rapid  $\sigma$ -bond rotation in intermediate 5. The reaction was also effective for a number of styrene derivatives (entries 5 and 7–15).  $\beta$ -Methylstyrene provided access to *anti*-product 15 in excellent yield and acceptable selectivity (entry 7; 89%, 4:1 *anti:syn*). Substituents on the aromatic ring were tolerated (entries 8–11); however, electron-withdrawing substituents significantly slowed the reaction. For example, a styrene derivative with a nitro group on the aromatic ring reached 45% completion (35% yield) after 24 h, the mass balance of the reaction being accounted for by unreacted starting material (entry 10). Increasing the steric demands of both the aromatic (entry 11; 90%, 5:1 *anti:syn*) and aliphatic (entry 12; 77%, 4:1 *anti:syn*) alkene substituents had little effect on the stereochemical outcome of the transformation.


From a mechanistic perspective, the addition of acetic acid to the dioxonium ion derived from a nonsymmetrical alkene can occur at two possible carbon centers. In the reaction of 4-methylstyrene 30 with peroxide 1 in the presence of acetic acid, the potential products are the isomeric esters 32 and 33 (Scheme 4). Interestingly, the addition of acetic acid occurs

Scheme 3. Substrate Scope<sup>a</sup>

<sup>a</sup>Yields quoted are isolated yields. All reactions run in duplicate. Stereoselectivities were determined by <sup>1</sup>H NMR spectroscopy of the crude reaction mixture.

exclusively at the benzylic center (compare to **B** in Scheme 2), resulting in *bis*-ester **32**. Independent synthesis of **33** (see Supporting Information) and careful examination of the crude reaction mixture showed that none of this *bis*-ester was formed to the detection limits of <sup>1</sup>H NMR spectroscopy. This shows that acetic acid addition is directed by electronic factors consistent with an ionic mechanism under the acidic reaction conditions employed. This finding was supported by the reaction of 2-substituted-1*H*-indenes with malonyl peroxide **1**,


Scheme 4. Selectivity of Acetic Acid Addition



which resulted in *anti*-diols with excellent selectivities (Scheme 3, entries 16–19; 8:1–13:1 *anti:syn*). These reactions proceeded with complete control over the regiochemistry of acetic acid addition as shown by the relative stereochemistry between the benzylic oxygen and the 2-substituent of the indene. This was also the case in the reactions of dihydronaphthalene derivatives (Scheme 3, entries 20 and 21).

Examination of the crude mixture by <sup>1</sup>H NMR spectroscopy from the reaction of stilbene **2** and peroxide **1** prior to hydrolysis (under the optimized reaction conditions) showed that none of acid **10** was present, indicating that fortuitous water was not responsible for erosion in stereoselectivity.<sup>16</sup> This reinforced our conclusions from the reaction conducted in the presence of acetic anhydride (Table 1, entry 10). Surprisingly, none of expected diester **8** was observed, decarboxylation having occurred during the course of the transformation to deliver **34** as the major product (Scheme 5).


Scheme 5. Rationalizing Selectivity Levels Observed



Considering the proposed mechanistic course of the reaction, we reasoned that the erosion of selectivity therefore arose through formation of the 7-membered ring intermediate **6**, which would lead to *syn*-diol **3** upon hydrolysis. This was confirmed by the isolation of **34** (31:1 *anti:syn*) and by subjecting it to basic hydrolysis to give *anti*-diol **4** (69% from alkene; 31:1 *anti:syn*). We therefore concluded that, in the case of *trans*-stilbene, compromise in selectivity occurs primarily through formation of **6** rather than through  $\sigma$ -bond rotation in intermediate **5**.

Further confirmation of the regioselectivity in the addition of acetic acid came from the use of isotopically labeled acetic acid (Scheme 6). Reaction of alkene **35** with malonyl peroxide **1** in the presence of labeled acetic acid gave adduct **36** with two

Scheme 6. Isotopic Labeling Experiments and X-ray Structure of Unlabeled 36



<sup>a</sup>Using 1.2 mmol NaIO<sub>4</sub> per g of silica gel 60 H (3.6 equiv).

labels incorporated into the structure of the *bis*-ester. The relative stereochemistry of 36 was confirmed by single crystal X-ray analysis of the <sup>16</sup>O variant. Treatment of 36 with methylamine led to *anti*-diol 37, where the label was incorporated exclusively at the benzylic center together with the amides 38 and 39. The location of the label in 37 was reinforced by oxidative cleavage of the diol under anhydrous reaction conditions<sup>18</sup> to give labeled benzaldehyde 40 and unlabeled cyclohexanecarboxaldehyde 41.

In summary, we have described a simple and effective method to alter the stereochemical course of the malonoyl peroxide-mediated metal-free alkene dihydroxylation through judicious choice of additive. Reaction of an alkene and peroxide 1 in the presence of water leads to the *syn*-1,2-diol, whereas use of dry acetic acid delivers the *anti*-1,2-diol. This powerful discovery arose through an intimate understanding of the mechanistic course of the reaction, which led us to effectively intercept an intermediate to deliver a complementary diastereoisomeric product using the same reagent. Ring-opening of the dioxonium intermediate proceeds with good regio- and stereoselectivity and bodes well for the development of additional bond-forming reactions. Development of methods involving alternative nucleophiles is ongoing and will be reported in due course.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b02674.

Analytical data, experimental procedures, and NMR spectra for all compounds reported (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: Nicholas.Tomkinson@strath.ac.uk

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors thank the EPSRC, GlaxoSmithKline, and the University of Strathclyde for financial support and the EPSRC Mass Spectrometry Service, Swansea, for high-resolution spectra.

## ■ REFERENCES

- (1) Kolb, H. C.; VanNieuwenhze, M. S.; Sharpless, K. B. *Chem. Rev.* **1994**, *94*, 2483.
- (2) (a) Bataille, C. J. R.; Donohoe, T. J. *Chem. Soc. Rev.* **2011**, *40*, 114. (b) Asghar, S. F.; Lewis, S. E. *Annu. Rep. Prog. Chem., Sect. B: Org. Chem.* **2011**, *107*, 34. (c) Christie, S. D. R.; Warrington, A. D. *Synthesis* **2008**, 1325.
- (3) Rawling, M. J.; Tomkinson, N. C. O. *Org. Biomol. Chem.* **2013**, *11*, 1434.
- (4) For selected recent examples of novel methods for the formation of the 1,2-diol functionality, see: (a) Dornan, P. K.; Wickens, Z. K.; Grubbs, R. H. *Angew. Chem., Int. Ed.* **2015**, *54*, 7134. (b) Monaco, M. R.; Prévost, S.; List, B. *Angew. Chem., Int. Ed.* **2014**, *53*, 8142. (c) Aciro, C.; Davies, S. G.; Kurosawa, W.; Roberts, P. M.; Russell, A. J.; Thomson, J. E. *Org. Lett.* **2009**, *11*, 1333.
- (5) Usui, Y.; Sato, K.; Tanaka, M. *Angew. Chem., Int. Ed.* **2003**, *42*, 5623.
- (6) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* **1980**, *102*, 5974.
- (7) Zhang, W.; Loebach, J. L.; Wilson, S. R.; Jacobsen, E. N. *J. Am. Chem. Soc.* **1990**, *112*, 2801.
- (8) Tu, Y.; Wang, Z.-X.; Shi, Y. *J. Am. Chem. Soc.* **1996**, *118*, 9806.
- (9) (a) Iida, T.; Yamamoto, N.; Matsunaga, S.; Woo, H.-G.; Shibasaki, M. *Angew. Chem., Int. Ed.* **1998**, *37*, 2223. (b) Vogl, E. M.; Matsunaga, S.; Kanai, M.; Iida, T.; Shibasaki, M. *Tetrahedron Lett.* **1998**, *39*, 7917. (c) Matsunaga, S.; Das, J.; Roels, J.; Vogl, E. M.; Yamamoto, N.; Iida, T.; Yamaguchi, K.; Shibasaki, M. *J. Am. Chem. Soc.* **2000**, *122*, 2252.
- (10) Schmidt, V. A.; Alexanian, E. J. *Angew. Chem., Int. Ed.* **2010**, *49*, 4491.
- (11) Giglio, B. C.; Schmidt, V. A.; Alexanian, E. J. *J. Am. Chem. Soc.* **2011**, *133*, 13320.
- (12) Prévost, C. *Compt. Rend.* **1933**, *196*, 1129.
- (13) For recent reports that develop the Prévost reaction, see: (a) Emmanuvel, L.; Shaikh, T. M. A.; Sudalai, A. *Org. Lett.* **2005**, *7*, 2250. (b) Zhong, W.; Yang, J.; Meng, X.; Li, Z. *J. Org. Chem.* **2011**, *76*, 9997. (c) Fujita, M.; Wakita, M.; Sugimura, T. *Chem. Commun.* **2011**, *47*, 3983. (d) Zhong, W.; Liu, S.; Yang, J.; Meng, X.; Li, Z. *Org. Lett.* **2012**, *14*, 3336.
- (14) (a) Griffith, J. C.; Jones, K. M.; Picon, S.; Rawling, M. J.; Kariuki, B. M.; Campbell, M.; Tomkinson, N. C. O. *J. Am. Chem. Soc.* **2010**, *132*, 14409. (b) Picon, S.; Rawling, M.; Campbell, M.; Tomkinson, N. C. O. *Org. Lett.* **2012**, *14*, 6250. (d) Jones, K. M.; Tomkinson, N. C. O. *J. Org. Chem.* **2012**, *77*, 921.
- (15) For use of peroxide 1 in the oxidation of aromatic rings, see: Dragan, A.; Kubczyk, T. M.; Rowley, J. H.; Sproules, S.; Tomkinson, N. C. O. *Org. Lett.* **2015**, *17*, 2618.
- (16) Rawling, M. J.; Rowley, J. H.; Campbell, M.; Kennedy, A. R.; Parkinson, J. A.; Tomkinson, N. C. O. *Chem. Sci.* **2014**, *5*, 1777.
- (17) Woodward, R. B.; Brutcher, F. V. *J. Am. Chem. Soc.* **1958**, *80*, 209.
- (18) Zhong, Y.-L.; Shing, T. K. M. *J. Org. Chem.* **1997**, *62*, 2622.