


Selective Functionalization of Tetrathiafulvalene Using Mg- and Zn-TMP-Bases: Preparation of Mono-, Di-, Tri-, and Tetrasubstituted Derivatives

Julia Nafe, Florian Auras, Konstantin Karaghiosoff, Thomas Bein, and Paul Knochel*

Department of Chemistry, Ludwig-Maximilians-Universität, Butenandtstrasse 5-13, 81377 München, Germany

Supporting Information



ABSTRACT: The tetrathiafulvalene-scaffold (TTF) reacts selectively in allylation, acylation, arylation, halogenation, and thiolation reactions via magnesium or zinc derivatives that are obtained by a direct metalation with Mg- and Zn-TMP-bases (TMP = 2,2,6,6-tetramethylpiperidyl). This stepwise functionalization provides access to a range of new mono-, di-, tri-, and tetra-functionalized TTF-derivatives and allows for fine-tuning of their energy levels.

Pioneered by the discovery of Wudl,¹ tetrathiafulvalenes (TTFs) have emerged into an important class of organic materials² owing to their unique optical, electronic, and magnetic properties.³ Much effort has been made to functionalize the TTF-scaffold,⁴ including C–H arylation⁵ and direct metalations.⁶ Although the lithiation of TTF (**1**) allows for an efficient functionalization with robust substituents, the preparation of TTFs featuring sensitive functional groups remains difficult due to the high reactivity of the carbon–lithium bond. Recently, we have reported a set of new metallic amides⁷ derived from 2,2,6,6-tetramethylpiperidine (TMP-H), such as TMPPMgCl·LiCl (**2**),⁸ TMP₂Mg·2LiCl,⁹ TMPZnCl·LiCl (**3**),¹⁰ and TMP₂Zn·2MgCl₂·2LiCl (**4**).¹¹ These bases are highly kinetically active due to the steric hindrance of the TMP-moiety and tolerate a wide range of functional groups.

Herein, we report the selective functionalization of the TTF-skeleton using Mg- and Zn-TMP-bases (**2–4**) and demonstrate the potential of the resulting new polyfunctionalized TTFs as molecular semiconductors.

Thus, the magnesiation of TTF (**1**) is conveniently achieved by the addition of TMPPMgCl·LiCl (**2**, 1.1 equiv) at 25 °C within 1 h leading to the magnesiated-TTF (**5**). This magnesium derivative can be treated with various electrophiles (E¹–X) providing a range of TTF-derivatives of type **6** in 55–92% yield (Scheme 1 and Table 1).

Scheme 1. Preparation of Functionalized TTFs (6**) via the Magnesiation of TTF (**1**) with TMPPMgCl·LiCl (**2**)**




Table 1. Preparation of 4-Substituted TTF-Derivatives of Type **6 by Magnesiation of TTF (**1**) with TMPPMgCl·LiCl (**2**)**

entry	electrophile	product	yield (%) ^a
1	I ₂		6a: R = I 55
2	(BrCl ₂ C) ₂		6b: R = Br 67
3	PhSO ₂ Cl		6c: R = Cl 65
4	MeSO ₂ SMe		6d: R = SMe 89
5	Me ₂ N=CH ₂ OCOCF ₃		6e: R = CH ₂ NMe ₂ 55
6	DMF		6f: R = H 60
7	NC-CO ₂ Et		6g: R = OEt 72
8	tBuCOCl		6h: R = tBu 76 ^b
9	3-Cl-C ₆ H ₄ COCl		6i: R = 3-Cl-C ₆ H ₄ 83 ^c
10	R = 4-Cl		6j: R = 4-Cl 86 ^d
11	R = 4-CO ₂ Et		6k: R = 4-CO ₂ Et 87 ^d
12	R = 4-OMe		6l: R = 4-OMe 61 ^d
13	R = 3-CF ₃		6m: R = 3-CF ₃ 92 ^d

^aIsolated yield of analytically pure product. ^bCuCN·2LiCl solution was added. ^cPd-catalyzed acylation reaction using 10 mol % Pd(PPh₃)₄.

^dCross-coupling conditions: ZnCl₂ transmetalation, 3 mol % Pd(dba)₂, 6 mol % tfp.

Received: September 18, 2015

Published: October 15, 2015

The halogenation of **5** (iodolysis, bromination, and chlorination) proceeds in moderate yields (55–67% yield; Table 1, entries 1–3) due to the limited stability of the heterocyclic halides (**6a–c**). Methylthiolation of **5** is performed using MeSO_2SMe affording the thioether (**6d**) in 89% yield (entry 4). An aminomethylation of **5** using the iminium salt $\text{Me}_2\text{NCH}_2\text{OCOCC}_3$ ¹² provides the amine **6e** in 55% yield (entry 5). The acylation of **5** can be directly achieved by the addition of DMF or ethyl cyanoformate leading to the aldehyde **6f** and the ester **6g** in 60–72% yield (entries 6–7; for single-crystal X-ray structure analysis of **6g**, see the Supporting Information, Figure S3). A copper-catalyzed acylation with pivaloyl chloride provides the ketone **6h** (76% yield; entry 8), whereas a Pd-catalyzed Negishi-acylation¹³ furnishes the ketone **6i** in 83% yield (entry 9). The arylation of **5** is achieved by a transmetalation with zinc chloride followed by a Negishi cross-coupling using 3 mol % $\text{Pd}(\text{dba})_2$ (dba = dibenzylideneacetone) and 6 mol % tfp (tri-2-furylphosphine)¹⁴ as a catalyst and an aryl iodide as an electrophile. Interestingly, electron-withdrawing, as well as electron-donating, groups can be attached to the electron-rich TTF-core producing the corresponding arylated TTF-derivatives (**6j–m**) in 61–92% yield (entries 10–13).

The preparation of symmetrically and asymmetrically disubstituted TTF-derivatives of type **7** can be achieved by a selective metalation of various monofunctionalized TTF-derivatives. The presence of an electron-withdrawing substituent, such as a chlorine or an acyl or a carboethoxy group, on the TTF-core directs the second metalation to the adjacent position.¹⁵ In the case of a chloride (**6c**) or a carboethoxy group (**6g**), the metalation is best performed with $\text{TMPPMgCl}\cdot\text{LiCl}$ (2). Thus, the treatment of 4-chloro-TTF (**6c**) with $\text{TMPPMgCl}\cdot\text{LiCl}$ (2) at 0 °C (0.5 h) followed by the copper-catalyzed allylation reaction with ethyl 2-(bromomethyl)-acrylate¹⁶ furnishes the disubstituted TTF **7a** in 85% yield (Table 2, entry 1). After transmetalation with zinc chloride, a Pd-catalyzed Negishi acylation reaction with benzoyl chloride and Negishi cross-coupling reactions with aryl iodides are achieved leading to 4,5-disubstituted TTF-derivatives (**7b–d**) in 78–92% yield (entries 2–4). Magnesiation of the ester **6g** is performed using $\text{TMPPMgCl}\cdot\text{LiCl}$ (2; 1.1 equiv) at –20 °C (0.5 h). Subsequent trapping with ethyl cyanoformate leads to the diester-TTF (**7e**) in 65% yield (entry 5). The thioethers **7f–g** were obtained in 59–65% yield by quenching the magnesiated TTF-derivative of **6g** with PhSO_2SPh and MeSO_2SMe (entries 6–7). In the case of a benzoyl substituent (**7e**), a metalation with a magnesium base is too harsh and leads to unwanted side reactions. However, a zirconation with $\text{TMP}_2\text{Zn}\cdot\text{2MgCl}_2\cdot\text{2LiCl}$ (4; 1.1 equiv, THF, 0 °C, 0.5 h) leads to the corresponding zirconated TTF in quantitative yield. After iodolysis, the corresponding iodide **7h** is obtained in 83% yield (entry 8).

Trifunctionalized TTF-derivatives are readily prepared starting from the 4,5-diethyl ester-TTF **7e** using the mild base $\text{TMPZnCl}\cdot\text{LiCl}$ (3; 1.3 equiv, THF, –30 °C, 0.5 h). Reactions with iodine, copper-catalyzed allylation with 3-bromocyclohexene, or Negishi cross-coupling reactions with aryl iodides produce the expected trisubstituted TTF-derivatives (**8a–e**) in 66–94% yield (Scheme 2).


Fully functionalized TTFs of type **9** are prepared by the zirconation of **8c–e** using $\text{TMPZnCl}\cdot\text{LiCl}$ (3; 1.3 equiv) at 0 °C within 0.5 h. Trapping the zirconated-TTF-derivative of **8c** and **8d** with iodine gives tetrasubstituted TTFs **9a** and **9b** in 76–88% yield (Table 3, entries 1 and 2). Furthermore, Negishi

Table 2. Preparation of 4,5-Disubstituted TTF-Derivatives of Type **7 by Metalation of the Monosubstituted TTFs **6c**, **6g**, and **6i** with Mg- and Zn-TMP-Bases**

entry	substrate	electrophile	product	yield (%) ^a
1	6c	$\text{Br}-\text{CH}_2-\text{CH}_2-\text{CO}_2\text{Et}$	7a	85 ^{b,c}
2	6c	$\text{Cl}-\text{C}_6\text{H}_4-\text{CH}_2-\text{C}_6\text{H}_5$	7b	83 ^{b,d}
3	6c	$\text{R}=\text{OMe}$	7c: R = OMe	78 ^{b,e}
4	6c	$\text{R}=\text{CN}$	7d: R = CN	92 ^{b,e}
5	6g	$\text{NC}-\text{CO}_2\text{Et}$	7e	65 ^f
6	6g	PhSO_2SPh	7f: R = Ph	59 ^f
7	6g	MeSO_2SMe	7g: R = Me	65 ^f
8	6i	I_2	7h	83 ^g

^aIsolated yield of analytically pure product. ^b $\text{TMPPMgCl}\cdot\text{LiCl}$ (1.1 equiv, 0 °C) was used. ^c $\text{CuCN}\cdot\text{2LiCl}$ solution was added. ^d ZnCl_2 solution was added. Pd-catalyzed acylation reaction using 10 mol % $\text{Pd}(\text{PPh}_3)_4$. ^eCross-coupling conditions: ZnCl_2 transmetalation, 3 mol % $\text{Pd}(\text{dba})_2$, 6 mol % tfp. ^f $\text{TMPPMgCl}\cdot\text{LiCl}$ (1.1 equiv, –20 °C) was used. ^g $\text{TMP}_2\text{Zn}\cdot\text{2MgCl}_2\cdot\text{2LiCl}$ (1.1 equiv, 0 °C) was used.

Scheme 2. Preparation of Trisubstituted TTF-Derivatives of Type **8 Using $\text{TMPZnCl}\cdot\text{LiCl}$ (3)**



cross-coupling, copper-catalyzed allylation reaction and Pd-catalyzed Negishi acylation reaction provide tetrasubstituted TTFs **9c–h** in 53–90% yield (entries 3–8; for single-crystal X-ray structure analysis of **9c** and **9d**, see the Supporting Information, Figures S4 and S5).

All TTF-derivatives feature a strong UV absorption and exhibit a broad absorption band in the visible spectral region (SI, Figure S1). While the VIS absorption is weak for the unsubstituted TTF, it can be significantly enhanced by attaching substituents. In particular, the nonsymmetric 1- and 3-fold substituted TTFs possess systematically higher extinction coefficients in the VIS region compared to their symmetrically functionalized counterparts. On the other hand, the larger conjugated π -systems of the 2- and 4-fold substituted molecules, respectively, extend the absorption further into the red.

Table 3. Preparation of Fully Functionalized TTF-Derivatives of Type 9 Using $\text{TMPZnCl}\cdot\text{LiCl}$ (3)

entry	substrate	electrophile	product, yield (%) ^a
1	8c	I_2	9a: R = CO_2Et , 88
2	8d	I_2	9b: R = CN, 76
3	8c	$\text{R}^2 = \text{CO}_2\text{Et}$	9c: $\text{R}^1 = \text{R}^2 = \text{CO}_2\text{Et}$, 90 ^b
4	8d	$\text{R}^2 = \text{CN}$	9d: $\text{R}^1 = \text{R}^2 = \text{CN}$, 63 ^b
5	8e	$\text{R}^2 = \text{OMe}$	9e: $\text{R}^1 = \text{R}^2 = \text{OMe}$, 63 ^b
6	8c	$\text{Br}-\text{CH}_2-\text{CH}_2-\text{CO}_2\text{Et}$	9f: 85 ^c
7	8c	$\text{C}_6\text{H}_4-\text{COCl}$	9g: 85 ^d
8	8d	$\text{C}_6\text{H}_4-\text{COCl}$	9h: 80 ^d

^aIsolated yield of analytically pure product. ^bCross-coupling conditions: 3 mol % $\text{Pd}(\text{dba})_2$, 6 mol % tfp, $\text{CuCN}\cdot 2\text{LiCl}$ solution was added. ^cPd-catalyzed acylation reaction using 10 mol % $\text{Pd}(\text{PPh}_3)_4$.

We then studied the effect of the nature and number of substituents on the energy levels of the TTF-derivatives using differential pulse voltammetry (DPV).¹⁷ The highest occupied molecular orbital (HOMO) energies were extracted from scans in the oxidation direction (SI, Figure S2) and referenced to the oxidation of ferrocene ($E_{\text{fc/fc+}} = -4.80$ eV vs vacuum).

We found that indeed the substitution has a profound effect on the position of the HOMO, spanning a range of more than 200 meV (Figure 1). While a single $-\text{CO}_2\text{Et}$ or $-\text{Cl}$ substituent




Figure 1. Highest occupied molecular orbital energies of selected substituted TTF-derivatives measured by differential pulse voltammetry.

shifts the HOMO upward (SI, Figure S2k and S2l), the energy levels of all multisubstituted TTFs are significantly lower than those for bare TTF. For the series of aryl substituents, the HOMO can be further fine-tuned by adjusting the electron-accepting end group. For the molecules of type **8** and **9** we observe deeper HOMO levels as the acceptor strength is increased from $-\text{OMe}$ to $-\text{CO}_2\text{Et}$ and $-\text{CN}$. This approach for systematic fine-tuning of the energy levels allows for matching the work function of contact layers or electrodes, such as gold, silver, or indium tin oxide, which is of key importance for possible applications of these materials in electronic devices.

In summary, the use of Mg- and Zn-TMP bases allows for a selective synthesis of mono-, di-, tri-, and tetrafunctionalized TTF-derivatives. Due to the gentle reaction conditions a wide range of sensitive functional groups are tolerated leading to symmetrically, as well as asymmetrically, substituted polyfunctionalized TTFs. This novel synthesis protocol allows for fine-tuning of optical properties and energy levels and thus provides a strategy for realizing tailor-made molecular semiconductors.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: [10.1021/acs.orglett.Sb02715](https://doi.org/10.1021/acs.orglett.Sb02715).

Detailed experimental procedures and characterization data for new compounds (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: paul.knochel@cup.uni-muenchen.de.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank the SFB 749 for financial support. We thank Rockwood Lithium (Hoechst) and BASF SE (Ludwigshafen) for the generous gift of chemicals. The authors are grateful for funding from the German Science Foundation (DFG; Research Cluster NIM) and the Free State of Bavaria (Research Network SolTech). The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 321339.

■ REFERENCES

- (a) Wudl, F.; Wobschall, D.; Hufnagel, E. *J. Am. Chem. Soc.* **1972**, *94*, 670. (b) Wudl, F.; Kruger, A. A.; Kaplan, M. L.; Hutton, R. S. *J. Org. Chem.* **1977**, *42*, 768.
- (a) Bunz, U. H. F. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 969. (b) Bunz, U. H. F.; Rotello, V. M. *Angew. Chem., Int. Ed.* **2010**, *49*, 3268.
- (a) Gorgues, A.; Hudhomme, P.; Sallé, M. *Chem. Rev.* **2004**, *104*, 5151. (b) Sarhan, A. E.-W. A. O. *Tetrahedron* **2005**, *61*, 3889. (c) Lorcy, D.; Bellec, N.; Fourmigué, M.; Avarvari, N. *Coord. Chem. Rev.* **2009**, *293*, 1398.
- (a) Narita, M.; Pittman, C. U., Jr. *Synthesis* **1976**, *1976*, 489. (b) Becker, J. Y.; Bernstein, J.; Ellern, A.; Gershtenman, H.; Khodorkovsky, V. *J. Mater. Chem.* **1995**, *5*, 1557. (c) John, D. E.; Moore, J. A.; Bryce, M. R.; Batsanov, A. S.; Leech, M. A.; Howard, J. A.

- K. *Synthesis* **1998**, **1998**, 826. (d) John, D. E.; Moore, J. A.; Bryce, M. R.; Batsanov, A. S.; Leech, M. A.; Howard, J. A. K. *J. Mater. Chem.* **2000**, **10**, 1273. (e) Iyoda, M.; Hara, K.; Kuwatani, Y.; Nagase, S. *Org. Lett.* **2000**, **2**, 2217. (f) Batsanov, A. S.; Bryce, M. R.; Chesney, A.; Howard, J. A. K.; John, D. E.; Moore, A. J.; Wood, C. L.; Gershtenman, H.; Becker, J. Y.; Khodorkovsky, V. Y.; Ellern, A.; Bernstein, J.; Perepichka, I. F.; Rotello, V.; Gray, M.; Cuello, A. O. *J. Mater. Chem.* **2001**, **11**, 2181. (g) Bouguessa, S.; Gouasmia, A. K.; Golhen, S.; Ouahab, L.; Fabre, J. M. *Tetrahedron Lett.* **2003**, **44**, 9275. (h) Bendikov, M.; Wudl, F.; Perepichka, D. F. *Chem. Rev.* **2004**, **104**, 4891. (i) Riedl, J.; Horáková, P.; Šebest, P.; Pohl, R.; Havran, L.; Fojta, M.; Hocek, M. *Eur. J. Org. Chem.* **2009**, **2009**, 3519. (j) Mitamura, Y.; Yorimitsu, H.; Oshima, K.; Osuka, A. *Chem. Sci.* **2011**, **2**, 2017.
- (5) Mitamura, Y.; Yorimitsu, H.; Oshima, K.; Osuka, A. *Chem. Sci.* **2011**, **2**, 2017.
- (6) (a) Green, D. C. *J. Chem. Soc., Chem. Commun.* **1977**, 161. (b) Green, D. C. *J. Org. Chem.* **1979**, **44**, 1476. (c) Hsu, S.-Y.; Chiang, L. Y. *J. Org. Chem.* **1987**, **52**, 3444. (d) Jørgensen, M.; Bechgaard, K. *Synthesis* **1989**, **1989**, 207. (e) Fabre, J.-M.; Garin, J.; Uriel, S. *Tetrahedron Lett.* **1991**, **32**, 6407. (f) Otsubo, T.; Kochi, Y.; Bitoh, A.; Ogura, F. *Chem. Lett.* **1994**, 2047. (g) Iyoda, M.; Kuwatani, Y.; Hara, K.; Ogura, E.; Suzuki, H.; Ito, H.; Mori, T. *Chem. Lett.* **1997**, 599. (h) Kuduva, S. S.; Avarvari, N.; Fourmigüé, M. *J. Chem. Soc., Dalton Trans.* **2002**, 3686. (i) Ueno, R.; Fujino, D.; Yorimitsu, H.; Osuka, A. *Chem. - Eur. J.* **2013**, **19**, 7156.
- (7) (a) Mulvey, R. E.; Mongin, F.; Uchiyama, M.; Kondo, Y. *Angew. Chem., Int. Ed.* **2007**, **46**, 3802. (b) Haag, B.; Mosrin, M.; Ila, H.; Malakhov, V.; Knochel, P. *Angew. Chem., Int. Ed.* **2011**, **50**, 9794. (c) Klatt, T.; Markiewicz, J. T.; Sämann, C.; Knochel, P. *J. Org. Chem.* **2014**, **79**, 4253.
- (8) (a) Armstrong, D. R.; Clegg, W.; Dale, S. H.; Hevia, E.; Hogg, L. M.; Honeyman, G. W.; Mulvey, R. E. *Angew. Chem., Int. Ed.* **2006**, **45**, 3775. (b) Mulvey, R. E. *Organometallics* **2006**, **25**, 1060. (c) Krasovskiy, A.; Krasovskaya, V.; Knochel, P. *Angew. Chem., Int. Ed.* **2006**, **45**, 2958. (d) Lin, W.; Baron, O.; Knochel, P. *Org. Lett.* **2006**, **8**, 5673. (e) García-Álvarez, P.; Graham, D. V.; Hevia, E.; Kennedy, A. R.; Klett, J.; Mulvey, R. E.; O'Hara, C. T.; Weatherstone, S. *Angew. Chem., Int. Ed.* **2008**, **47**, 8079. (f) Mosrin, M.; Knochel, P. *Org. Lett.* **2008**, **10**, 2497. (g) Mosrin, M.; Boudet, N.; Knochel, P. *Org. Biomol. Chem.* **2008**, **6**, 3237. (h) Piller, F. M.; Knochel, P. *Synthesis* **2011**, **2011**, 1751. (i) Kunz, T.; Knochel, P. *Angew. Chem., Int. Ed.* **2012**, **51**, 1958. (j) Armstrong, D. R.; Crosbie, E.; Hevia, E.; Mulvey, R. E.; Ramsay, D. L.; Robertson, S. D. *Chem. Sci.* **2014**, **5**, 3031. (k) Léon, T.; Quinio, P.; Chen, Q.; Knochel, P. *Synthesis* **2014**, **46**, 1374. (l) Dagousset, G.; François, C.; Léon, T.; Blanc, R.; Sansiaume-Dagousset, E.; Knochel, P. *Synthesis* **2014**, **46**, 3133. (m) Nafe, J.; Herbert, S.; Auras, F.; Karaghiosoff, K.; Bein, T.; Knochel, P. *Chem. - Eur. J.* **2015**, **21**, 1102.
- (9) (a) Campbell, M.; Snieckus, V. *Encyclopedia of Reagents for Organic Synthesis*; Wiley: New York, 1995. (b) Clososki, G. C.; Rohrbogner, J.; Knochel, P. *Angew. Chem., Int. Ed.* **2007**, **46**, 7681. (c) Rohrbogner, J.; Clososki, G. C.; Knochel, P. *Angew. Chem., Int. Ed.* **2008**, **47**, 1503. (d) Boudet, N.; Dubbaka, S. R.; Knochel, P. *Org. Lett.* **2008**, **10**, 1715. (e) Rohrbogner, J.; Wagner, A. J.; Clososki, G. C.; Knochel, P. *Org. Synth.* **2009**, **86**, 374. (f) Despotopoulou, C.; Gignoux, C.; McConnell, D.; Knochel, P. *Synthesis* **2009**, **2009**, 3661. (g) Despotopoulou, C.; Klier, L.; Knochel, P. *Org. Lett.* **2009**, **11**, 3326. (h) Wunderlich, S. H.; Rohrbogner, C. J.; Unsinn, A.; Knochel, P. *Org. Process Res. Dev.* **2010**, **14**, 339. (i) Piller, F. M.; Bresser, T.; Fischer, M. K. R.; Knochel, P. *J. Org. Chem.* **2010**, **75**, 4365. (j) Bresser, T.; Knochel, P. *Angew. Chem., Int. Ed.* **2011**, **50**, 1914.
- (10) (a) Turck, A.; Plé, N.; Mojovic, L.; Quéguiner, G. *J. Heterocycl. Chem.* **1990**, **27**, 1377. (b) Mojovic, L.; Turck, A.; Plé, N.; Dorsy, M.; Ndzi, B.; Quéguiner, G. *Tetrahedron* **1996**, **52**, 10417. (c) Knochel, P.; Millot, N.; Rodriguez, A. L.; Tucker, C. E. *Org. React.* **2001**, **58**, 417. (d) Wunderlich, S.; Knochel, P. *Chem. Commun.* **2008**, 6387. (e) Mosrin, M.; Knochel, P. *Org. Lett.* **2009**, **11**, 1837. (f) Mosrin, M.; Monzon, G.; Bresser, T.; Knochel, P. *Chem. Commun.* **2009**, 5615. (g) Monzon, G.; Knochel, P. *Synlett* **2010**, **2010**, 304. (h) Bresser, T.; Monzon, G.; Mosrin, M.; Knochel, P. *Org. Process Res. Dev.* **2010**, **14**, 1299. (i) Duez, S.; Steib, A. K.; Manolikakes, S. M.; Knochel, P. *Angew. Chem., Int. Ed.* **2011**, **50**, 7686. (j) Klier, L.; Bresser, T.; Nigst, T. A.; Karaghiosoff, K.; Knochel, P. *J. Am. Chem. Soc.* **2012**, **134**, 13584. (k) Unsinn, A.; Ford, M. J.; Knochel, P. *Org. Lett.* **2013**, **15**, 1128. (l) Haas, D.; Mosrin, M.; Knochel, P. *Org. Lett.* **2013**, **15**, 6162. (m) Klatt, T.; Blümke, T. D.; Ganiek, M. A.; Knochel, P. *Synthesis* **2014**, **46**, 290. (n) Blanc, R.; Groll, K.; Bernhardt, S.; Stockmann, P. N.; Knochel, P. *Synthesis* **2014**, **46**, 1052. (o) Shen, Z.-L.; Sommer, K.; Knochel, P. *Synthesis* **2015**, **47**, 2617.
- (11) (a) Wunderlich, S. H.; Knochel, P. *Angew. Chem., Int. Ed.* **2007**, **46**, 7685. (b) Wunderlich, S. H.; Knochel, P. *Org. Lett.* **2008**, **10**, 4705. (c) Kienle, M.; Dunst, C.; Knochel, P. *Org. Lett.* **2009**, **11**, 5158. (d) Mosrin, M.; Knochel, P. *Chem. - Eur. J.* **2009**, **15**, 1468. (e) Dunst, C.; Kienle, M.; Knochel, P. *Synthesis* **2010**, **2010**, 2313.
- (12) (a) Ahond, A.; Cavé, A.; Kan-Fan, C.; Husson, H.-P.; De Rostolan, J.; de, J.; Potier, P. *J. Am. Chem. Soc.* **1968**, **90**, 5622. (b) Kinast, G.; Tietze, L.-F. *Angew. Chem., Int. Ed. Engl.* **1976**, **15**, 239. (c) Sweeney, J.; Perkins, G. *e-EROS Encyclopedia of Reagents in Organic Synthesis*; Wiley: Hoboken, NJ, 2005; DOI: [10.1002/047084289X.rt237.pub.2](https://doi.org/10.1002/047084289X.rt237.pub.2). (d) Corpet, M.; Gosmini, C. *Synthesis* **2014**, **46**, 2258. (e) Werner, V.; Ellwart, M.; Wagner, A. J.; Knochel, P. *Org. Lett.* **2015**, **17**, 2026.
- (13) (a) Negishi, E.; Valente, L. F.; Kobayashi, M. *J. Am. Chem. Soc.* **1980**, **102**, 3298. (b) Negishi, E. *Acc. Chem. Res.* **1982**, **15**, 340. (c) Negishi, E.; Bagheri, V.; Chatterjee, S.; Luo, F.-T.; Miller, J. A.; Stoll, A. T. *Tetrahedron Lett.* **1983**, **24**, 5181.
- (14) Farina, V.; Krishnan, B. *J. Am. Chem. Soc.* **1991**, **113**, 9585.
- (15) (a) Snieckus, V. *Chem. Rev.* **1990**, **90**, 879. (b) Whisler, M. C.; MacNeil, S.; Snieckus, V.; Beak, P. *Angew. Chem., Int. Ed.* **2004**, **43**, 2206.
- (16) Villieras, J.; Rambaud, M. *Synthesis* **1982**, **1982**, 924.
- (17) (a) Frost, J. M.; Faist, M. A.; Nelson, J. *Adv. Mater.* **2010**, **22**, 4881. (b) Calik, M.; Auras, F.; Salonen, L. M.; Bader, K.; Grill, I.; Handloser, M.; Medina, D. D.; Dogru, M.; Löbermann, F.; Trauner, D.; Hartschuh, A.; Bein, T. *J. Am. Chem. Soc.* **2014**, **136**, 17802. (c) Auras, F.; Li, Y.; Löbermann, F.; Döblinger, M.; Schuster, J.; Peter, L. M.; Trauner, D.; Bein, T. *Chem. - Eur. J.* **2014**, **20**, 14971.