


# Indium-Catalyzed [2 + 2] Cycloaddition of Allylsilanes to Internal Alkynes

Kazuhiro Okamoto,\* Takuya Shimbayashi, Eisuke Tamura, and Kouichi Ohe\*

Department of Energy and Hydrocarbon Chemistry, Graduate School of Engineering, Kyoto University, Katsura, Nishikyo-ku, Kyoto 615-8510, Japan


**S** Supporting Information

**ABSTRACT:** We have developed an indium-catalyzed [2 + 2] cycloaddition of allylsilanes to alkynes leading to selective cyclobutenone formation. The resulting cyclobutenones were readily converted to the oxidized products by Tamao–Fleming oxidation or the ring-opened products by an electrocyclic reaction.



Small-ring molecules are generally used as highly reactive synthetic intermediates because of their ring-strain energy.<sup>1</sup> Of such molecules, cyclobutenes are useful compounds because they readily undergo stereoregular electrocyclic ring-opening reactions to afford the corresponding 1,3-dienes.<sup>2</sup> The most direct and efficient method for preparing cyclobutenes is [2 + 2] cycloaddition of alkenes with alkynes that proceeds by photoirradiation,<sup>3</sup> transition metal catalysis,<sup>4</sup> or Lewis acid catalysis.<sup>5</sup> Of various alkenes, allylsilanes are often used as the allyl synthons in Lewis acid mediated allylation reactions such as Mukaiyama– or Sakurai–Hosomi-type reactions.<sup>6</sup> For [2 + 2] cycloaddition, there have been a few reports on Lewis acid mediated reactions of allylsilanes with electron-deficient alkynes. In such reports, there remain three points requiring improvement: (1) a stoichiometric Lewis acid such as TiCl<sub>4</sub> is required, (2) the scope of alkynes is limited to terminal alkynes, and (3) a further [2 + 2] cycloaddition may occur depending on the reaction conditions (Scheme 1),<sup>7</sup> in which the substrate scope is

**Scheme 1. Lewis Acid Mediated [2 + 2] Cycloaddition Reaction of Alkenes with Alkynes**



**Existing Problem**

- stoichiometric amount of Lewis acid is required
- applicable only for terminal alkynes
- difficult to suppress further cycloaddition

**Planned work**

- catalytic cycloaddition
- application to internal alkynes

limited because only terminal alkynes can be applied.<sup>7</sup> Therefore, it is important to develop a [2 + 2] cycloaddition reaction with internal alkynes to provide synthetic diversity.

Meanwhile, we have studied the direct addition reactions of alkenes to  $\alpha,\beta$ -unsaturated ketones controlled by the orbital–interaction effect of silicon substituents and have reported an acid-catalyzed direct conjugate addition of styrene derivatives to

$\beta$ -silylenones, as well as a scandium-catalyzed [3 + 2] cycloaddition reaction of allylsilanes to  $\beta$ -silylenones.<sup>8</sup> In this paper, we report an indium-catalyzed [2 + 2] cycloaddition reaction of allylsilanes to internal conjugated alkynes.

First, the reaction of alkyne **1a** and allyltriisopropylsilane **2a** in 1,2-dichloroethane was performed in the presence of various acid catalysts. Although Me<sub>3</sub>SiOTf and HOTf were effective in the conjugate alkenylation that we have previously reported, they exhibited no catalytic activity, in the present reaction (Table 1, entries 1 and 2). Scandium(III) triflate, which is the best catalyst for the previous report on [3 + 2] cycloaddition, showed high catalytic activity, but gave only a 7%

**Table 1. Lewis Acid Catalyzed [2 + 2] Cycloaddition of Allylsilane **2a** with Alkyne **1a**<sup>a</sup>**

entry	catalyst	temp (°C)	<b>1a</b> conv (%) <sup>b</sup>	<b>3aa</b> yield (%) <sup>b</sup>
1	Me <sub>3</sub> SiOTf	25	0	0
2	HOTf	25	6	0
3	Sc(OTf) <sub>3</sub>	25	66	7
4	Al(OTf) <sub>3</sub>	25	71	17
5	Ga(OTf) <sub>3</sub>	25	79	38
6	In(OTf) <sub>3</sub>	25	78	61
7	InCl <sub>3</sub>	25	66	47
8	InBr <sub>3</sub>	0	96	81 (73) <sup>c</sup>
9	InI <sub>3</sub>	−40	98	58
10	In(NTf <sub>2</sub> ) <sub>3</sub>	−40	84	53
11	TiCl <sub>4</sub> (1.0 equiv)	0	100	16
12	Me <sub>2</sub> AlCl (1.0 equiv)	0	2	2

<sup>a</sup>Reactions were carried out with alkyne **1a** (0.20 mmol), allylsilane **2a** (0.40 mmol), and catalyst (20  $\mu$ mol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl (1.5 mL).  
<sup>b</sup>Determined by <sup>1</sup>H NMR. <sup>c</sup>Isolated yield.


**Received:** October 23, 2015

**Published:** November 19, 2015

yield of the target cyclobutene. As a result of screening other metal(III) triflate salts, indium(III) triflate was proven to be the most effective catalyst in the [2 + 2] cycloaddition reaction (entries 4–6). Then, we examined other indium salts and suitable reaction temperatures (entries 7–10). While indium(III) chloride showed low catalytic activity, indium(III) bromide served as the most effective catalyst at 0 °C and gave the cycloadduct in 73% isolated yield. In the case of indium(III) iodide or bis(triflyl)imide, yields of the product were lower even at low temperature (–40 °C) because of their high reactivity. Other typical Lewis acids, such as TiCl<sub>4</sub> and Me<sub>2</sub>AlCl, gave the cycloadduct in only very low yields even when the stoichiometric amount of these reagents was used.<sup>9</sup>


Various combinations of alkynes and allylsilanes afforded the corresponding cyclobutenones in high yields under the optimized reaction conditions (Scheme 2). As a terminal substituent of alkynes, the cyclohexyl group at R<sup>1</sup> could be altered to both *n*-butyl and *tert*-butyl groups (3ba and 3ca).<sup>10</sup> These results show the suitability of various alkyl groups, including primary, secondary, and tertiary alkyl groups. The cycloadducts were obtained in good yields when the alkynes possess various aromatic and heteroaromatic rings such as benzene, naphthalene, thiophene, and benzofuran (3da–3ga). As a silyl substituent of allylsilanes, the *tert*-butyldiphenylsilyl group (*t*-BuPh<sub>2</sub>Si) and triphenylsilyl group (Ph<sub>3</sub>Si), which can be used for further transformations such as Tamao–Fleming oxidation, could also be applied (3db and 3dc).<sup>11</sup> The scope of alkynes was also expanded to β-silylalkynes. The reaction of alkyne 1h with allylsilane 2b gave the cycloadduct 3hb selectively, while the reaction of alkyne 1i and allylsilane 2a underwent [3 + 2] cycloaddition involving the rearrangement of a silyl group in addition to the main [2 + 2] cycloaddition reaction, resulting in a mixture of two products, 3ia and 4ia, in a total yield of 87% with a ratio of ca. 1.4:1. In our previous report, the β-silicon substituents on the α,β-unsaturated carbonyl compounds exerted a remarkable influence on the product selectivity and also affected the selectivity in the present reaction.<sup>8,12</sup>

It is worth noting that the selectivity of the cycloaddition pathway was switched by using a stoichiometric amount of TiCl<sub>4</sub>, and the [3 + 2] cycloaddition product 4ia was selectively obtained in 67% yield (eq 1).<sup>13</sup>



Because the cyclobutenones obtained in the present reaction have large strain energies, they are expected to undergo thermal electrocyclic reactions to afford the corresponding ring-opened products. The reaction of cycloadduct 3aa for 2 h in refluxing toluene afforded the dienone 5aa in 22% yield together with silylenolate 6aa in 12% yield (eq 2). Silylenolate 6aa was considered to be formed by the thermal sigmatropic rearrangement of 5aa involving both C–Si bond cleavage and O–Si bond formation. Increasing the reaction time to 24 h resulted in


## Scheme 2. Indium-Catalyzed [2 + 2] Cycloaddition of Allylsilanes 2 to Alkynes 1<sup>a</sup>



<sup>a</sup>Reactions were carried out with alkyne 1 (0.20 mmol), allylsilane 2 (0.40 mmol), and InBr<sub>3</sub> (20 μmol) in ClCH<sub>2</sub>CH<sub>2</sub>Cl (1.5 mL). Isolated yields are shown. <sup>b</sup>Reactions were performed at 40 °C for 42 h. <sup>c</sup>Determined by the <sup>1</sup>H NMR spectrum of the mixture of 3ia and 4ia after column chromatography.

the selective formation of 6aa in high yield. The silyl moiety of enolate 6aa was readily removed by treatment with tetrabutylammonium fluoride, dienone 7aa being obtained quantitatively as a single geometrical isomer (eq 3).

The silicon substituents on the cyclobutenones were oxidatively transformed as mentioned above. Cyclobutenone 3ab, which possesses a SiPh<sub>2</sub>(*t*-Bu) group, was first treated with fluoroboric acid followed by aqueous hydrogen peroxide as an oxidant and KHF<sub>2</sub> to give alcohol 8ab in good yield without any side reaction on the cyclobutene moiety (eq 4). It is of great interest that the generally highly reactive cyclobutene ring was tolerated under such reaction conditions.



In summary, we have developed an indium-catalyzed [2 + 2] cycloaddition of allylsilanes and electron-deficient alkynes, which would serve as an efficient method for the synthesis of cyclobutenones. The selectivity of the cycloaddition mode was switched by changing the silyl substituents at the  $\beta$ -position of alkynes and the catalyst. The usefulness of the present reaction was also demonstrated by the geometry-selective thermal electrocyclic ring-opening reaction of the cyclobutenones, as well as the oxidative transformation of the silicon substituents.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.5b03065.

Experimental procedures and compound characterization data (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: kokamoto@scl.kyoto-u.ac.jp.

\*E-mail: ohe@scl.kyoto-u.ac.jp.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI (Grant-in-Aid for Scientific Research (C); Grant Number 26410119).

## ■ REFERENCES

- (1) Anslyn, E. V.; Dougherty, D. A. *Strain and Stability. Modern Physical Organic Chemistry*; University Science Books: Sausalito, CA, 2006; pp 65–144.
- (2) For reviews on synthetic utilities of cyclobutenes, see: (a) Namyslo, J. C.; Kaufmann, D. E. *Chem. Rev.* **2003**, *103*, 1485. (b) Murakami, M.; Miyamoto, Y.; Hasegawa, M.; Usui, I.; Matsuda, T. *Pure Appl. Chem.* **2006**, *78*, 415.
- (3) (a) Pappas, S. P.; Pappas, B. C.; Portnoy, N. A. *J. Org. Chem.* **1969**, *34*, 520. (b) Serve, M. P.; Rosenberg, H. M. *J. Org. Chem.* **1970**, *35*, 1237. (c) Bloomfield, J. J.; Owsley, D. C. *J. Am. Chem. Soc.* **1971**, *93*, 782. (d) Koft, E. R.; Smith, A. B., III. *J. Am. Chem. Soc.* **1984**, *106*, 2115. (e) Serebryakov, E. P. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **1984**, *33*, 120.
- (4) (a) Lautens, M.; Klute, W.; Tam, W. *Chem. Rev.* **1996**, *96*, 49. (b) Hilt, G.; Paul, A.; Treutwein, J. *Org. Lett.* **2010**, *12*, 1536. (c) López-Carrillo, V.; Echavarren, A. M. *J. Am. Chem. Soc.* **2010**, *132*,

9292. (d) Motokura, K.; Nakayama, K.; Miyaji, A.; Baba, T. *ChemCatChem* **2011**, *3*, 1419. (e) Nishimura, A.; Ohashi, M.; Ogoshi, S. *J. Am. Chem. Soc.* **2012**, *134*, 15692. (f) Sakai, K.; Kochi, T.; Kakiuchi, F. *Org. Lett.* **2013**, *15*, 1024.

(5) (a) Snider, B. B. *J. Org. Chem.* **1976**, *41*, 3061. (b) Snider, B. B.; Brown, L. A.; Conn, R. S. E.; Killinger, T. A. *Tetrahedron Lett.* **1977**, *18*, 2831. (c) Clark, R. D.; Untch, K. G. *J. Org. Chem.* **1979**, *44*, 248. (d) Snider, B. B.; Rodini, D. J.; Conn, R. S. E.; Sealfon, S. *J. Am. Chem. Soc.* **1979**, *101*, 5283. (e) Fienemann, H.; Hoffmann, H. M. R. *J. Org. Chem.* **1979**, *44*, 2802. (f) Rosenblum, M.; Scheck, D. *Organometallics* **1982**, *1*, 397. (g) Quendo, A.; Rousseau, G. *Tetrahedron Lett.* **1988**, *29*, 6443. (h) Faron, K. L.; Wulff, W. D. *J. Am. Chem. Soc.* **1988**, *110*, 8727. (i) Ishihara, K.; Fushimi, M. *J. Am. Chem. Soc.* **2008**, *130*, 7532.

(6) For reviews, see: (a) Hosomi, A. *Acc. Chem. Res.* **1988**, *21*, 200. (b) Fleming, I.; Dunoguès, J.; Smithers, R. *Org. React.* **1989**, *37*, 57. (c) Fleming, I. *Comprehensive Organic Synthesis*, 2nd ed.; Elsevier: 1991; Vol. 2, p 56310.1016/B978-0-08-052349-1.00041-X. (d) Miura, K.; Hosomi, A. *Allylsilanes, Allenylsilanes, and Propargylsilanes. In Main Group Metals in Organic Synthesis*; Yamamoto, H., Oshima, K., Eds.; Wiley-VCH: Weinheim, 2006; pp 489–533.

(7) (a) Hojo, M.; Tomita, K.; Hirohara, Y.; Hosomi, A. *Tetrahedron Lett.* **1993**, *34*, 8123. (b) Monti, H.; Audran, G.; Monti, J.-P.; Léandri, G. *Synlett* **1994**, *1994*, 403. (c) Monti, H.; Audran, G.; Léandri, G.; Monti, J.-P. *Tetrahedron Lett.* **1994**, *35*, 3073. (d) Knölker, H.-J.; Baum, G.; Graf, R. *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 1612. (e) Hojo, M.; Murakami, C.; Nakamura, S.; Hosomi, A. *Chem. Lett.* **1998**, 331. (f) Inanaga, K.; Takasu, K.; Ihara, M. *J. Am. Chem. Soc.* **2005**, *127*, 3668. (g) Takasu, K.; Hosokawa, N.; Inanaga, K.; Ihara, M. *Tetrahedron Lett.* **2006**, *47*, 6053.

(8) (a) Okamoto, K.; Tamura, E.; Ohe, K. *Angew. Chem., Int. Ed.* **2013**, *52*, 10639. (b) Okamoto, K.; Tamura, E.; Ohe, K. *Angew. Chem., Int. Ed.* **2014**, *53*, 10195.

(9) We attempted further cycloaddition of cyclobutenone **3hb** with allylsilane **2a**, but failed to obtain the desired product.

(10) In the reaction of terminal alkynone (R<sup>1</sup> = H), complex mixtures were obtained and the target cycloadduct was not detected.

(11) The reaction of alkynone **1a** with less hindered allyltrimethylsilane gave the [2 + 2] cycloadduct in low yield (6%).

(12) Simple alkenes such as styrene did not cause the cycloaddition with alkynones under the present conditions.

(13) [3 + 2]-Cycloaddition of allylsilanes with enones: (a) Knölker, H.-J.; Jones, P. G.; Pannek, J.-B. *Synlett* **1990**, *1990*, 429. (b) Knölker, H.-J.; Foitzik, N.; Graf, R.; Pannek, J.-B. *Tetrahedron* **1993**, *44*, 9955. (c) Knölker, H.-J.; Foitzik, N.; Goemann, H.; Graf, R.; Jones, P. G.; Wanzl, G. *Chem. - Eur. J.* **1997**, *3*, 538. (d) Knölker, H.-J. *J. Prakt. Chem./Chem.-Ztg.* **1997**, *339*, 304. For a review, see: (e) Schmidt, A. W.; Knölker, H.-J. *Synlett* **2010**, *2010*, 2207.