


Total Synthesis of the Antitumor Macrolides, (+)-Brefeldin A and 4-Epi-Brefeldin A from D-Glucose: Use of the Padwa Anionic Allenylsulfone [3 + 2]-Cycloadditive Elimination To Construct Trans-Configured Chiral Cyclopentane Systems

Ziyue Xiong and Karl J. Hale*

The School of Chemistry & Chemical Engineering and the Centre for Cancer Research and Cell Biology (CCRCB), The Queen's University Belfast, Stranmillis Road, Belfast BT9 5AG, Northern Ireland, United Kingdom

S Supporting Information

ABSTRACT: A new synthesis of (+)-brefeldin A is reported via Padwa allenylsulfone [3 + 2]-cycloadditive elimination. Cycloadduct **13** was initially elaborated into iodide **27**, which, following treatment with Zn, gave aldehyde **28** whose C(9) stereocenter was epimerized. Further elaboration into enoate **38** and Julia–Kocienski olefination with **5** subsequently afforded **39**, which was deprotected at C(1) and O(15). Yamaguchi macrolactonization of the *seco*-acid thereafter afforded a macrocycle that underwent O-desilylation and inversion at C(4) to give (+)-brefeldin A following deprotection.



The structurally complex antitumor macrolide, (+)-brefeldin A,¹ has occupied an almost iconic position within the field of stereocontrolled natural product total synthesis,² due to the significant challenges it poses for complex chiral cyclopentane ring construction. In this connection, and following a recent successful deployment of the Padwa allenylsulfone anionic [3 + 2]-cycloadditive–elimination³ in the synthesis of (–)-echinosporin,⁴ we became interested in evaluating whether this novel cyclopentene ring-assembly method might prove useful for the stereoselective construction of chiral bicyclic cyclopentanoids with a *trans*-ring junction, and in this connection, (+)-brefeldin A immediately sprung to mind as a target.

(+)-Brefeldin A has elicited considerable medicinal interest over the years,⁵ due to the fact that its water-soluble 7-*N,N*-dimethylglycinate pro-drug, breflate, was reported to be a powerful inhibitor of human melanoma xenograft growth in mice.⁶ Despite these early exciting findings, subsequent more detailed pharmacological evaluation of breflate and other brefeldin A pro-drugs at higher dosages (20 mg/kg) did eventually reveal that they could bring about seizures in mice and cause noticeable neurodegeneration.⁷ Ultimately, these observations led to molecules of the brefeldin class not proceeding into human clinical development. It is now believed that much of the toxicity of (+)-brefeldin A derives from its blockade of the interaction between the adenosine diphosphate ribosylation factor 1 (Arf1)–GDP (guanosine diphosphate) complex with the Sec7 domains of various Arf–GTPase exchange factors, which interferes with their normal functioning.^{7–9} Despite these problems, work has continued on the synthesis of new brefeldin A analogues,⁵ with many teams

retaining the hope that they might identify a structurally modified congener that will have an improved activity/toxicity profile. It was with such thoughts in mind, that we too commenced synthetic efforts on (+)-brefeldin A, and herein, we now report a new, fully stereocontrolled, enantioselective synthesis.


In our original retrosynthetic plan for (+)-brefeldin A (**1**) (Scheme 1), we sought to access **1** from the *seco*-acid **2** by regioselective macrolactonization. Compound **2** would emerge from **3** by cleavage of all the protecting groups, while **3** would derive from **6** by the successful implementation of *E*-olefin cross-metathesis and Julia/Kocienski olefination¹⁰ tactics on **6** and **4**, respectively. Aldehyde **6** would possibly emerge from **8** by pyrrolidine-induced epimerization, while **8** might be securable from the pyranoside **9** by Vasella reductive ring cleavage.¹¹ Further retrosynthetic tracing of **9** to **12** suggested compounds **10** and **11** as intermediates, with **11** emerging from a sulfone carbanion oxidation on **12** and **10** from a ketone reduction. A double Mitsunobu inversion¹² on **10** with benzoic acid and an O-desilylation and iodination would thereafter complete the route to **9**. The requisite Padwa anionic [3 + 2]-cycloadduct **13** had already been prepared from **16** and **17** during our development of a new pathway to (–)-echinosporin,⁴ and so now we envisioned converting **13** into **12** by a simple catalytic hydrogenation reaction.

With this summary of our strategy in mind, we repeated the work of Flasz and Hale⁴ to acquire cycloadduct **13** in the previously reported 56% yield (Scheme 2). The alkene in **13** was

Received: July 9, 2016

Published: August 19, 2016

Scheme 4. Our Synthesis of the New Alkenyl Aldehyde 29




aldehyde also did not occur; a 3 h reaction time typically minimized this event and gave **28** in 95% yield. A variety of bases were then investigated to perform the desired epimerization of **28** into **29**, but pyrrolidine soon emerged as the reagent of choice, giving a high yield of product with high stereoselectivity.


A fresh disappointment beset us when we discovered that the alkene in **29** was not a productive partner in olefin cross-metathesis with excess methyl acrylate to obtain **30** (Scheme 4). Given this undesired outcome, we decided to investigate whether a Wittig reaction might prove useful for enoate elaboration, but clearly, the adoption of this tactic would require reduction of the C(9)-aldehyde and protection of the resulting alcohol to enable the C(3)-aldehyde to be oxidatively unveiled. There was also the issue of C(4) inversion and macrolactonization to contend with much later in the synthesis. After due consideration, an orthogonal silyl ether protecting group strategy was eventually adopted, with aldehyde **29** being converted into differentially protected tri-*O*-silyl ether **33** (Scheme 5).

A four-step protocol accomplished this task (Scheme 5). In this, NaBH_4 reduction of **29** delivered a primary alcohol that underwent *O*-silylation with TBSCl and imidazole. Alkene **31** then had its *O*-benzoate reductively removed with $i\text{-Bu}_2\text{AlH}$ at low temperature. The resulting alcohol **32** was then *O*-silylated with TBDPSCl and imidazole in CH_2Cl_2 at rt. Initially, the use of cat. OsO_4 and NaIO_4 was pursued for the oxidative cleavage of alkene **33**, to obtain **35**, but this gave rise to a complex reaction mixture. In the end, the alkene of **33** was best dihydroxylated under standard Upjohn conditions with cat. OsO_4 and excess *N*-methylmorpholine *N*-oxide.¹⁵ This afforded a single diol **34**, whose stereochemistry was tentatively assigned as *anti*, using Kishi's empirical rule.¹⁶ This diol was then oxidatively cleaved with $\text{Pb}(\text{OAc})_4$, and the resulting aldehyde **35** Wittig olefinated to obtain enoate **36** as a single geometric isomer. Selective *O*-desilylation of **36** with cat. PPTS/EtOH at rt over 60 h cleanly gave alcohol **37**, which was converted into the aldehyde **38** by TEMPO/ $\text{PhI}(\text{OAc})_2$ oxidation. Julia/Kocienski olefination¹⁰ of **38** with the *N*-phenyltetrazolylsulfone **5** furnished the all (*E*)-alkene **39** as a single stereoisomer in 68% yield. Our new abridged synthesis of **5**^{2e} from **45**¹⁷ is presented in Scheme 6.

Ceric ammonium nitrate cleavage of the PMP group¹⁸ from **39** then ensued; the methyl ester was likewise cleaved with aq LiOH.

Scheme 5. Completion of Our Syntheses of (+)-Brefeldin A (**1**) and 4-Epi-Brefeldin A (**44**) from Alkenyl Aldehyde 29

A Yamaguchi macrolactonization¹⁹ of the *seco*-acid **40** was now accomplished at high dilution, as described in Scheme 5. Macrolactone **41** was formed in excellent yield, in what was a very clean reaction. Selective C(4) *O*-desilylation of **41** could subsequently be accomplished cleanly and rapidly with 5 equiv of $n\text{-Bu}_4\text{NF}$ in THF over 1 h. The resulting alcohol was then subjected to Mitsunobu inversion with Ph_3P /DIAD and *p*-nitrobenzoic acid.^{12c} The desired product **43** was isolated pure in 65% yield; elimination product **42** was also isolated in 17% yield by SiO_2 flash chromatography. With pure **43** in hand, (+)-brefeldin A was readily accessed after HF-pyridine-induced

Scheme 6. Synthesis of the *N*-Phenyltetrazolylsulfone 5

O-desilylation and brief treatment (0.5 h) with K_2CO_3 (1 equiv) in MeOH at 0 °C. Following final purification by SiO_2 flash chromatography, the natural product was obtained in 70% yield for the two steps and was spectroscopically identical with the spectra reported for (+)-brefeldin A in the literature in CD_3OD ^{2f} and CDCl_3 .²⁰ Cleavage of the *O*-silyl protecting groups from 41 with *n*-Bu₄NF in THF afforded 4-epi-brefeldin A,^{2h} as well.

In summary, a new total synthesis of (+)-brefeldin A was achieved from the Padwa [3 + 2]-cycloadduct 13. The utility of the latter cycloaddition was demonstrated for the preparation of trans-configured chiral cyclopentanes. New types of brefeldin analogues should now prove accessible.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b02002.

Full experimental procedures for all steps, as well as copies of the IR, HMRS, and ¹H/¹³C NMR spectra of every intermediate (PDF)

■ AUTHOR INFORMATION

Corresponding Author

*E-mail: k.j.hale@qub.ac.uk.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank QUB for a studentship and for funding this work.

■ REFERENCES

- (1) (a) Singleton, V. L.; Bohonos, N.; Ullstrup, A. J. *Nature* **1958**, *181*, 1072. (b) Weber, H. P.; Hauser, D.; Sigg, H. P. *Helv. Chim. Acta* **1971**, *54*, 2763.
- (2) For previous (+)-brefeldin A total syntheses in reverse chronological order, see: (a) Fuchs, M.; Furstner, A. *Angew. Chem., Int. Ed.* **2015**, *54*, 3978. (b) Wu, Y.; Gao, J. *Org. Lett.* **2008**, *10*, 1533. (c) Seo, S.-Y.; Jung, J.-K.; Paek, S.-M.; Lee, Y.-S.; Kim, S.-H.; Suh, Y.-G. *Tetrahedron Lett.* **2006**, *47*, 6527. (d) Wu, Y.; Shen, X.; Yang, Y.-Q.; Hu, Q.; Huang, J.-H. *Tetrahedron Lett.* **2004**, *45*, 199. (e) Trost, B. M.; Crawley, M. L. *J. Am. Chem. Soc.* **2002**, *124*, 9328. (f) Kim, D.; Lee, J.; Shim, P. J.; Lim, J. I.; Jo, H.; Kim, J. *Org. Chem.* **2002**, *67*, 764. (g) Suh, Y.-G.; Jung, J.-K.; Seo, S.-Y.; Min, K.-H.; Shin, D.-Y.; Lee, Y.-S.; Kim, S.-H.; Park, H.-J. *J. Org. Chem.* **2002**, *67*, 4127. (h) Wang, Y.; Romo, D. *Org. Lett.* **2002**, *4*, 3231. (i) Haynes, R. K.; Lam, W. W.-L.; Yeung, L.-L.; Williams, I. D.; Ridley, A. C.; Starling, S. M.; Vonwiller, S. C.; Hambley, T. W.; Lelandais, P. J. *Org. Chem.* **1997**, *62*, 4552. (j) Carnell, A. J.; Casy,

- G.; Gorins, G.; Kompany-Saeid, A.; McCague, R.; Olivo, H. F.; Roberts, S. M.; Willetts, A. J. *J. Chem. Soc., Perkin Trans. 1* **1994**, 3431. (k) Miyaoka, H.; Kajiwara, M. *J. Chem. Soc., Chem. Commun.* **1994**, 483. (l) Solladié, G.; Lohse, O. *J. Org. Chem.* **1993**, *58*, 4555. (m) Taber, D. F.; Silverberg, L. J.; Robinson, E. D. *J. Am. Chem. Soc.* **1991**, *113*, 6639. (n) Nokami, J.; Ohkura, M.; Dan-Oh, Y.; Sakamoto, Y. *Tetrahedron Lett.* **1991**, *32*, 2409. (o) Corey, E. J.; Carpino, P. *Tetrahedron Lett.* **1990**, *31*, 7555. (p) Hatakeyama, S.; Sugawara, K.; Kawamura, M.; Takano, S. *Synlett* **1990**, 691. (q) Gais, H.-J.; Lukas, K. L. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 142. (r) Le Drian, C.; Greene, A. E. *J. Am. Chem. Soc.* **1982**, *104*, 5473. (s) Corey, E. J.; Wollenberg, R. H.; Williams, D. R. *Tetrahedron Lett.* **1977**, *18*, 2243.

- (3) (a) Padwa, A.; Yeske, P. E. *J. Am. Chem. Soc.* **1988**, *110*, 1617. (b) Padwa, A.; Yeske, P. E. *J. Org. Chem.* **1991**, *56*, 6386. (c) Padwa, A.; Watterson, S. H.; Ni, Z. *J. Org. Chem.* **1994**, *59*, 3256. (d) Nunez, A., Jr.; Martin, M. R.; Fraile, A.; Garcia Ruano, J. L. *Chem. - Eur. J.* **2010**, *16*, 5443.

- (4) Flasz, J. T.; Hale, K. J. *Org. Lett.* **2012**, *14*, 3024.

- (5) (a) Anadu, N. O.; Davisson, V. J.; Cushman, M. J. *Med. Chem.* **2006**, *49*, 3897. (b) Seehafer, K.; Rominger, F.; Helmchen, G.; Langhans, M.; Robinson, D. G.; Ozata, B.; Brugger, B.; Strating, J. R. P. M.; van Kuppeveld, F. J. N.; Klein, C. D. *J. Med. Chem.* **2013**, *56*, 5872. (c) Forster, S.; Persch, E.; Tverskoy, O.; Rominger, F.; Helmchen, G.; Klein, C.; Gonen, B.; Brugger, B. *Eur. J. Org. Chem.* **2011**, *2011*, 878.

- (6) (a) Carter, C. A.; Waud, W. R.; Plowman, J. *Proc. Am. Assoc. Cancer Res.* **1995**, *36*, 386. (b) Phillips, L. R.; Wolfe, T. L.; Malspeis, L.; Supko, J. G. *J. Pharm. Biomed. Anal.* **1998**, *16*, 1301.

- (7) Tosca, P.; Merrill, J.; Ryan, M.; Knostman, K.; Johnson, J.; Peggins, J.; Tomaszewski, K. *Cancer Res.* **2009**, *69*, S102.

- (8) Peyroche, A.; Antonny, B.; Robineau, S.; Acker, J.; Cherfils, J.; Jackson, C. L. *Mol. Cell* **1999**, *3*, 275. (b) Chardin, P.; McCormick, F. *Cell* **1999**, *97*, 153.

- (9) Colanzi, A.; Grimaldi, G.; Catara, G.; Valente, C.; Cericola, C.; Liberali, P.; Ronci, M.; Lalioti, V. S.; Bruno, A.; Beccari, A. R.; Urbani, A.; De Flora, A.; Nardini, M.; Bolognesi, M.; Luini, A.; Corda, D. *Proc. Natl. Acad. Sci. U. S. A.* **2013**, *110*, 9794.

- (10) Blakemore, P. R.; Cole, W. J.; Kocienski, P. J.; Morley, A. *Synlett* **1998**, 1998, 26.

- (11) Bernet, B.; Vasella, A. *Helv. Chim. Acta* **1979**, *62*, 1990.

- (12) (a) Mitsunobu, O. *Synthesis* **1981**, 1981, 1. (b) Hughes, D. L. *Org. React.* **1992**, *42*, 335. (c) Martin, S. F.; Dodge, J. A. *Tetrahedron Lett.* **1991**, *32*, 3017.

- (13) Davis, F. A.; Sheppard, A. C. *Tetrahedron* **1989**, *45*, 5703.

- (14) Barton, D. H. R.; McCombie, S. W. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1574.

- (15) VanRheenen, V.; Kelly, R. C.; Cha, D. Y. *Tetrahedron Lett.* **1976**, *17*, 1973.

- (16) Cha, J. K.; Christ, W. J.; Kishi, Y. *Tetrahedron* **1984**, *40*, 2247.

- (17) Chaudhari, D. A.; Kattanguru, P.; Fernandes, R. A. *RSC Adv.* **2015**, *5*, 42131.

- (18) Fukuyama, T.; Laird, A. A.; Hotchkiss, L. M. *Tetrahedron Lett.* **1985**, *26*, 6291.

- (19) Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.

- (20) Glaser, R.; Shiftan, D.; Froimowitz, M. *Magn. Reson. Chem.* **2000**, *38*, 274.