

Borane-Catalyzed Selective Hydrosilylation of Internal Ynamides Leading to β -Silyl (Z)-Enamides

Youngchan Kim, †,‡,§ Ramesh B. Dateer,‡,†,§ and Sukbok Chang*,‡,†©

Supporting Information

ABSTRACT: We have developed a borane-catalyzed regioand stereoselective hydrosilylation of internal ynamides for the first time. The scope of ynamide substrates and silane reactants was broad under mild reaction conditions, affording synthetically versatile β -silyl (Z)-enamide products. The observed stereoselectivity was reasoned to be due to the β -silicon effect on a postulated ketene iminium intermediate.

$$R = -[N] \xrightarrow{\text{Cat.} \\ B(C_6F_5)_3} \text{ Phose } [N] \xrightarrow{\text{Phose } \\ Si = H} [N] \xrightarrow{\text{Si}} [N]$$

$$\beta \text{-silicon effect} \xrightarrow{\text{Si}} [N]$$

$$\bullet \text{ The first example of ynamide hydrosilylation}$$

$$\bullet \text{ Synthetically versatile } \beta \text{-silyl } (Z) \text{-enamide products}$$

$$\bullet \text{ Transition metal-free}$$

$$\bullet \text{ High regio- and stereoselectivity}$$

Vinylsilanes are versatile compounds that serve as an important synthetic building unit. In fact, they are frequently utilized in a broad range of transformations, including Tamao—Fleming oxidation, electrophilic substitution, and crosscoupling reactions. Among various preparative methods for vinylsilanes, hydrosilylation of alkynes is one of the most straightforward, convenient, and atom-economical procedures. While transition-metal catalysts such as platinum, ruthenium, rhodium, or iridium have been well studied for the hydrosilylation of alkynes (Scheme 1a), control of regio- and/or stereoselectivity is often problematic. In this context, unsym-

Scheme 1. Hydrosilylation of Unsymmetrical Internal Alkynes

a. Catalysis by transition metals or Lewis acids

$$R^1$$
 R^2 $Cat.$ R^1 R^2 R^3 R^4 R^5 R^5

- Limitation in functional group compatibility
- b. Present Study: Borane-catalyzed hydrosilylation of ynamides

$$R = [N] \xrightarrow{\text{Cat.} \atop B(C_6F_5)_3} \begin{bmatrix} \bigcirc \\ H^-B(C_6F_5)_3 \\ Si \end{bmatrix} [N]$$

$$Si = [N]$$

$$Si = [N]$$

$$\beta$$
-silicon effect

- · The first example of ynamide hydrosilylation
- Synthetically versatile β-silyl (Z)-enamide products
- · Transition metal-free
- · High regio- and stereoselectivity

metrical internal alkynes are especially challenging substrates. As a result, several approaches have been investigated to improve the limitations: 6 intramolecular hydrosilylation, 7 the use of polarized alkynes, 8 or the presence of internal directing groups. 9

Lewis acids have been shown to be effective for the hydrosilylation of alkynes, wherein Lewis acidic metal species are initially inserted into triple bonds and then transmetalation by silanes is followed to afford vinylsilane products. For instance, Molander and Yamamoto independently reported organo-aluminum or yttrium-catalyzed hydrosilylation of alkynes. Although these procedures were successfully applied to internal alkynes as well as terminal ones, there is room to improve the regioselectivity. In addition, compatibility of labile functional groups was rather insufficient mainly due to the acidic nature of the metal species (Scheme 1a). 11

We herein report the borane-catalyzed selective hydrosilylation of ynamides for the first time (Scheme 1b). The scope was found to be broad with high functional group tolerance under mild conditions. Moreover, the reaction occurs with excellent regio- and stereoselectivity to afford synthetically valuable β -silyl (Z)-enamide products. Based on the mechanistic studies, including computational calculations, the β -silicon effect was hypothesized to be responsible for the observed excellent stereoselectivity.

We commenced our study by monitoring a test reaction of a representative ynamide 1a with diethylsilane (2.0 equiv) in the presence of $B(C_6F_5)_3$ catalyst (5 mol %) in chloroform-d (Table 1). The progress of hydrosilylation was checked by 1H NMR spectroscopy. The hydrosilylation reaction proceeded with moderate efficiency at room temperature to afford β -silyl (Z)-enamide 2a (66% 1H NMR yield after 24 h, entry 1).

Received: November 21, 2016 Published: December 12, 2016



[†]Department of Chemistry, Korea Advanced Institute of Science and Technology (KAIST), Daejeon 34141, Korea

[‡]Center for Catalytic Hydrocarbon Functionalization, Institute for Basic Science (IBS), Daejeon 34141, Korea

Organic Letters Letter

Table 1. Optimization of Reaction Conditions

Ph
$$\frac{Ts}{N}$$
 Me + $\frac{Et_2SiH_2}{(2.0 \text{ equiv})}$ $\frac{cat. B(C_6F_5)_3}{solvent}$ $\frac{Ph}{HEt_2Si}$ $\frac{N}{Me}$ He $\frac{N}{N}$ $\frac{Ts}{Me}$ $\frac{2a}{(Z/E, >20:1)^b}$

entry	$B(C_6F_5)_3 \text{ (mol \%)}$	solvent	time (h)	$yield^{b}$ (%)
1 ^c	5	CDCl ₃	24	66
2	5	$CDCl_3$	6	94
3	3	CDCl ₃	6	96
4^d		$CDCl_3$	24	NR
5 ^e	3	CDCl ₃	24	71
6	3	toluene- d_8	24	79
7	3	THF- d_8	24	NR

^aConditions: **1a** (0.2 mmol), diethylsilane (2.0 equiv), and $B(C_6F_5)_3$ catalyst in CDCl₃ (0.8 mL). ^bDetermined by ¹H NMR analysis of the crude mixture (internal standard: mesitylene). ^cAt 25 °C. ^dAt 80 °C. ^eDiethylsilane (1.1 equiv).

Importantly, the isolated product was determined to display complete control of regio- and stereoselectivity as determined by NMR spectroscopy (see the Supporting Information for details). This result was significant in that while $B(C_6F_5)_3$ was known to catalyze hydrosilylation of unsaturated bonds such as olefins, carbonyls, imines, carboxylic acids, and nitriles, 12 only limited examples of internal alkynes were known to undergo the boranecatalyzed hydrosilylation. 13 Therefore, we determined to optimize the present hydrosilylation further by evaluating various reaction parameters.

The hydrosilylation efficiency was significantly improved by running the reaction at higher temperature even with lower catalyst loading and in shorter time, and excellent product yield was obtained in 6 h at 60 °C using 3 mol % of B(C_6F_5)₃ catalyst (entry 3). No reaction was observed in the absence of borane species (entry 4). The product yield was slightly decreased with the use of 1.1 equiv of silylating reagent (entry 5). Solvents other than chloroform were less effective in the current hydrosilylation of ynamide (entries 6 and 7).

With the above optimized reaction conditions in hand, we next explored the scope of N-sulfonyl ynamides (Scheme 2). Substrates bearing various para-substituents in the phenvlacetylenic moiety smoothly underwent the desired hydrosilylation leading to the corresponding β -silyl (Z)-enamides in high yields (2a-f), which displayed complete control of regioand stereoselectivity. Notably, the reaction efficiency was not affected by the electronic variation, and substrates having either electron-donating (Me or OMe) or -withdrawing substituents (Cl or CF₃) were hydrosilylated in good yields. The presence of ortho substituents in the phenyl moiety of ynamides did not give any detrimental effect on the desired reaction (2g and 2h). Ynamide derivatives having naphthyl or alkenyl side chains were also viable for the present hydrosilylation to furnish the corresponding products (2i and 2j, respectively). In the latter case, in particular, the olefinic double bond remained intact to prove that the present hydrosilylation is highly chemoselective.

When alkyl ynamides were subjected to the standard conditions, the desired hydrosilylation also took place smoothly with complete control of regio- and stereoselectivity. For instance, butyl-, cyclopropyl-, and cyclohexyl ynamides were all reacted to afford the corresponding β -silyl (Z)-enamides in good yields (2k-m). Replacement of the N-tosyl (Ts) group with the N-mesyl (Ms) did not hamper the reactivity and selectivity (2n).

Scheme 2. Substrate Scope of Ynamides

^aConditions: substrate (0.2 mmol), silane (2.0 equiv), and B(C_6F_5)₃ (3 mol %) in CHCl₃ (0.8 mL) at 60 °C. ^bDetermined by ¹H NMR analysis of the crude mixture (internal standard: mesitylene). ^cFor 12 h at 60 °C. ^dFor 0.5 h at 25 °C.

n-pentyl, 79% (2q) cyclohexyl, 52% (2r)^c

ORTEP of 2

In addition, variation of *N*-substituents in substrates from *N*-methyl to phenyl, cyclopropyl, *n*-pentyl, or cyclohexyl was compatible with the present conditions to afford the corresponding products in satisfactory yields (**2o**-**r**). The structure of **2r** was confirmed by X-ray crystallographic analysis. On the other hand, the reaction of acetylenic terminal ynamides and internal *N*-acyl ynamides was unsuccessful, and the desired vinylsilanes were not obtained.

The scope of silanes was next briefly examined (Scheme 3). Diphenylsilane, as another type of secondary hydrosilane in addition to dietheylsilane, worked well in the current reactions, still displaying excellent regio- and stereoselectivity (2s). Tertiary silylating reagents such as dimethylphenylsilane, triethylsilane, and benzyldimethylsilane were all efficient for the present hydrosilylation (2s-v). However, the reaction with triisopropylsilane or diethoxymethylsilane was ineffective even at elevated temperature.

The synthetic utility of the present $B(C_6F_5)_3$ -catalyzed hydrosilylation of ynamides was briefly examined. The reaction procedure was convenient to carry out on a gram scale with slightly lower loading of catalyst, still maintaining the high efficiency and excellent selectivity (eq 1). In addition, the silyl group in the obtained vinylsilane products was easily removed by silver fluoride, affording synthetically versatile (E)-enamide 3 without scrambling the double bond (eq 2). The structure of 3 was confirmed by X-ray crystallographic analysis (see the Supporting Information).

Organic Letters Letter

Scheme 3. Substrate Scope of Silanes^a

"Conditions: 1a (0.2 mmol), silane (2.0 equiv), and $B(C_6F_5)_3$ (3 mol %) in $CHCl_3$ (0.8 mL) at 60 °C. Determined by 1H NMR analysis of the crude mixture (internal standard: mesitylene). At 80 °C for 12 h.

We wondered about the rationale on the observed regio- and stereoselectivity in the present borane-catalyzed hydrosilyation of ynamides. In accordance with the precedents to form a borane-silane adduct I in situ, ¹⁴ the first step is believed to be a selective addition of ynamide to the silylium species, leading to ketene iminium species II (Scheme 4). ¹⁵ A subsequent hydride attack to

Scheme 4. Mode of the Present Hydrosilylation of Ynamides

the assumed intermediate II by borohydride III can proceed via two paths: (a) to the opposite side from the silyl group or (b) to the same side relative to the silyl moiety. While experimental results showed the exclusive formation of β -silyl (Z)-enamide products indicating the preference of path (a), computational calculations on this process were performed.

Calculations based on the density functional theory (DFT) showed that in a reaction of the presupposed ketene iminium intermediate (INT) with borohydride the solution-phase free

energy barriers differ 3.1 kcal/mol between anti-addition (10.8 kcal/mol) and syn-addition (13.9 kcal/mol, Figure 1). This difference in the free energy barriers ($\Delta\Delta G^{\ddagger}$) is in reasonably good agreement with the experimental results, suggesting that the anti-addition path will be predominant.




Figure 1. Relative energetics in the product-determining transition states [values in the parentheses are the relative free energies in kcal/mol; color code: H (white), B (pink), C (gray), N (purple), O (red), F (green), Si (gold), and S (yellow)].

Analysis of the geometry-optimized structure of ketene iminium species provided an additional insight into the reason for the observed stereoselectivity (Figure 2). The difference in




Figure 2. (a) Geometry-optimized structure of ketene iminium species and (b) schematic presentation of β-silicon effect.

dihedral angles between Si-C(1)-C(2) and C(3)-C(1)-C(2) was determined to be 18° . This geometrical bias can be attributed to the β -silicon effect to overlap the C(1)-Si and empty p-orbital of a carbocationic C(2) atom. ¹⁶ This alignment will accordingly decrease the dihedral angle of Si-C(1)-C(2) relative to that of C(3)-C(1)-C(2), ¹⁷ thus favoring the borohydride attack to ketene iminium species via the sterically more accessible *anti*-addition path.

In conclusion, the $B(C_6F_5)_3$ -catalyzed highly regio- and stereoselective hydrosilylation of ynamides has been developed to deliver synthetically versatile β -silyl (Z)-enamide products. The scope of ynamides and silanes was broad, and the reaction proceeds efficiently under mild and convenient conditions. The observed stereoselectivity was rationalized to result from a β -silicon effect of a postulated ketene iminium intermediate.

Organic Letters Letter

ASSOCIATED CONTENT

S Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.orglett.6b03485.

Complete experimental details and relevant spectra for all important compounds (PDF)

Crystallographic data for 2r (CIF)

Crystallographic data for 3 (CIF)

AUTHOR INFORMATION

Corresponding Author

*E-mail: sbchang@kaist.ac.kr.

ORCID ®

Sukbok Chang: 0000-0001-9069-0946

Author Contributions

§Y.K. and R.B.D. contributed equally.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This research was supported by the Institute for Basic Science (IBS-R010-D1). We thank Dr. Hyun-Ji Song (Center for Catalytic Hydrocarbon Functionalization, Institute for Basic Science) for helpful discussions.

REFERENCES

- (1) (a) Corey, J. Y. Chem. Rev. 2011, 111, 863. (b) Langkopf, E.; Schinzer, D. Chem. Rev. 1995, 95, 1375. (c) Marciniec, B. Catalysis of hydrosilylation of carbon-carbon multiple bonds: Recent progress; Springer: New York, 2002.
- (2) (a) Blumenkopf, T. A.; Overman, L. E. Chem. Rev. 1986, 86, 857. (b) Bunlaksananusorn, T.; Rodriguez, A. L.; Knochel, P. Chem. Commun. 2001, 745. (c) Denmark, S. E.; Liu, J. H. C. Angew. Chem., Int. Ed. 2010, 49, 2978. (d) Denmark, S. E.; Pan, W. Org. Lett. 2001, 3, 61. (e) Denmark, S. E.; Regens, C. S. Acc. Chem. Res. 2008, 41, 1486. (f) Fleming, I.; Henning, R.; Plaut, H. J. Chem. Soc., Chem. Commun. 1984, 29. (g) Nakao, Y.; Hiyama, T. Chem. Soc. Rev. 2011, 40, 4893. (h) Tamao, K.; Akita, M.; Maeda, K.; Kumada, M. J. Org. Chem. 1987, 52, 1100. (i) Tamao, K.; Ishida, N.; Tanaka, T.; Kumada, M. Organometallics 1983, 2, 1694. (j) McAdam, C. A.; McLaughlin, M. G.; Cook, M. J. Org. Chem. Front. 2015, 2, 510. (k) McIntosh, M. C.; Weinreb, S. M. J. Org. Chem. 1991, 56, 5010.
- (3) (a) Lim, D. S. W.; Anderson, E. A. Synthesis 2012, 44, 983. (b) Trost, B. Science 1991, 254, 1471.
- (4) (a) Chaulagain, M. R.; Mahandru, G. M.; Montgomery, J. Tetrahedron 2006, 62, 7560. (b) De Bo, G.; Berthon-Gelloz, G.; Tinant, B.; Markó, I. E. Organometallics 2006, 25, 1881. (c) Gallenkamp, D.; Fürstner, A. J. Am. Chem. Soc. 2011, 133, 9232. (d) Iglesias, M.; Sanz Miguel, P. J.; Polo, V.; Fernández-Alvarez, F. J.; Pérez-Torrente, J. J.; Oro, L. A. Chem. Eur. J. 2013, 19, 17559. (e) Na, Y.; Chang, S. Org. Lett. 2000, 2, 1887. (f) Trost, B. M.; Ball, Z. T.; Laemmerhold, K. M. J. Am. Chem. Soc. 2005, 127, 10028. (g) Zhou, H.; Moberg, C. Org. Lett. 2013, 15, 1444.
- (5) (a) Belger, C.; Plietker, B. Chem. Commun. 2012, 48, 5419. (b) Itami, K.; Mitsudo, K.; Nishino, A.; Yoshida, J.-i. J. Org. Chem. 2002, 67, 2645. (c) McBee, J. L.; Escalada, J.; Tilley, T. D. J. Am. Chem. Soc. 2009, 131, 12703. (d) Mo, Z.; Xiao, J.; Gao, Y.; Deng, L. J. Am. Chem. Soc. 2014, 136, 17414. (e) Sanada, T.; Kato, T.; Mitani, M.; Mori, A. Adv. Synth. Catal. 2006, 348, 51. (f) Wu, W.; Li, C.-J. Chem. Commun. 2003, 1668. (g) Zuo, Z.; Yang, J.; Huang, Z. Angew. Chem., Int. Ed. 2016, 55, 10839. (h) McLaughlin, M. G.; Cook, M. J. Chem. Commun. 2011, 47, 11104. (i) Tojo, S.; Isobe, M. Tetrahedron Lett. 2005, 46, 381. (j) Yong, L.; Kirleis, K.; Butenschön, H. Adv. Synth. Catal. 2006, 348, 833.

- (6) (a) Berthon-Gelloz, G.; Schumers, J.-M.; De Bo, G.; Markó, I. E. J. Org. Chem. 2008, 73, 4190. (b) Ding, S.; Song, L.-J.; Chung, L. W.; Zhang, X.; Sun, J.; Wu, Y.-D. J. Am. Chem. Soc. 2013, 135, 13835. (c) Rooke, D. A.; Ferreira, E. M. J. Am. Chem. Soc. 2010, 132, 11926.
- (7) (a) Denmark, S. E.; Pan, W. Org. Lett. **2002**, 4, 4163. (b) Marshall, J. A.; Yanik, M. M. Org. Lett. **2000**, 2, 2173. (c) Trost, B. M.; Ball, Z. T. J. Am. Chem. Soc. **2003**, 125, 30.
- (8) (a) Ding, S.; Song, L.-J.; Wang, Y.; Zhang, X.; Chung, L. W.; Wu, Y.-D.; Sun, J. Angew. Chem., Int. Ed. 2015, 54, 5632. (b) Hamze, A.; Provot, O.; Alami, M.; Brion, J.-D. Org. Lett. 2005, 7, 5625. (c) Song, L.-J.; Ding, S.; Wang, Y.; Zhang, X.; Wu, Y.-D.; Sun, J. J. Org. Chem. 2016, 81, 6157. (d) Stork, G.; Jung, M. E.; Colvin, E.; Noel, Y. J. Am. Chem. Soc. 1974, 96, 3684. (e) Sumida, Y.; Kato, T.; Yoshida, S.; Hosoya, T. Org. Lett. 2012, 14, 1552. (f) Sun, J.; Deng, L. ACS Catal. 2016, 6, 290. (g) Trost, B. M.; Ball, Z. T. J. Am. Chem. Soc. 2004, 126, 13942.
- (9) (a) Huang, K.-H.; Isobe, M. Eur. J. Org. Chem. 2014, 2014, 4733.
 (b) Kawasaki, Y.; Ishikawa, Y.; Igawa, K.; Tomooka, K. J. Am. Chem. Soc. 2011, 133, 20712.
- (10) (a) Holthausen, M. H.; Mehta, M.; Stephan, D. W. Angew. Chem., Int. Ed. 2014, 53, 6538. (b) Molander, G. A.; Retsch, W. H. Organometallics 1995, 14, 4570. (c) Pérez, M.; Hounjet, L. J.; Caputo, C. B.; Dobrovetsky, R.; Stephan, D. W. J. Am. Chem. Soc. 2013, 135, 18308. (d) Sudo, T.; Asao, N.; Gevorgyan, V.; Yamamoto, Y. J. Org. Chem. 1999, 64, 2494. (e) Sudo, T.; Asao, N.; Yamamoto, Y. J. Org. Chem. 2000, 65, 8919.
- (11) (a) Asao, N.; Sudo, T.; Yamamoto, Y. *J. Org. Chem.* **1996**, *61*, 7654. (b) Kato, N.; Tamura, Y.; Kashiwabara, T.; Sanji, T.; Tanaka, M. *Organometallics* **2010**, *29*, 5274.
- (12) (a) Blackwell, J. M.; Morrison, D. J.; Piers, W. E. Tetrahedron 2002, 58, 8247. (b) Blackwell, J. M.; Sonmor, E. R.; Scoccitti, T.; Piers, W. E. Org. Lett. 2000, 2, 3921. (c) Gandhamsetty, N.; Jeong, J.; Park, J.; Park, S.; Chang, S. J. Org. Chem. 2015, 80, 7281. (d) Gandhamsetty, N.; Joung, S.; Park, S.-W.; Park, S.; Chang, S. J. Am. Chem. Soc. 2014, 136, 16780. (e) Gandhamsetty, N.; Park, S.; Chang, S. J. Am. Chem. Soc. 2015, 137, 15176. (f) Gevorgyan, V.; Rubin, M.; Liu, J.-X.; Yamamoto, Y. J. Org. Chem. 2001, 66, 1672. (g) Kim, Y.; Chang, S. Angew. Chem., Int. Ed. 2016, 55, 218. (h) Oestreich, M.; Hermeke, J.; Mohr, J. Chem. Soc. Rev. 2015, 44, 2202. (i) Parks, D. J.; Piers, W. E. J. Am. Chem. Soc. 1996, 118, 9440. (j) Rubin, M.; Schwier, T.; Gevorgyan, V. J. Org. Chem. 2002, 67, 1936. (k) Gandhamsetty, N.; Park, J.; Jeong, J.; Park, S.-W.; Park, S.; Chang, S. Angew. Chem., Int. Ed. 2015, 54, 6832. (l) Simonneau, A.; Oestreich, M. Angew. Chem., Int. Ed. 2013, 52, 11905. (m) Süsse, L.; Hermeke, J.; Oestreich, M. J. Am. Chem. Soc. 2016, 138, 6940.
- (13) (a) Curless, L. D.; Ingleson, M. J. Organometallics 2014, 33, 7241. (b) Keess, S.; Simonneau, A.; Oestreich, M. Organometallics 2015, 34, 790.
- (14) (a) Hermeke, J.; Mewald, M.; Oestreich, M. J. Am. Chem. Soc. 2013, 135, 17537. (b) Houghton, A. Y.; Hurmalainen, J.; Mansikkamäki, A.; Piers, W. E.; Tuononen, H. M. Nat. Chem. 2014, 6, 983. (c) Nikonov, G. I.; Vyboishchikov, S. F.; Shirobokov, O. G. J. Am. Chem. Soc. 2012, 134, 5488. (d) Parks, D. J.; Blackwell, J. M.; Piers, W. E. J. Org. Chem. 2000, 65, 3090. (e) Rendler, S.; Oestreich, M. Angew. Chem., Int. Ed. 2008, 47, 5997.
- (15) (a) DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. Chem. Rev. 2010, 110, 5064. (b) Evano, G.; Coste, A.; Jouvin, K. Angew. Chem., Int. Ed. 2010, 49, 2840.
- (16) (a) Dallaire, C.; Brook, M. A. Organometallics 1990, 9, 2873. (b) Dallaire, C.; Brook, M. A. Organometallics 1993, 12, 2332. (c) Lambert, J. B.; Chelius, E. C. J. Am. Chem. Soc. 1990, 112, 8120. (d) Lambert, J. B.; Emblidge, R. W.; Malany, S. J. Am. Chem. Soc. 1993, 115, 1317. (e) Lambert, J. B.; Finzel, R. B. J. Am. Chem. Soc. 1982, 104, 2020. (f) Lambert, J. B.; Wang, G. T.; Finzel, R. B.; Teramura, D. H. J. Am. Chem. Soc. 1987, 109, 7838. (g) Pitt, C. G. J. Organomet. Chem. 1973, 61, 49. (h) Sommer, L. H.; Dorfman, E.; Goldberg, G. M.; Whitmore, F. C. J. Am. Chem. Soc. 1946, 68, 488.
- (17) (a) Wierschke, S. G.; Chandrasekhar, J.; Jorgensen, W. L. *J. Am. Chem. Soc.* **1985**, *107*, 1496. (b) Zhang, W.; Stone, J. A.; Brook, M. A.; McGibbon, G. A. *J. Am. Chem. Soc.* **1996**, *118*, 5764.