

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

### SYNTHESIS OF 1,2- AND 1,4-*bis*[ $\alpha$ -(2,4,6-TRINITROANILINO)-ACETAMIDO]BENZENE AS NEW INSENSITIVE EXPLOSIVES

B. P. Bandgar<sup>a</sup>; V. R. Damal<sup>b</sup>; S. S. Pandit<sup>b</sup>

<sup>a</sup> Organic Chem. Research Lab., School of Chemical Sciences, Swami Ramanand Teerth Marathwada University, Nanded, Maharashtra, INDIA <sup>b</sup> Padmashri Vikhe-Patil College, Ahmednagar, INDIA

**To cite this Article** Bandgar, B. P. , Damal, V. R. and Pandit, S. S.(1998) 'SYNTHESIS OF 1,2- AND 1,4-*bis*[ $\alpha$ -(2,4,6-TRINITROANILINO)-ACETAMIDO]BENZENE AS NEW INSENSITIVE EXPLOSIVES', *Organic Preparations and Procedures International*, 30: 6, 703 – 705

**To link to this Article:** DOI: 10.1080/00304949809355328

**URL:** <http://dx.doi.org/10.1080/00304949809355328>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

**SYNTHESIS OF 1,2- AND 1,4-bis[ $\alpha$ -(2,4,6-TRINITROANILINO)-  
ACETAMIDO]BENZENE AS NEW INSENSITIVE EXPLOSIVES<sup>†</sup>**

Submitted by  
(07/10/98)

B. P. Bandgar\*,<sup>††</sup> V. R. Damal<sup>†††</sup> and S. S. Pandit<sup>††</sup>


<sup>††</sup> *Organic Chem. Research Lab., School of Chemical Sciences  
Swami Ramanand Teerth Marathwada University  
Nanded- 431606, Maharashtra, INDIA*

<sup>†††</sup> *Padmashri Vikhe-Patil College  
Pravaranagar (Loni)-413713, Dist. Ahmednagar, INDIA*

There is a critical need in industry for explosive compositions which would be safe, reliable and stable at above 350° temperatures. For example, it is sometimes necessary to shoot explosive devices into hot wells at temperatures which may reach as high as 200-300°. In the steel industry, open hearth furnaces are tapped with explosive shaped charges in tap holes at temperatures above 500°. The explosives available at present, which have the best possible thermal properties, must have the shaped charges jacketed with heavy insulation. Even so, the charge must be initiated within 3-4 min. from the time it is set in place or it may fail due to thermal instability.<sup>1</sup> There is also a growing demand in connection with space programs for explosive compounds which are stable at elevated temperatures and low pressure. The temperature of conventional explosives subjected to aerodynamic heating on high speed aircrafts may rise above 80°, hence making them unsuitable for space aircraft applications.<sup>2</sup>

Explosives with improved high temperature properties, usually called "Heat Resistant Explosives" have emerged to meet such requirements. Nitro compounds such as TNT (2,4,6-trinitrotoluene) and picric acid have been found to be very useful heat resistant explosives.<sup>2-4</sup> They are of special interest because of their ability to withstand the high temperature and low pressure encountered in space aircraft applications.<sup>1-4</sup> No doubt, the manufacture of these explosives is so crucial that sometimes, even complex synthetic routes are adopted for the synthesis of heat resistant explosives. The importance of nitro compounds as heat resistant explosives led us to investigate the synthesis of new highly insensitive explosives such as 1,2-bis[ $\alpha$ -(2,4,6-trinitroanilino)acetamido]benzene and 1,4-bis[ $\alpha$ -(2,4,6-trinitroanilino)-acetamido]benzene (*Scheme*).

These two new explosives are condensation products of 1,2- and 1,4-phenylenediamines with 2,4,6 trinitroanilinoacetic acid chloride [TNAAC, **3**]. 2,4,6-Trinitroanilinoacetic acid [TNAAC, **2**] and its lead salts have been reported as energetic materials and additives in propellant formulations.<sup>5</sup> The theoretically calculated velocities of detonation [VOD] and impact sensitivities for **4** and **5** indicate these to be highly explosives but stable<sup>3,5</sup> up to 400°.



i)  $\text{HCl}\cdot\text{NH}_2\text{CH}_2\text{COOH}$ ,  $\text{Na}_2\text{CO}_3$  ii)  $\text{PCl}_3$  iii) *o*-phenylenediamine iv) *p*-phenylenediamine

### Scheme

### EXPERIMENTAL SECTION

All chemicals were of analytical grade. Solvents were distilled before use.  $^1\text{H}$  NMR spectra were recorded on a 90 MHz varian FT90A nmr instrument and IR spectra were recorded on a FT-IR Paragon-500 Perkin Elmer instrument.

**Synthesis of 2,4,6-Trinitroanilinoacetic Acid (TNAA, 2).**- A mixture of glycine (412 mg, 5 mmol),  $\text{Na}_2\text{CO}_3$  (530 mg, 5 mmol) in water (20 mL) and picryl chloride (1238 mg, 5 mmol) in acetone (20 mL) was stirred at  $25^\circ$  for 1h. The reaction mixture was cooled to  $10^\circ$  and acetone was removed under reduced pressure. The solid obtained was dissolved in a minimum amount of distilled water and neutralized with dil. HCl. The product was collected, dried under vacuum at  $40^\circ$  and recrystallized from hot water to give 801 mg (58%) of yellow solid, mp.  $158\text{-}160^\circ$  (dec.); IR: 3568, 3498, 3322, 3088, 2924, 2564, 1870, 1724, 1684, 1626, 1590,  $1534\text{ cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{D}_2\text{O}$ ):  $\delta$  3.9 (s, 2H,  $\text{CH}_2$ ), 8.8 (s, 2H, Ar-H).

*Anal.* Calcd. for  $\text{C}_{12}\text{H}_6\text{O}_8\text{N}_4$ : C, 33.57; H, 2.09; N, 19.38. Found: C, 33.64; H, 2.02; N, 19.44

**2,4,6-Trinitroanilinoacetyl Chloride (TNAAC, 3).**- A mixture of TNAA (1430 mg, 5 mmol) and  $\text{PCl}_5$  (1043 mg, 5 mmol) in chloroform (25 mL) was refluxed for 3 h. Then the reaction mixture was cooled to  $0^\circ$ . The product was collected, washed with cold hexane, dried and recrystallized from hot water to yield 809 mg (53%) of yellow solid, mp.  $101\text{-}104^\circ$ ; IR:  $1780\text{ cm}^{-1}$  (C=O),  $3300\text{ cm}^{-1}$  (NH);  $1560\text{ cm}^{-1}$  ( $\text{NO}_2$ );  $^1\text{H}$  NMR:  $\delta$  4.0 (d, 2H,  $\text{CH}_2$ ), 9.0 (s, 2H, Ar-H), 9.5 (m, 1H, NH).

*Anal.* Calcd. for  $\text{C}_{12}\text{H}_5\text{ClO}_7\text{N}_4$ : C, 31.52; H, 1.64; N, 18.39. Found: C, 31.43; H, 1.58; N, 18.48

**Synthesis of 1,2- and 1,4-bis-[ $\alpha$ -(2,4,6-Trinitroanilino)acetamido]benzene (4 and 5).**- A mixture of *o*-phenylenediamine or *p*-phenylenediamine (270 mg, 2.5 mmol) and TNAAC (1525 mg, 5 mmol)

in DMF (50 mL) was stirred at 25° for 3 h. The orange colored product formed was collected, washed and dried. Yield of **4** = 805 mg (25%), mp. 410° (dec.); IR: 1652, 1620 cm<sup>-1</sup> (C=O), 3448, 3316 cm<sup>-1</sup> (NH), 1574 cm<sup>-1</sup> (NO<sub>2</sub>); <sup>1</sup>H NMR: δ 4.3 (d, 4H, 2xCH<sub>2</sub>), 8.2 (s, 4H, Ar-H), 9.2 (s, 4H, Ar-H), 9.5 (m, 2H, NH), 9.7 (m, 2H, NH).

*Anal.* Calcd. for C<sub>22</sub>H<sub>16</sub>N<sub>10</sub>O<sub>8</sub>: C, 48.18; H, 2.92; N, 25.55. Found: C, 48.10; H 2.81; N, 25.66

Yield of **5**: (1095 mg, 34%), mp. 430° (dec.); IR: 1650, 1630 cm<sup>-1</sup> (C=O), 3400, 3310 cm<sup>-1</sup> (NH), 1570 cm<sup>-1</sup> (NO<sub>2</sub>); <sup>1</sup>H NMR: δ 4.4 (d, 4H, 2xCH<sub>2</sub>), 8.5 (s, 4H, Ar-H), 9.3 (s, 4H, Ar-H), 9.4 (m, 2H, NH), 9.6 (m, 2H, NH).

*Anal.* Calcd for C<sub>22</sub>H<sub>16</sub>N<sub>10</sub>O<sub>8</sub>: C, 48.18; H, 2.92; N, 25.55. Found: C, 48.22, H, 2.87, N, 25.47

**Acknowledgement.**- VRD thanks Principal Dr. P. S. Gunjal and Dr. J. N. Nigal for their encouragement.

### REFERENCES

† This paper is dedicated to Prof. M. S. Wadia (University of Pune) on the occasion of his 60th birthday. For accompanying paper, see B. P. Bandgar, S. A. Kulkarni and J. N. Nigal, *Org. Prep. Proced. Int.*, **30**, 706 (1998).

1. R. J. Spaer and W. S. Wilson, *J. Energy Materials*, **2**, 61 (1984).
2. G. A. Olah and D. R. Squire, *Chemistry of Energetic Materials*, Academic Press, New York (1991).
3. R. Meyer and J. Kohler, *Explosives* 4th Ed. VCH, Weinheim (1993).
4. K. U. B. Rao, R. R. Soman and H. Singh, *J. Energy Materials*, **8**, 99 (1990).
5. R. R. Soman, M.Sc. Thesis, University of Pune (1988).

\*\*\*\*\*