

This article was downloaded by:

On: 26 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

### NEW AND IMPROVED SYNTHESIS OF ( $\pm$ )-4-BROMO-7,8-DIMETHOXY-ISOCHROMAN-3-ONE, A KEY INTERMEDIATE FOR THE ELABORATION OF ( $\pm$ )-*cis*-ALPININE AND ( $\pm$ )-*cis*-ALPINIGENINE

Darío A. Bianchi<sup>a</sup>; Teodoro S. Kaufman<sup>a</sup>

<sup>a</sup> Institute de Química Orgánica de Síntesis -IQUIOS- (CONICET-UNR) and Facultad de Ciencias Bioquímicas y Farmacéuticas Universidad Nacional de Rosario, Rosario, REPÚBLICA ARGENTINA

**To cite this Article** Bianchi, Darío A. and Kaufman, Teodoro S.(2004) 'NEW AND IMPROVED SYNTHESIS OF ( $\pm$ )-4-BROMO-7,8-DIMETHOXY-ISOCHROMAN-3-ONE, A KEY INTERMEDIATE FOR THE ELABORATION OF ( $\pm$ )-*cis*-ALPININE AND ( $\pm$ )-*cis*-ALPINIGENINE', Organic Preparations and Procedures International, 36: 2, 195 – 199

**To link to this Article:** DOI: 10.1080/00304940409355398

**URL:** <http://dx.doi.org/10.1080/00304940409355398>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


**NEW AND IMPROVED SYNTHESIS OF ( $\pm$ )-4-BROMO-7,8-DIMETHOXY-  
ISOCHROMAN-3-ONE, A KEY INTERMEDIATE FOR THE ELABORATION  
OF ( $\pm$ )-*cis*-ALPININE AND ( $\pm$ )-*cis*-ALPINIGENINE**

Submitted by  
(11/10/03)


Darío A. Bianchi and Teodoro S. Kaufman\*

*Instituto de Química Orgánica de Síntesis –IQUIOS– (CONICET-UNR)  
and Facultad de Ciencias Bioquímicas y Farmacéuticas  
Universidad Nacional de Rosario, Suipacha 531  
S2002LRK Rosario, REPÚBLICA ARGENTINA  
E-mail: tkaufman@fbioyf.unr.edu.ar*

The Rhoeadine-Papaverrubine alkaloids constitutes a small family of natural products, elaborated almost exclusively by the *Papaver* genus; many of the members of this family have potentially useful biological activity.<sup>1</sup> In 1982, Ahmad and Snieckus<sup>2</sup> reported a convergent entry into the rhoeadan skeleton and a formal total synthesis of ( $\pm$ )-*cis*-alpinigenine (1) and the related ( $\pm$ )-*cis*-alpinine (2), using bromolactone 3 as key intermediate. The bromolactone was prepared by these authors employing different approaches; however, regioselectivity problems arising



during isochromanone ring formation and in the final bromination step under free radical conditions, lead to mixtures of regioisomeric isochroman-3-one derivatives, as well as nuclear and benzylic brominated products; moreover, in spite that improved syntheses of precursors of 3 have been reported,<sup>3,4</sup> the efficiency of the overall sequence demonstrated to be rather poor. Here, we report a convenient, alternative and efficient synthesis of bromolactone 3, starting from 2,3-dimethoxytoluene (4), which proceeds through the known glyoxylate 5.<sup>5</sup>



Reagents and conditions: a. See ref. 5 (75%); b) NaCNBH<sub>3</sub>, AcOHgl, EtOH, c) Camphorsulfonic acid (5 mol%), MeOH, RT, 6 h (89%); d) 1. SMe<sub>2</sub>, NBS, CH<sub>2</sub>Cl<sub>2</sub>, RT→0°C; 2.  $\alpha$ -hydroxylactone 7, 0°C→RT, overnight (89%)

The sequence is based on the acid-catalyzed lactonization of **6**, which avoids production of regioisomeric isochroman-3-ones and a bromodehydroxylation reaction of the resulting hydroxylactone **7**, which does not generate nuclearily halogenated products. Selective reduction of **5** with sodium cyanoborohydride in ethanol, to which glacial acetic acid was added in order to accelerate the reaction, smoothly and cleanly afforded 91% of mandelate **6**. Next, lactonization was carried out by exposure of **6** to camphorsulfonic acid in dry methanol at room temperature,<sup>6</sup> furnishing 89% of  $\alpha$ -hydroxyketone **7** as a solid. Finally, bromination of **7** was undertaken; however, the  $\text{PPh}_3\cdot\text{Br}_2$  adduct in DMF<sup>7</sup> gave disappointingly low yields (Table), while the  $\text{PPh}_3\text{-NBS}$  reagent<sup>8</sup> furnished only 41% of **3** (Entries 1 and 2) and the conditions shown in Entry 3 provided 32% of the related chloride.

**Table.** Elaboration of Bromolactone **3** by Bromodehydroxylation of **7**

Entry N <sup>o</sup>	Reagent	Conditions	Yield (%)
1	$\text{PPh}_3\cdot\text{Br}_2$	DMF, RT	< 10
2	$\text{PPh}_3$ , NBS	$\text{CH}_2\text{Cl}_2$ , 0°C	41
3	$\text{MeSO}_2\text{Cl}$ , $\text{EtN}_3$ , LiBr, $\text{Bu}_4\text{N}^+\text{Br}$	$\text{CH}_2\text{Cl}_2$ , 0°C→RT, overnight	-
4	$\text{SMe}_2$ , NBS	$\text{CHCl}_3$ , 0°C→60°C, 30 min	< 30
5	$\text{SMe}_2$ , NBS	$\text{CHCl}_3$ , 0°C	38
6	$\text{SMe}_2$ , NBS	$\text{CH}_2\text{Cl}_2$ , -10°C→reflux	56
7	$\text{SMe}_2$ , NBS	$\text{CH}_2\text{Cl}_2$ , -20°C→RT	59
8	$\text{SMe}_2$ , NBS (RT)	$\text{CH}_2\text{Cl}_2$ , 0°C	70
9	$\text{SMe}_2$ , NBS (RT)	$\text{CH}_2\text{Cl}_2$ , 0°C→RT, overnight	89

Yields improved when the  $\text{SMe}_2\text{-NBS}^9$  couple was employed. Preparation of the reagent at 0°C in  $\text{CHCl}_3$  followed by reflux in the presence of **7**, produced **3** (< 30%) and inseparable side products (Entry 4). However, when the reaction was heated at lower temperature, 56% of the product was recovered; yields did not improve significantly when the reagents were mixed at -20°C and allowed to attain room temperature (Entries 5 and 7). However, preparation of the  $\text{SMe}_2\text{-NBS}$  reagent at room temperature (Entries 8 and 9) produced considerable enhancements in generation of brominated product, furnishing the best yields (89%) when the reaction was left overnight at room temperature. Under these conditions, the overall yield of **3** from **4** was 54%, more than twice the reported yields for the elaboration of this key intermediate. Compound **3** is highly sensitive to moisture; its spectral data fully agreed with those reported in the literature.

In conclusion, we have developed a simple, practical and efficient synthesis of bromolactone **3**, which proceeds in six high-yield steps from commercially available 2,3-dimethoxytoluene; this procedure is devoid of the drawbacks affecting previous syntheses.

## EXPERIMENTAL SECTION

The melting points were taken on an Ernst Leitz Wetzlar model 350 hot-stage microscope and are uncorrected. FT-IR spectra were determined with a Bruker IFS 25 infrared spectrophotometer. The  $^1\text{H}$  and  $^{13}\text{C}$ -NMR spectra were acquired with a Bruker AC200-E spectrometer (200.13 MHz for  $^1\text{H}$ ), employing  $\text{CDCl}_3$  as solvent; the chemical shifts are expressed in  $\delta$  ppm downfield from the internal standard (TMS). High resolution mass spectral data were obtained from the Kent Mass Spectrometry Unit 1 (Kent, UK). Reagents and solvents were used as received; dry MeOH was obtained by distillation from magnesium methoxide. Dry  $\text{CHCl}_3$  and  $\text{CH}_2\text{Cl}_2$  were accessed by reflux over  $\text{P}_2\text{O}_5$ , followed by distillation. Flash column chromatographies were carried out with silica gel 60 H and eluted with hexane-EtOAc employing gradient techniques. All new compounds gave single spots on TLC plates (Merck, art. 5554) run in different solvent systems. Chromatographic spots were detected by exposure to UV light (254 nm) followed by spraying with ethanolic *p*-anisaldehyde/sulfuric acid reagent and careful heating of the plates for better selectivity.

**(±)-Ethyl {2-[(Acetoxy)methyl]-3,4-dimethoxyphenyl}hydroxyacetate (6).**- Under a dry nitrogen atmosphere, a solution of keto ester **5<sup>5</sup>** (1001.4 mg, 3.23 mmol) in absolute ethanol (40 mL) was successively treated with glacial acetic acid (0.184 mL, 3.23 mmol) and sodium cyanoborohydride (223.7 mg, 3.55 mmol) and the mixture was stirred overnight at room temperature. The reaction was quenched with 1N NaOH (5 mL), diluted with brine (10 mL) and extracted with EtOAc (3 x 30 mL). The combined organic extracts were washed once with brine (5 mL), dried ( $\text{Na}_2\text{SO}_4$ ), concentrated under reduced pressure and chromatographed to provide mandelate **6** (918 mg, 91%), as a colorless oil; IR (film): 3460, 2980, 2850, 1740, 1730, 1600, 1500, 1460, 1280, 1090 and 810  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR:  $\delta$  1.22 (s, 3H,  $J = 8.6$ ,  $\text{OCH}_2\text{Me}$ ), 1.60 (bs, 1H, OH), 2.07 (s, 3H,  $\text{MeCO}_2$ ), 3.84 (s, 3H, OMe), 3.86 (s, 3H, OMe), 4.20 (dq, 2H,  $J = 1.0, 8.6$ ,  $\text{OCH}_2\text{Me}$ ), 5.35 (s, 2H,  $\text{ArCH}_2\text{O}$ ), 5.40 (s, 1H,  $\text{ArCHOH}$ ), 6.92 (d, 1H,  $J = 8.6$ , ArH) and 7.10 (d, 1H,  $J = 8.6$ , ArH);  $^{13}\text{C}$  NMR:  $\delta$  13.88 ( $\text{MeCH}_2$ ), 20.86 ( $\text{MeCO}_2$ ), 55.56 (OMe-7), 57.60 ( $\text{ArCH}_2\text{O}$ ), 61.20 (OMe-8), 61.97 ( $\text{MeCH}_2\text{O}$ ), 69.77 ( $\text{ArCHOH}$ ), 112.71 (C-5), 122.97 (C-6), 128.03 (C-2), 130.88 (C-1), 148.46 (C-4), 152.65 (C-3), 170.73 (C=O) and 173.60 (C=O). HRMS for  $\text{C}_{15}\text{H}_{20}\text{O}_7$ . Calcd.: 312.12087. Found: 312.12087.

*Anal.* Calcd for  $\text{C}_{15}\text{H}_{20}\text{O}_7$ : C, 57.69; H, 6.45. Found: C, 57.60; H, 6.52

**(±)-4-Hydroxy-7,8-dimethoxy-isochroman-3-one (7).**- A mixture of mandelate **6** (250 mg, 0.80 mmol) and camphorsulfonic acid (93 mg, 0.4 mmol) in anhydrous MeOH (8 mL) was warmed to 40°C and stirred for 6 h. After cooling, the reaction mixture was diluted with brine (10 mL), and extracted with EtOAc (3 x 10 mL); the organic extracts were washed with brine (10 mL), dried over  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo*. Chromatography of the residue give hydroxylactone **7** (160 mg, 89%), as a white solid mp. 126-128°C (hexane-EtOAc). IR (KBr): 3466, 2980, 2870, 1734, 1496, 1456, 1380, 1276, 1258, 1206, 1142, 1086, 1024, 818 and 806

$\text{cm}^{-1}$ ;  $^1\text{H}$  NMR:  $\delta$  3.62 (bs, 1H,  $w_{1/2} = 8$  Hz, OH), 3.87 (s, 3H, OMe), 3.88 (s, 3H, OMe), 5.10 (bs, 1H, H-4), 5.16 (d, 1H,  $J = 14.5$ ,  $\text{ArCH}_2\text{O}$ ), 5.64 (d, 1H,  $J = 14.5$ ,  $\text{ArCH}_2\text{O}$ ), 6.98 (d, 1H,  $J = 8.4$ , H-6) and 7.30 (d, 1H,  $J = 8.4$ , H-5);  $^{13}\text{C}$  NMR:  $\delta$  55.86 (OMe-7), 61.04 (OMe-8), 64.06 (C-1), 67.24 (C-4), 113.01 (C-6), 118.92 (C-7a), 122.90 (C-5), 126.59 (C-4a), 144.40 (C-7), 151.66 (C-8) and 173.76 (C-3). HRMS for  $\text{C}_{11}\text{H}_{12}\text{O}_5$ . Calcd.: 224.06847. Found: 224.06823.

*Anal.* Calcd. for  $\text{C}_{11}\text{H}_{12}\text{O}_5$ : C, 58.93; H, 5.39; Found: C, 58.98; H, 5.35

**(±)-4-Bromo-7,8-dimethoxy-isochroman-3-one (3).**- Under a dry nitrogen atmosphere, a stirred solution of NBS (159 mg, 0.89 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (2 mL) was treated dropwise with methyl sulfide (0.082 mL, 1.37 mmol) at room temperature. The mixture was cooled to  $0^\circ\text{C}$ , a solution of hydroxylactone **7** (100 mg, 0.35 mmol) in  $\text{CH}_2\text{Cl}_2$  (2 mL) was introduced dropwise and the system was left to attain room temperature. When the reaction was completed (TLC), brine (5 mL) was added and the products were extracted with EtOAc (3 x 15 mL); the combined organic extracts were washed once with brine (5 mL), dried ( $\text{Na}_2\text{SO}_4$ ), concentrated and chromatographed, furnishing **3** (114 mg, 89%) as an oil which crystallized upon standing in the refrigerator, furnishing a solid mp.  $121\text{-}123^\circ\text{C}$ , *lit.*<sup>2</sup>  $122\text{-}123^\circ\text{C}$ ; IR (film): 3000, 2950, 2840, 1750, 1600, 1490, 1380, 1280, 1160, 1090, 1049, 990, 970, 895, 820, 765 and  $690\text{ cm}^{-1}$ ;  $^1\text{H}$  NMR:  $\delta$  3.87 (s, 3H, OMe), 3.89 (s, 3H, OMe), 5.40 (s, 1H, H-4), 5.44 (d, 1H,  $J = 15.3$ ,  $\text{ArCH}_2\text{O}$ ), 5.61 (d, 1H,  $J = 15.3$ ,  $\text{ArCH}_2\text{O}$ ), 6.92 (d, 1H,  $J = 8.4$ , H-6) and 7.12 (d, 1H,  $J = 8.4$ , H-5);  $^{13}\text{C}$  NMR:  $\delta$  39.46 (C-4), 55.83 (C-7), 60.96 (C-7), 65.16 (C-1), 112.90 (C-6), 123.83 (C-8a), 123.91 (C-5), 126.76 (C-4a), 144.95 (C-7), 153.45 (C-8) and 165.66 (C-3); EI-MS,  $m/z$  (%): 288, 286 ( $\text{M}^+$ , 6), 208 (26), 207 (100), 179 (24) and 151 (41).

**Acknowledgments.**- The authors acknowledge CONICET, ANPCyT, SECyT-UNR and Fundación Antorchas for financial support. D. A. B. thanks CONICET for a fellowship. Helpful assistance provided by Lic. Federico Rúa is also gratefully acknowledged.

## REFERENCES

1. M. W. Scheve, *Acta Biol. Med. Ger.*, **35**, 1019 (1976); (b) D. Walterová, K. Hátle, A. Nemecková and B. Vecerek, *Cesk Farm.*, **27**, 314 (1978); *Chem. Abstr.*, **90**, 145784 (1979).
2. I. Ahmad and V. Snieckus, *Can. J. Chem.*, **60**, 2678 (1982).
3. W. Nagata, K. Okada and T. Aoki, *Synthesis*, 365 (1979), and references cited therein.
4. N. S. Narasimhan, R. S. Mali and B. S. Kulkarni, *Tetrahedron Lett.*, 2797 (1981).
5. T. S. Kaufman, *Heterocycles*, **55**, 323 (2001).

6. (a) S. K. Taylor, J. A. Fried, Y. N. Grassi, A. E. Marolewski, E. A. Pelton, T.-J. Poel, D. S. Rezanka and M. R. Whittaker, *J. Org. Chem.*, **58**, 7304 (1993); (b) T. Konoike and Y. Araki, *J. Org. Chem.*, **59**, 7849 (1994).
7. J. B. P. A. Wijnberg, P. G. Wiering and H. Steinberg, *Synthesis*, 901 (1981).
8. B. Imperiali and R. S. Roy, *J. Org. Chem.*, **60**, 1891 (1995).
9. E. J. Corey, C. U. Kim and M. Takeda, *Tetrahedron Lett.*, 4339 (1972).