

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

### DIRECT $\alpha$ -HYDROXYLATION OF ARYL KETONES USING A HYPERVALENT IODINE(III) SULFONATE

H. -Y. Huang<sup>a</sup>; R. -S. Hou<sup>b</sup>; H. -M. Wang<sup>b</sup>; L. -C. Chen<sup>a</sup>

<sup>a</sup> Graduate Institute of Pharmaceutical Sciences, College of Pharmacy, Kaohsiung Medical University, Kaohsiung, Taiwan Republic of CHINA <sup>b</sup> Chung Hwa College of Medical Technology, Tainan, Taiwan Republic of CHINA

**To cite this Article** Huang, H. -Y. , Hou, R. -S. , Wang, H. -M. and Chen, L. -C.(2006) 'DIRECT  $\alpha$ -HYDROXYLATION OF ARYL KETONES USING A HYPERVALENT IODINE(III) SULFONATE', *Organic Preparations and Procedures International*, 38: 5, 473 – 476

**To link to this Article:** DOI: 10.1080/00304940609356439

**URL:** <http://dx.doi.org/10.1080/00304940609356439>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

8. J. S. Yadav, B. V. S. Reddy, A. D. Krishna and T. Swamy, *Tetrahedron Lett.*, **44**, 6055 (2003).
9. The bioassay of herbicidal activity was evaluated by the Research Institute of Elemento-Organic Chemistry of Nankai University.
10. W. O. Kermack, W. H. Perkin, Jr., and R. Robinson, *J. Chem. Soc.*, **119**, 1602 (1921); J. G. Cannon, B. J. Demopoulos, J. P. Long, J. R. Flynn and F. M. Sharabi, *J. Med. Chem.*, **24**, 238 (1981).

\*\*\*\*\*

**DIRECT  $\alpha$ -HYDROXYLATION OF ARYL KETONES  
USING A HYPERVALENT IODINE(III) SULFONATE**


*Submitted by* H.-Y. Huang<sup>†</sup>, R.-S. Hou<sup>††</sup>, H.-M. Wang<sup>††</sup>, and L.-C. Chen<sup>\*†</sup>  
(01/25/06)

<sup>†</sup> *Graduate Institute of Pharmaceutical Sciences, College of Pharmacy  
Kaohsiung Medical University, Kaohsiung  
Taiwan 807, Republic of CHINA*

<sup>††</sup> *Chung Hwa College of Medical Technology  
Tainan, Taiwan 717, Republic of CHINA  
e-mail: m715003@cc.kmu.edu.tw*

$\alpha$ -Hydroxyketones are widely used intermediates and are important constituents of many biologically important natural products.<sup>1</sup> The methods used for their synthesis involve the direct  $\alpha$ -hydroxylation of ketones under basic<sup>2</sup> or acidic conditions.<sup>3</sup> Only a few reports in the literature account for the direct  $\alpha$ -hydroxylation of ketones under neutral conditions.

Recently, hypervalent iodine(III) reagents have been used extensively in organic synthesis due to their low toxicity, ready availability and easy handling.<sup>4</sup> As a part of our on-going studies to utilize hypervalent iodine (III) reagents in organic synthesis, we report here a new and direct method for the synthesis of  $\alpha$ -hydroxy aryl ketones by the reaction of aryl ketones with [hydroxy-(2,4-dinitrobenzenesulfonyloxy)iodo] benzene (HDNIB) under mild conditions. The required HDNIB was prepared in satisfactory yields from the reaction of 2,4-dinitrobenzenesulfonic acid with phenyliodine (III) diacetate (PIDA).<sup>5</sup> Treatment of aryl ketones (**1**) with HDNIB in CH<sub>3</sub>CN at reflux for 1 h produced the  $\alpha$ -(2,4-dinitrobenzenesulfonyloxy)-ketone intermediates (**2**),<sup>5</sup> which can then undergo hydrolysis with DMSO-H<sub>2</sub>O system at room temperature for 2 h to give  $\alpha$ -hydroxyaryl ketones (**3**) in good yields as shown in the *Scheme*.



- a)  $\text{R}_1 = \text{C}_6\text{H}_5$ ,  $\text{R}_2 = \text{H}$ ; b)  $\text{R}_1 = 4\text{-MeC}_6\text{H}_4$ ,  $\text{R}_2 = \text{H}$ ; c)  $\text{R}_1 = 4\text{-MeOC}_6\text{H}_4$ ,  $\text{R}_2 = \text{H}$ ; d)  $\text{R}_1 = 4\text{-ClC}_6\text{H}_4$ ,  $\text{R}_2 = \text{H}$ ; e)  $\text{R}_1 = 4\text{-BrC}_6\text{H}_4$ ,  $\text{R}_2 = \text{H}$ ; f)  $\text{R}_1 = 3\text{-Furyl}$ ,  $\text{R}_2 = \text{H}$ ; g)  $\text{R}_1 = 3\text{-Thienyl}$ ,  $\text{R}_2 = \text{H}$ ; h)  $\text{R}_1 = \text{C}_6\text{H}_5$ ,  $\text{R}_2 = \text{Me}$ ; i)  $\text{R}_1 = \text{C}_6\text{H}_5$ ,  $\text{R}_2 = \text{C}_6\text{H}_5$

Our experiments involving a one-pot procedure for the preparation of  $\alpha$ -hydroxy-aryl ketones (**3**) by treatment of HDNIB with aryl ketones and subsequent hydrolysis using DMSO- $\text{H}_2\text{O}$  at room temperature was successful. The results are summarized in the Table. When the reaction was conducted by replacing HDNIB with HTIB (Koser's reagent)<sup>6</sup> under the same conditions, the preparation of 2-hydroxy-1-phenylethanone (**3a**) requires refluxing for 7 h. This observation clearly demonstrated that the leaving ability of  $-\text{ODNs}$  is superior to  $-\text{OTs}$  in nucleophilic substitution reactions.

**Table.** Preparation of  $\alpha$ -Hydroxyketones **3a-i**

Cmpd <sup>a</sup>	Yield (%)	mp. (°C)	lit. mp. (°C)
<b>3a</b>	88	87-88	80-82 <sup>7</sup>
<b>3b</b>	85	79-80	81-83 <sup>7</sup>
<b>3c</b>	83	104-105	99-101 <sup>7</sup>
<b>3d</b>	86	120-121	118-120 <sup>7</sup>
<b>3e</b>	84	133-134	134-136 <sup>7</sup>
<b>3f</b>	78	80-81	79-81 <sup>7</sup>
<b>3g</b>	76	71-72	73-74 <sup>8</sup>
<b>3h</b>	82	Oil	Oil <sup>7</sup>
<b>3i</b>	72	129-131	126-128 <sup>7</sup>

a) All products are known compounds and their physical constants, IR and <sup>1</sup>H NMR spectra correspond to those reported in literature.

In summary, the method described herein provides a good approach for the synthesis of  $\alpha$ -hydroxyarylktones by the reaction of aryl ketones with hypervalent iodine(III) sulfonate (HDNIB) and subsequent hydrolysis using DMSO- $\text{H}_2\text{O}$  at room temperature in a one-pot procedure.

## EXPERIMENTAL SECTION

All melting points are uncorrected. The IR spectra were recorded on a Shimadzu IR-27 G spectrophotometer.  $^1\text{H}$  NMR spectra were recorded on a Varian Unity Plus 400 MHz. Chemical shifts ( $\delta$ ) were measured in ppm with respect to TMS. MS were obtained on a JEOL JMS D-300 instrument.

**Typical Procedure.**- A mixture of acetophenone (120 mg, 1.0 mmol) and HDNIB (468 mg, 1.0 mmol) in acetonitrile (20 mL) was heated at reflux for 1 h. After the reaction mixture had been cooled to room temperature, 12 mL of DMSO- $\text{H}_2\text{O}$  (1 : 2) was added and the mixture was stirred at room temperature for 2 h. After removal of acetonitrile under reduced pressure, the aqueous phase was extracted with  $\text{CH}_2\text{Cl}_2$ . The combined organic extracts were concentrated under reduced pressure. The resulting residue was then dissolved in  $\text{Et}_2\text{O}$ , washed with  $\text{H}_2\text{O}$ , and dried over  $\text{MgSO}_4$ . The solvent was evaporated off and the residue was purified by chromatography on silica gel eluting with AcOEt-*n*-hexane (1:3) to give 120 mg (88%) of **3a**. IR (KBr): 3425, 1684  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  3.25-3.75 (br s, 1H), 4.88 (s, 2H), 7.49-7.94 (m, 5H); MS (EI)  $m/z$  : 136 ( $\text{M}^+$ ), 105, 83, 57, 45.

**Acknowledgement.**- We gratefully acknowledge the National Science Council of the Republic of China for the financial support of this work (Grant No. 94-2113-M-037-008).

## REFERENCES

1. B. Raduchel, *Synthesis*, 292 (1980).
2. A. B. Jones, "Comprehensive Organic Synthesis", Vol. 7, CH. 2 and 3, B. M. Trost, ed., Pergamon Press. Oxford, 1991.
3. A. Mckillop, J. D. Hunt, and E. C. Taylor, *J. Org. Chem.*, **37**, 3381 (1972); R. M. Moriarty, B. A. Berglund, and R. Penmasta, *Tetrahedron Lett.*, **33**, 6065 (2002).
4. Reviews, see: D. F. Banks, *Chem. Rev.*, **66**, 243 (1966); A. Varvoglis, *Chem. Soc. Rev.*, **10**, 377 (1981); G. F. Koser, in "The Chemistry of Functional Groups", Supplement D, ed. by S. Patai and Z. Rappoport, Wiley, New York, NY, 1983, CH. 18 and 25; A. Varvoglis, *Synthesis*, 709 (1984); R. M. Moriarty and O. Parkash, *Acc. Chem. Res.*, **19**, 244 (1986); M. Ochiai and Y. Nagao, *Yuki Gosei Kagaku Kyokaishi*, **44**, 660 (1986); R. M. Moriarty, R. K. Vaid, and G. F. Koser, *Synlett*, 365 (1990); A. Varvoglis, "The Organic Chemistry of Polycordinated Iodine", VCH, New York, NY, 1992; Y. Kita, H. Tohma, and T. Yakura, *Trends Org. Chem.*, **3**, 113 (1992); Y. Kita and H. Tohma, *Farumashia*, **28**, 984 (1992); P. J. Stang, *Angew. Chem. Int. Ed. Engl.*, **31**, 274 (1992); T. Kitamura, *Yuki Gosei Kagaku Kyokaishi*, **53**, 893 (1995); P. J. Stang and V. V. Zhdankin, *Chem. Rev.*, **96**, 1123 (1996); T. Kitamura and Y. Fujiwara, *Org. Prep. Proc. Int.*, **29**, 411 (1997).
5. J. C. Lee, J.-H. Choi, and Y. C. Lee, *Synlett*, 1563 (2001).

6. R. M. Moriarty, R. K. Vaid, and G. F. Koser, *Synlett*, 365 (1990); G. F. Koser, *Aldrichim. Acta*, **34**, 89 (2001).
7. Y.-Y. Yuan and Z.-C. Chen, *Synth. Commun.*, **32**, 1875 (2002).
8. R. M. Moriarty, M. P. Duncan, and O. Prakash, *J. Chem. Soc. Perkin Trans. I.*, 1781 (1987).

\*\*\*\*\*


### IMPROVED METHOD FOR THE CYCLIZATION OF *ortho*-ALKYNYLBENZENEDIAZONIUM SALTS

Submitted by  
(02/07/06)

Nataliya A. Zolnikova, Lidiya G. Fedenok\* and Nikolay E. Polyakov

*Institute of Chemical Kinetics and Combustion,  
Siberian Branch of Russian Academy of Sciences  
Institutskaya 3, 630090 Novosibirsk, RUSSIA  
e-mail: fedenok@kinetics.nsc.ru*

In the last decade, the Richter reaction has elicited considerable interest among chemists for two reasons: 1) the requirement to produce condensed heteroaromatic compounds with high biological activity, and 2) creation of convenient methods for introduction of alkyne groups into the aromatic ring, thereby making *vic*-alkynylaminoarenes accessible.<sup>1</sup> The cyclization of *ortho*-alkynylbenzenediazonium salts, discovered by Richter,<sup>2</sup> has been applied to the preparation of 4-hydroxycinnoline derivatives (*Eq. 1*).<sup>3,4</sup>



In the standard procedure, the conversion of anilines **1** into cinnolines **3** can be realized in a one-pot preparation. At the same time, the Richter reaction for the preparation of some cinnoline derivatives is limited because of its rigorous conditions. Acidity and high temperature lead to poor results in the conversion of *vic*-alkynylaminoarenes because of the concurrent hydrolysis of some functional groups. Transformation of the amino group (compound **1**) to the diazonium salt (compound **2**) enhances the reactivity of substituents; for example, all attempts to perform the cyclization of the 3-diethylamino-6-(heptyn-1-yl)-1,4-naphthoquinone-5-diazonium