

Correction to Convergent Kilo-Scale Synthesis of a Potent Renin Inhibitor for the Treatment of Hypertension

Louis-Charles Campeau,* Sarah J. Dolman, Danny Gauvreau, Ed Corley, Jinchu Liu, Erin N. Guidry, Stéphane G. Ouellet, Dietrich Steinhuebel, Mark Weisel, and Paul D. O'Shea

2011, 15, 1138–1148; 10.1021/op2001063

The first three sentences of the third paragraph of page 1141 should read thusly, along with an additional reference at the end of the third sentence:

Olefin **16** was subjected to chiral hydrogenation previously developed in our laboratories. Initial conditions were Ru(cod)-(methallyl)₂/SL-J212-1 (10 mol % premixed Ru/ligand in DCM) at room temperature, 500 psig, in 2-MeTHF, with 1.2 equiv of HBF₄·OEt₂. Excess tetrafluoroboric acid was required to protonate the pyridyl-nitrogen, which otherwise acts as a catalyst poison.¹

■ REFERENCES

(1) Previous work on asymmetric hydrogenation of tetrasubstituted olefins shows that acid is also required for precatalyst formation: (a) Dobbs, D. A.; Vanhessche, K. P. M.; Brazi, E.; Rautenstrauch, V.; Lenoir, J.-Y.; Genet, J.-P.; Wiles, J.; Bergens, S. H. *Angew. Chem., Int. Ed.* **2000**, *39*, 1992. (b) Wiles, J. A.; Bergens, S. H.; Vanhessche, K. P. M.; Dobbs, D. A.; Rautenstrauch, V. *Angew. Chem., Int. Ed.* **2001**, *40*, 914, 1573. (c) Dupau, P.; Bruneau, C.; Dixneuf, P. H. *Adv. Synth. Catal.* **2001**, *343*, 331. (d) Tang, W.; Wu, S.; Zhang, X. *J. Am. Chem. Soc.* **2003**, *125*, 9570.

Published: May 29, 2012

