

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

CORRIGENDUM

To cite this Article (1994) 'CORRIGENDUM', Organic Preparations and Procedures International, 26: 3, 391 – 392

To link to this Article: DOI: 10.1080/00304949409458447

URL: <http://dx.doi.org/10.1080/00304949409458447>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>


This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Corrigendum

Volume 25, p. 708 (1993):

structure 3 should be as follows:



Volume 25, p. 709 (1993):

line 6 should read: nm (log ϵ 4.14-4.02).²

Table should read:

TABLE 1. Yields and Physical Constants of 2-Imidazolones 2^a

Compd	mp. (°C)	Yield (%)	Time (hrs)	¹ H NMR (δ)	Elemental Analyses (Found)		
					C	H	N
2a	206	75	2	7.23-8.19 (m, 5H, Ar-H) 10.53 (s, 2H, NH)	62.40 (62.10)	4.05 (4.17)	24.27 (24.41)
2b	165-166	70	3	2.33 (s, 3H, CH ₃), 7.14-8.16 (m, 4H, Ar-H), 10.38 (s, 2H, NH) D ₂ O-exchangeable	64.17 (63.90)	4.81 (5.10)	22.46 (22.54)
2c	190	70	4		64.17 (64.25)	4.81 (5.10)	22.46 (22.76)
2d	212	65	6	3.25 (s, 3H, OCH ₃), 7.16-8.00 (m, 4H, Ar-H) 10.23(broad, 2H, NH) D ₂ O-exchangeable	59.11 (58.90)	4.43 (4.67)	20.68 (20.38)
2e	160	40	6	6.66-7.93 (m, 4H, Ar-H) 8.60 (s, 2H, NH), D ₂ O- exchangeable	52.04 (51.81)	2.89 (3.02)	20.24 (20.54)
2f	242	70	3	6.85-8.05 (m, 4H, Ar-H), 8.57 (s, 2H, NH) ppm	52.04 (52.15)	2.89 (2.92)	20.24 (20.24)
2g	120	70	3	8.14-9.14 (m, 4H, Ar-H), 12.16 (s, 2H, NH) ppm	49.54 (49.40)	2.77 (2.65)	25.68 (25.40)

a) Except for colorless **2e**, all compounds are yellow and were crystallized from ethanol.

lines 4 and 5 of **EXPERIMENTAL SECTION** should read:

spectrophotometer as nujol mulls. The ¹H NMR spectra were run on JEOL FT-NMR FX-90Q spectrometer at the probe temperature (27°) in dimethyl sulfoxide-*d*₆ solution using TMS as an internal

Vol. 25, p. 710 (1993):

line 4 should read:

1-Methyl-3-(5-phenyl-2H-imidazol-2-one-4-yl)-2-thiourea (3a). Typical Procedure.- 4-Amino-5-

line 6 should read:

0.01 mole) in dry benzene (20 mL) for 1 hr on a steam bath; a solid mass was formed. The solvent

line 13 should read:

amino-5-phenyl-2H-imidazol-2-one in 30 mL ethanol was treated with freshly prepared 2,4-dinitro-

reference 3 should read:

3. a) T. S. Oakwood and C. A. Weisgerber, *Org. Syn. Coll. Vol. III*, 112 (1955); b) F. Asinger, A. Saus, H. Offermanns and H. D. Hahn, *Ann.*, 922 (1966); c) H. Brachwitz and K. Werner, Ger. (East) Patent 64276 (1968); *Chem. Abstr.*, 38610c, (1969); d) R. L. Soulen, S. C. Carlson and F. Lang, *J. Org. Chem.*, 479 (1973); e) J. F. Normant and C. Piechucki, *Bull. Soc. Chim. Fr.*, 2402 (1972).