

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

A FACILE SYNTHESIS OF 9-HYDROXYBENZO[a]PYRENE

Shoujun Chen^{ab}; Chunhua Wang^a; Peter P. Fu^a

^a Division of Biochemical Toxicology, National Center for Toxicological Research, Jefferson, AR, USA ^b

Department of Medicinal Chemistry, College of Pharmacy, The University of Michigan, Ann Arbor, MI, USA

To cite this Article Chen, Shoujun , Wang, Chunhua and Fu, Peter P.(1997) 'A FACILE SYNTHESIS OF 9-HYDROXYBENZO[a]PYRENE', *Organic Preparations and Procedures International*, 29: 1, 131 – 134

To link to this Article: DOI: 10.1080/00304949709355177

URL: <http://dx.doi.org/10.1080/00304949709355177>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

6. J. H. Looker, *J. Org. Chem.*, **17**, 510 (1952).
7. A. Ault, "*Techniques and Experiments for Organic Chemistry, 3rd Edition*" (Allyn and Bacon, Boston: 1973) p. 236-237.
8. An Aldrich "coffee pot" Kugelrohr, cat. Z10,046-3, was used. The receiving bulb was cooled in an ice bath.
9. The product at this point exhibited satisfactory ¹H NMR and GC-MS spectra, and had a mp. 100-102°.


A FACILE SYNTHESIS OF 9-HYDROXYBENZO[a]PYRENE

Submitted by Shoujun Chen^{*†}, Chunhua Wang and Peter P. Fu
(03/05/96)

*Division of Biochemical Toxicology
National Center for Toxicological Research
Jefferson, AR 72079, USA*

One important way to elucidate the mechanisms of metabolic activation of polyaromatic hydrocarbons is to study the DNA adducts formed thereof.¹ Synthetic DNA adducts are also molecular biomarkers for measuring human exposures to chemical carcinogens.² It is well documented that the *bay-region* diol epoxides of BAP^{**} are the principal active forms of this carcinogen³ and the key step in its tumor initiation is thought to be the covalent binding of these epoxides to DNA.⁴ Although the corresponding K-region epoxides were shown not to be responsible for the binding to DNA found in cellular systems,⁵ evidence indicated that 9-hydroxy-BAP-4,5-oxide might also be an active metabolite of BAP which could bind to nucleic acids *in vivo*.⁶⁻⁸ In order to assess the human health risk posed by BAP, it is necessary and timely to synthesize the DNA adduct of 9-hydroxy-BAP-4,5-oxide as a biomarker. The key intermediate is 9-hydroxy BAP (**6**) which was synthesized by Harvey,⁶ Yagi⁹ and Sims¹⁰ from 9,10-dihydrobenzo(a)pyren-7(8H)-one. However, these syntheses require either several steps under harsh conditions^{9,10} or the use of expensive and hazardous osmium tetroxide.⁶

We now report here a facile synthetic route to 9-hydroxy-BAP from a relatively more economical starting material, 7-hydroxy-7,8,9,10-tetrahydro-BAP (**1**) which was converted to 7-*p*-chlorobenzoyloxy-7,8,9,10-tetrahydro-BAP (**2**) quantitatively by treatment with *p*-chlorobenzoyl chloride, pyridine and catalytic amount of DMAP in chloroform at room temperature. 7-*p*-Chlorobenzoyloxy-7,8-dihydro-BAP (**3**) can be obtained selectively when **2** is treated with DDQ in hot benzene. The process can be easily monitored by TLC on silica gel.



It should be pointed out that, in our hands, both 7,8,9,10-tetrahydro-BAP and the 7-acetoxy derivative of **2** lack this selectivity. Treatment of either compounds with DDQ leads to the formation of BAP without control. The arene **3** thus obtained can be employed in the Prevost reaction without isolation to give **4** in a "one pot" procedure. After debenzoylation¹¹ and dehydration^{12,13} in the usual manner, 9-hydroxy-BAP (**6**) can be obtained with a total yield of 31% (from **1**). The structure of **6** was confirmed by the comparison of its UV/VIS and proton NMR spectra with those of an authentic sample prepared through other methods.^{6,14} The experimental result obtained here is consistent with those predicted by molecular orbital theoretical calculation.¹⁵

EXPERIMENTAL SECTION

Melting points were measured on an Electrothermal melting point apparatus and are not corrected. UV/VIS spectra were obtained on a Beckman DU-65 spectrophotometer; ¹H-NMR spectra were taken with a Varian Gemini-300 instruments. Silver benzoate and 7,8,9,10-tetrahydro-7-hydroxy BAP (**1**) were purchased from Aldrich Chem. Co. The former was recrystallized from water before use.

7-(p -Chlorobenzoyloxy)-7,8,9,10-tetrahydro-BAP (2) was prepared from **1** and p -chlorobenzoyl chloride in chloroform in quantitative yield, mp 154-156°;

Anal. Calcd. for $\text{C}_{27}\text{H}_{19}\text{ClO}_2$: C, 78.92; H, 4.67. Found: C, 78.57; H, 4.67

¹H NMR(CDCl_3): δ 2.12-2.24 (m, 2 H_9), 3.30-3.44 (m, 2 H_8), 3.61-3.72 (dt, 2 H_{10}), 6.66-6.72 (t, 1 H_7), 7.3-8.3 (m, 12 H_{aryl}).

7-(p -Chlorobenzoyloxy)-9,10-dibenzoyloxy-7,8,9,10-tetrahydrobenzo-[a]pyrene (4).- DDQ (0.12 g, 0.53 mmol) was added to a hot solution of **2** (0.22g, 0.54 mmol) in benzene (20 mL) under N_2 , and the mixture was stirred at 70°. The progress of the reaction was monitored by TLC. After consumption of the starting material (*ca* 40 min), the hot solution was filtered quickly through a column of silica gel and Celite, and washed thoroughly with benzene. The filtrate was concentrated to *ca* 20 mL

under reduced pressure without heating. In another flask, a mixture of silver benzoate (0.28g, 1.2 mmol) and iodine (0.14 g, 0.55 mmol) in benzene (20 mL) was heated to reflux for 30 min. To this yellow suspension was added the above benzene solution of **3** and the reaction mixture was heated to reflux for 1hr. After being cooled, the reaction mixture was filtered through a layer of celite. The filtrate was concentrated, chromatographed on silica gel and eluted with a mixture of hexane and ethyl acetate (10:1, v/v) to give 0.18g (52% based on **2**) of **4** as a light yellow solid, mp 201-203°;

Anal. Calcd. for $C_{41}H_{27}ClO_4$: C, 79.53; H, 4.40. Found: C, 79.34; H, 4.52

1H NMR ($CDCl_3$): δ 0.8-0.9 (m, 1 H_{8a}), 1.2-1.3 (m, 1 H_{8b}), 6.0 (d, 1 H_{10} , $J = 2.4$ Hz), 6.9-7.0 (m, 1 H_9), 7.2-8.4 (m, 25 H, $H_{aryl} + H_7$).

7,8,9-Trihydroxy-7,8,9,10-tetrahydrobenzo[a]pyrene (5).- To a stirred solution of **4** (0.18 g, 0.28 mmol) in THF (5 mL) and methanol (95 mL) was added sodium methoxide (86 mg, 1.6 mmol) and the resultant solution was heated to reflux for 0.5 hr, and then cooled to room temperature; 15 mL of cold water was added to the reaction solution which was stirred at rt for 1 hr and extracted with EtOAc (3 x 15 mL). The EtOAc extracts was washed with water (2 x 10 mL). Removal of EtOAc under reduced pressure followed by chromatography on Florisil^R afforded 0.06 g (72%) of **5** as a white solid, mp 215-216° (dec);

Anal. Calcd. for $C_{20}H_{16}O_3$: C, 78.92; H, 5.31. Found: C, 78.61; H, 5.49

1H NMR (CD_3OD): δ 2.2-2.3 (m, 1 H_{8a}), 2.55-2.65 (m, 1 H_{8b}), 4.35-4.45 (m, 1 H_9), 5.23 (t, 1 H_7 , $J = 4.7$ Hz), 5.58 (d, 1 H_{10} , $J = 3.8$ Hz), 8.0-8.3 (m, 9H, 8 $H_{aryl} + OH$), 8.61 (s, 1H, OH), 8.64 (s, 1H, OH).

9-Hydroxybenzo[a]pyrene (6).- A solution of **5** (54 mg, 0.18 mmol) and $PTSH_2O$ (34 mg, 0.18 mmol) in dry benzene (20 mL) was heated to reflux under an atmosphere of argon for 1 hr, and it was then allowed to cool to rt, washed with water (2 x 10 mL), dilute sodium bicarbonate solution (1 x 10 mL) and water (10 mL). Evaporation of the benzene solution followed by chromatography on silica gel afforded 0.04 g (84%) of **6** as a crystalline solid, mp 194-196°, lit.¹⁰ mp 196°; UV/VIS spectra were identical with those reported.^{10,16} 1H NMR (THF- d_8): δ 7.44 (d, 1 H_8 , $J = 8$ Hz), 7.80-8.26 (m, 7 H), 8.39 (s, 1 H_{10}), 8.51 (s, 1 H_9), 9.01 (d, 1 H_{11} , $J = 9$ Hz), 9.11 (br, 1 H, OH).

REFERENCES

** The abbreviations used are: BAP, Benzo[a]pyrene; DDQ, 2,3-Dichloro-5,6-dicyano-1,4-benzoquinone; DMAP, 4-Dimethylaminopyridine; EtOAc, Ethyl acetate; $PTSH_2O$, *p*-Toluenesulfonic acid hydrate.

† Current address: Department of Medicinal Chemistry, College of Pharmacy, the University of Michigan, Ann Arbor, MI 48109-1055, USA.

1. A. Dipple, R. C. Moschel and C. A. Bigger, "Polycyclic Aromatic Hydrocarbons: Chemical Carcinogens", ed. C.E.Searles, pp 41-163, Washington D.C., American Chemical Society (1984).
2. a) R. W. Giese, in: *Biomonitoring and Carcinogen Risk Assessment*, R. C. Garner, P. B. Farmer, G. T. Steel, A. S. Wright, eds., pp 247-253, Oxford University Press, New York, NY (1991); b)

- US Dept. of Health and Human Service, *Environ. Health Perspect.*, **99**, 3 (1993).
3. R. G. Harvey, *Acc. Chem. Res.*, **14**, 218 (1981).
 4. S. C. Cheng, B. D. Hilton, J. M. Roman and A. Dipple, *Chem. Res. Toxicol.*, **2**, 334 (1989).
 5. A. Dipple, in "DNA Adduct: Identification and Biological Significance", pp 107-129, eds. K. Hemminki, A. Dipple, D. E. G. Shuker, F. F. Kadlubar, D. Segerback and H. Bartsch, LARC Scientific Publication No.125, Lyon, Int. Agency for Research on Cancer, LARC (1994).
 6. R. G. Harvey and C. Cortez, *Carcinogenesis*, **4**, 941 (1983).
 7. W. Bochnitschek, A. Seidel, H. Kung and F. Oesch, *Angew. Chem. Int. Ed. Engl*, **24**, 699 (1985).
 8. M. Degawa, S. J. Stern, M. V. Martin, F. P. Guengerich, P. P. Fu, K. F. Ilett, R. K. Kaderlik and F. F. Kadlubar, *Cancer Research*, **54**, 4915 (1994).
 9. H. Yagi, G. M. Holder, P. M. Pansette, O. Hernandez, H. J. Yeh, R. A. LeMahieu and D. M. Jerina, *J. Org. Chem.*, **41**, 977 (1976).
 10. P. Sims, *J. Chem. Soc. C*, 32 (1968).
 11. Y.-S. Wu, J.-S. Lai and P. P. Fu, *J. Org. Chem.*, **58**, 7283 (1993).
 12. P. P. Fu and R. G. Harvey, *J. Org. Chem.*, **48**, 1534 (1983).
 13. S. K. Yang, P. P. Roller and H. V. Gelboin, *Biochemistry*, **16**, 3680 (1977).
 14. Compound **6** was converted to 9-acetoxy-BAP in 92% yield by stirring with Ac₂O and pyridine at rt for 12 hrs. 9-Acetoxy-BAP has been previously used as a precursor for the preparation of 9-hydroxy-BAP.⁶
 15. P. P. Fu and R. G. Harvey, *Tetrahedron*, **34**, 857 (1978).
 16. R. I. Freudenthal and P. Jones (Eds), "Carcinogenesis: A Comprehensive Survey", p 410, Raven Press, 1976.