


This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

SYNTHESIS OF SUBSTITUTED 1-TETRALONES

Punit Kumar^a

^a Indian Institute of Chemical Technology, Hyderabad, INDIA

To cite this Article Kumar, Punit(1997) 'SYNTHESIS OF SUBSTITUTED 1-TETRALONES', *Organic Preparations and Procedures International*, 29: 4, 477 – 480

To link to this Article: DOI: 10.1080/00304949709355222

URL: <http://dx.doi.org/10.1080/00304949709355222>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.


The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

SYNTHESIS OF SUBSTITUTED 1-TETRALONES

Submitted by Punit Kumar
(05/17/96)


Indian Institute of Chemical Technology
Hyderabad 500 007, INDIA

Substituted 1-tetralone derivatives (**1**) are valuable intermediates in organic synthesis. In fact, 6-methoxy-1-tetralone¹ (**2**), 1-tetralone² (**3**), and 5,6-methylenedioxy-1-tetralone³ are the starting materials in the synthesis of several complex biologically active compounds. In 1972, Heck reported the conjugate addition of benzyl chloride to methyl acrylate, promoted by Pd(OAc)₂, leading to the formation of methyl 4-phenyl-3-butenolate (69%) and methyl 4-phenyl-2-butenolate (9%)⁴. Since




that time, relatively little interest has been shown in this reaction. The existing approaches⁵ to the synthesis of substituted 1-tetralone derivatives require long or complex reaction sequences, sometimes leading to the formation of isomeric compounds. For instance, preparation of compound **5** by Friedel-Crafts acylation of benzene with methylsuccinic anhydride followed by reduction of one of the benzylic ketones leads to an isomeric mixture. As a part of our current interest in abortifacient drug Mifepristone⁶ and ABT-200³, a potent α -2 antagonist and norepinephrine uptake inhibitor currently undergoing testing at Abbot Labs, we developed an alternate process, based on the Heck reaction to prepare 6-methoxy-1-tetralone (**2**), 1-tetralone (**3**), 5,6-dimethoxy-1-tetralone (**4**) and 2-methyl-1-tetralone.

3-Methoxybenzyl chloride (**6**) and ethyl acrylate were heated at 100° in the presence of Pd(OAc)₂ and tributylamine for 9 h to give **7** in 85% yield. The ¹H NMR spectrum of **7** revealed the presence of 3- and 2-butenates as regioisomers. Subsequent reduction of **7** over Pd-C in methanol at




a) Pd(OAc)₂, Bu₃N, 100°, 9 h; b) i) H₂, Pd-C, MeOH, 50 psi, 5 h; ii) 6N, KOH, EtOH, Δ , 2 h;
c) PPE, CHCl₃, RT, 7 h.

50 psi followed by hydrolysis gave the acid **8**. Compound **8** was treated with polyphosphoric ester (PPE)⁷ in CHCl_3 at room temperature to give **2** (77.8% overall yield) along with 8-methoxy-1-tetralone (**9**) (8%). Similarly, benzyl chloride (**10**) was converted into 1-tetralone (**3**) (overall yield 77%). The synthesis of 5,6-dimethoxy-1-tetralone (**4**) (overall yield 67%) was also completed by the above approach starting from 2,3-dimethoxybenzyl chloride (**11**)⁸.



In order to prepare 2-methyl-1-tetralone (**5**), the reaction of benzyl chloride (**10**) and ethyl methacrylate in the presence of $\text{Pd}(\text{OAc})_2$ at 100° for 11 h followed by catalytic reduction produced the product **12**. Subsequent ester hydrolysis of **12** and cyclization with PPE- CHCl_3 gave **5** (overall 51.6% yield).



EXPERIMENTAL SECTION

Proton magnetic resonance spectra were recorded in CDCl_3 on Varian FT-200 MHz (Gemini) spectrophotometer using TMS as the internal standard. Infra Red spectra were scanned on Shimadzu IR-470 with sodium chloride optics. Mass spectra were recorded on VG 7070 H. Melting points were recorded on Buchi 535 melting point apparatus and are uncorrected. All evaporations were carried under reduced pressure on rotary evaporators below 50° .

Ethyl 4-(3-Anisyl)-2-butenolate and Ethyl 4-(3-Anisyl)-3-butenolate (7).- A mixture of 3-methoxybenzylchloride (15.6 g, 100 mmol), $\text{Pd}(\text{OAc})_2$ (0.24 g, 1 mmol), tri-*n*-butylamine (37.07 g, 200 mmol) and ethyl acrylate (15.0 g, 150 mmol) was heated at 100° for 9 h. The reaction mixture was poured in water and extracted with ethyl acetate. The organic layer was washed with 1N HCl, water, dried (Na_2SO_4) and concentrated. The residue was chromatographed on silica gel by using ethyl acetate:light petroleum (5:95) to give **7** as a pale brown viscous oil, isolated in 85% yield (18.7 g). ^1H NMR: δ 1.28 (t, 3H, $J = 7.3$ Hz), 3.21 (d, 2H, $J = 5.8$ Hz), 3.80 (s, 3H), 4.16 (q, 2H, $J = 7.3$ Hz), 6.15-7.3 (m, 6H).

Anal. Calcd. for $\text{C}_{13}\text{H}_{16}\text{O}_3$: C, 70.88; H, 7.32. Found: C, 70.77; H, 7.34

6-Methoxy-1-tetralone (2).- Compound **7** (18.7 g, 84.09 mmol) and 10% Pd-C (2 g) in methanol (100 mL) was hydrogenated at 50 psi for 5 h. After the catalyst was filtered through celite, the filtrate was diluted with 6N KOH (50 mL) and heated under reflux for 2 h. The reaction was acidified with concentrated HCl and methanol was removed under reduced pressure. The aqueous layer was extracted with ethyl acetate, dried (Na_2SO_4) and concentrated. The residue was dissolved in chloro-

form (100 mL) and PPE (19 g) was added. The resulting mixture was stirred at room temperature for 7 h, decomposed with water. The chloroform layer was separated, dried (Na_2SO_4), concentrated and the residue was chromatographed on silica gel by using ethyl acetate:light petroleum (5:95) to give **2** as a white crystalline solid, mp. 76-76.6°; lit.^{5f} mp. 75-77°, isolated in 92% yield (13.7 g). $^1\text{H NMR}$: δ 2.04 (m, 2H), 2.50 (t, 2H, $J = 6.3$ Hz), 2.86 (t, 2H, $J = 6.3$ Hz), 3.77 (s, 3H), 6.61 (d, 1H, $J = 1.5$ Hz), 6.73 (dd, 1H, $J = 1.5$ Hz, 9.0 Hz), 7.93 (d, 1H, $J = 9$ Hz). IR (CHCl_3): 1670 cm^{-1} . MS: m/z 176 (M^+).

Further elution gave 8-methoxy-1-tetralone as a colorless liquid, bp. 96-98°/ 0.02 mm; lit.⁹ bp. 94-96° / 0.02 mm isolated in 8% yield (1.1g). $^1\text{H NMR}$: δ 2.06 (m, 2H), 2.61 (t, 2H, $J = 6.2$ Hz), 2.91 (t, 2H, $J = 6.2$ Hz), 3.89 (s, 3H), 6.78 (m, 2H), 7.34 (t, 1H, $J = 8.0$ Hz).

1-Tetralone (3), isolated as a light yellow liquid, bp. 136-138°/ 15 mm; lit.¹⁰ bp. 135-137° / 15 mm in 77% yield. $^1\text{H NMR}$: δ 2.06 (m, 2H), 2.56 (t, 2H, $J = 6.8$ Hz), 2.88 (t, 2H, $J = 6.8$ Hz), 7.13-7.24 (m, 2H), 7.36 (m, 1H), 7.95 (d, 1H, $J = 8.1$ Hz). IR (Neat): 1680 cm^{-1} ; MS: m/z 146 (M^+).

5,6-Dimethoxy-1-tetralone (4), white crystalline solid, mp. 103-104.6°; lit.¹¹ mp. 104-105°, isolated in 67% yield. $^1\text{H NMR}$: δ 2.02-2.17 (m, 2H), 2.56 (t, 2H, $J = 6.4$ Hz), 2.94 (t, 2H, $J = 6.4$ Hz), 3.79 (s, 3H), 3.90 (s, 3H), 6.82 (d, 1H, $J = 8.6$ Hz), 7.78 (d, 1H, $J = 8.6$ Hz). IR (CHCl_3): 1665 cm^{-1} ; HRMS: 206.0944. Calcd. for $\text{C}_{12}\text{H}_{14}\text{O}_3$: 206.0942.

Ethyl 2-Methyl-4-phenylbutenoate (12).- A mixture of benzyl chloride (12.6 g, 100 mmol), $\text{Pd}(\text{OAc})_2$ (0.24 g, 1 mmol) and ethyl methacrylate (20 mL, 183 mmol) tri-*n*-butylamine (37.07 g, 200 mmol) was heated at 100° for 11 h. After usual work-up, the residue was hydrogenated over 10% Pd-C (1.5 g) at 50 psi in methanol (50 mL) for 6 h to give **12** as a colourless viscous oil, isolated in 60% yield (12.1 g). $^1\text{H NMR}$: δ 1.16 (d, 3H, $J = 6.4$ Hz), 1.25 (t, 3H, $J = 6.4$ Hz), 1.65 (m, 1H), 1.95 (m, 1H), 2.36 (m, 1H), 2.55 (t, 1H, $J = 6.4$ Hz), 4.08 (q, 2H, $J = 6.4$ Hz), 7.0-7.2 (m, 5H).

2-Methyl-1-tetralone (5).- Compound **12** (12.0 g, 58.2 mmol) was hydrolysed with 6N KOH (20 mL) and then cyclised with PPE (12 g) in chloroform (30 mL) as reported above to provide **5** as a colorless oil, isolated in 88% yield (8.26g). $^1\text{H NMR}$: δ 1.30 (d, 3H, $J = 6.0$ Hz), 1.7-3.1 (m, 5H), 7.15-7.4 (m, 3H), 8.0 (d, 1H, $J = 9$ Hz). IR (Neat): 1680 cm^{-1} ; HRMS: 160.0897. Calcd. for $\text{C}_{11}\text{H}_{12}\text{O}$: 160.0888.

Acknowledgments. - The author is thankful to CSIR for providing fellowship and to Dr. M. K Gurjar and Dr. T. R. Bhardwaj for providing the much needed encouragement.

REFERENCES

1. a) S. N. Ananchenko and I. V. Torgov, *Tetrahedron Lett.*, 1553 (1963); b) S. N. Ananchenko, V. Ye. Limanov, V. N. Leonov, V. N. Rzhazhnikov and I. V. Torgov, *Tetrahedron*, **18**, 1355 (1962); c) H. Bayer, C. Batzl, R. W. Hartman and A. J. Mannschreck, *J. Med. Chem.* **34**, 2685 (1991); d) C. Almansa, L. A. Gomez, F. L. Cavalcanti, R. Rodriguez, E. Carcellar, J. Bartroli, J. C. Rafanell and J. Forn, *ibid.*, **36**, 2121 (1993).
2. M. F. El-Zohry and A. M. El-Khawaga, *J. Org. Chem.*, **55**, 4036 (1990).
3. a) R. E. Zella, A. A. Hancock, S. A. Buckener, F. Z. Basha, K. Tietze, J. F. DeBarnadis and M. D. Meyer, *Bioorganic and Medicinal Chem. Lett.*, **4**, 1319 (1994); b) M. Deshpande, *Interna-*

- tional Conference on Organic Synthesis-10*, held at Bangalore, India from December 12-16, 1994, Page 87.
4. a) R. F. Heck and J. P. Nolley, Jr., *J. Org. Chem.*, **37**, 2320 (1972); b) R. F. Heck, *Org. React.*, **27**, 345 (1982).
 5. a) H. W. Thompson, *J. Chem. Soc.*, 2310 (1932); b) V. C. E. Burnop, G. H. Elliot and R. P. Linstead, *ibid.*, 727 (1940); c) D. G. Thomas and A. N. Nathan, *J. Am. Chem. Soc.*, **70**, 331 (1948); d) W. S. Johnson and H. J. Glenn, *ibid.*, **71**, 1092 (1949); e) G. Stork, *ibid.*, **69**, 576 (1947); f) D. Papa, *ibid.*, **71**, 3246 (1949); g) D. Papa, E. Schwenk and H. Breiger, *J. Org. Chem.*, **14**, 366 (1949); h) D. Papa and E. Schwenk, US Patent 2, 475, 718, July 12, 1949; Chem. Abstr., **43**, 7510i (1949).
 6. a) A. Ulmann and C. Dubois, *Acta Obstet. Gynecol. Scand. Suppl.* **149**, 9 (1989). b) O. Heikinheimo, O. Ylikorkala and P. Lahtenmaki, *Ann. Med.* (Hagerstown, Md), **22**, 75 (1990).
 7. M. P. Cava, M. V. Lakshmikantham and M. J. Mitchell, *J. Org. Chem.*, **34**, 2665 (1969).
 8. Compound **11** was prepared in two steps by reduction of 2,3-dimethoxybenzaldehyde with sodium borohydride in methanol (88%) followed by treatment of the alcohol for 6 h with triphenylphosphine in refluxing carbon tetrachloride (85%).
 9. B. Tarnchompoo, C. Thebtaranonth and Y. Thebtaranonth, *Synthesis*, 785 (1986).
 10. *Org. Synth.*, Coll. Vol., **4**, 900.
 11. N. F. Elmore and T. J. King, *J. Chem. Soc.*, 4425 (1961).

DETHIOACETALIZATION OF DITHIOLANES WITH FERRIC NITRATE AND SILICA GEL IN HEXANE

Submitted by Masao Hirano*, Ken Ukawa, Shigetaka Yakabe and Takashi Morimoto*
(12/09/97)

*Department of Applied Chemistry, Faculty of Technology
Tokyo University of Agriculture and Technology
Koganei, Tokyo 184, JAPAN*

The importance of protection/deprotection processes for certain functionalities in the synthesis of organic compounds has led to the development of a large numbers of protective groups and of methods for their removal.¹ Of the typical deprotective procedures (*e.g.* chemically, photolyti-