
This article was downloaded by:

On: 27 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information: http://www.informaworld.com/smpp/title~content=t902189982

α,β-UNSATURATED SULFONES BY THE WITTIG REACTION OF STABLE YLIDE WITH ALDEHYDES UNDER MICROWAVE IRRADIATION

Cong Fu^a; Caiding Xu^a; Zhi-Zhen Huang^a; Xian Huang^a

^a Department of Chemistry, Hangzhou University, Zhejiang, PR CHINA

To cite this Article Fu, Cong , Xu, Caiding , Huang, Zhi-Zhen and Huang, Xian(1997) ' α , β -UNSATURATED SULFONES BY THE WITTIG REACTION OF STABLE YLIDE WITH ALDEHYDES UNDER MICROWAVE IRRADIATION', Organic Preparations and Procedures International, 29: 5, 587 — 589

To link to this Article: DOI: 10.1080/00304949709355238 URL: http://dx.doi.org/10.1080/00304949709355238

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: http://www.informaworld.com/terms-and-conditions-of-access.pdf

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Volume 29, No. 5, 1997 **OPPI BRIEFS**

6. G. Bram, J. Sansoulet, H. Galons, Y. Bensaid and C. Combet-Farnoux and M. Miocque, Tetrahedron Lett., 26, 4601 (1985).

- 7. a). J. H. Fendler, "Membrane Mimetic Chemistry", Wiley Interscience, New York, 1982; b) M. Gratzel and K. Kalyanasundaram, "Surfactant Solutions: Kinetics and Catalysis in Microheterogeneous systems", Surfactant Science Series, Vol 38, Marcel Dekker, New York, 1991.
- 8. a) D.A. Jaeger and M. R. Frey, J. Org. Chem., 47, 311 (1982); b) V. Ramesh and V. Ramamoorthy, J. Org. Chem., 49, 536 (1984); c) R. Germani, P. Ponti, G. Savelli, N. Spreti, C. A. Bunton and J. R. Moffat, J. Chem. Soc., Perkin Trans II, 401 (1989); d) L.Wei, A. Lucas, J. Yue and B. R. Lennox, Langmuir., 7, 1336 (1991) e) C. D. Mudaliar and S. H. Mashraqui, J. Chem. Res. (S), 174 (1994).
- 9. However, ethyl cyanoacetate failed to react with either benzylidene acetophenone or benzylidene acetone under our standard micellar conditions. To understand this unusual result, we are presently investigating these reactions at higher concentration of alkali and micelles.
- 10. K R. Nivalkar, C. D. Mudaliar and S. H. Mashraqui, J. Chem. Res. (S), 98 (1992).
- 11. A. Garcia-Raso, J. Garcia Raso, B. Campaner, R. Mestres and J.V. Sinisterra, Synthesis, 1037 (1982).
- 12. P. Laszlo, M. Moutaufier and L. S. Randriamahefa, Tetrahedron Lett., 31, 4867 (1990).
- 13. K. Watanabe, K. Miyazu and K. Irie, Bull. Chem. Soc. Jpn, 55, 3212 (1982).

α,β-UNSATURATED SULFONES BY THE WITTIG REACTION OF STABLE YLIDE WITH ALDEHYDES UNDER MICROWAVE IRRADIATION

Submitted by (01/17/96)

Cong Fu, Caiding Xu, Zhi-Zhen Huang and Xian Huang*

Department of Chemistry, Hangzhou University

Zhejiang 310028, P. R. CHIN A

α,β-Unsaturated sulfones are important intermediates in organic synthesis and may be obtained from the condensation of arylsulfonylacetic acid with aldehydes.² Kobayashi et al. found that tosyl chloride underwent addition with vinylarenes catalyzed by a ruthenium(II) complex; subsequent dehydrochlorination gave the trans-α,β-unsaturated sulfones.³ The reduction of E-β-iodo-α,β-unsaturated sulfones with zinc in acetic acid afforded Z-tosyl-2-arylethenes. α,β-Unsaturated sulfones may also be obtained from the phase-transfer catalyzed condensation of sulfones with aldehydes.⁵ Many

OPPI BRIEFS Volume 29, No. 5, 1997

reactions, such as the Diels-Alder,⁶ Claisen,⁶ ene reactions,⁶ dehydration and nucleophilic substitutions⁷ may be conducted safely in commercial microwave ovens with remarkable rate enhancements

and dramatic reduction of reaction times compared with conventional heating. Previous papers described the synthesis of ethyl cinnamates and chalcones *via* microwave-induced Wittig reactions, and this paper reports the synthesis of α,β -unsaturated sulfones *via* the Wittig reaction under microwave irradiation.

As is well known, the Wittig reaction of stable ylides proceeds slowly using the conventional methods; ^{9,10} for example, the condensation of *p*-tosylmethylene triphenylphosphorane with *p*-nitrobenzaldehyde in chloroform required three days at room temperature to give a 53% yield of 1-(*p*-nitrophenyl)-2-(*p*-tosyl)ethylene. ⁹ Our experiments showed that, *p*-tosylmethylene triphenylphosphorane absorbed on silica gel, underwent the Wittig reaction completely under the irradiation of microwave after 15 min. to afford the *trans* isomer as the sole products. Silica gel was used as an effective support for this reaction, without which the yield of the Wittig reaction decreased dramatically (26%). The *trans* configuration of products was established by ¹H NMR spectra in which the coupling constants of vinylic hydrogens were in the range of 15-16Hz. This method was also adapted to aliphatic aldehydes.

TABLE. Synthesis of α,β-Unsaturated Sulfones

Aldehyde	Product	Yield (%)	mp. (°C)	lit. mp. (°C)
C ₆ H ₅ CHO	C ₆ H ₅ CH=CHTs	66	120-121	120-1213
4-NO ₂ C ₆ H ₄ CHo	4-NO ₂ C ₆ H ₄ CH=CHTs	58	182-184	183-184 ¹⁰
4-CIC ₆ H ₄ CHO	4-CIC ₆ H ₄ CH=CHTs	63	151-152	151-152 ³
3-NO ₂ C ₆ H ₄ CHO	3-NO ₂ C ₆ H ₄ CH=CHTs	57	145-146	146-147 ³
4-CH ₃ C ₆ H ₄ CHO	4-CH ₃ C ₆ H ₄ CH=CHTs	70	154-155	154-155 ³
2.4-C1 ₂ C ₆ H ₃ CHO	2,4-Cl ₂ C ₆ H ₃ CH=CHTs	57	128-129	129-129.5 ³
(CH ₃) ₃ CCH ₂ CHO	(CH ₃) ₃ CCH ₂ CH=CHTs	53	oil ^a	oil ¹¹
C ₆ H ₅ CH ₂ CHO	C ₆ R ₅ CH ₂ CH=CHTs	62	118-120	12011

a) ¹H NMR (CDCl₃): δ 1.0 (s, 9H), 2.2-2.5 (m, 5H), 6.4 (d, 1H, J = 15Hz), 6.8 (d, 1H, J = 15Hz), 7.3-7.8 (m, 4H)

EXPERIMENTAL SECTION

Melting points are uncorrected. ¹H NMR spectra were obtained with F 90Q spectrometer in CDCl₃ using TMS as internal standard. National NN-5252 domestic microwave oven was used.

Typical Procedure.- A mixture of *p*-tosylmethylene triphenylphosphorane⁹ (1mmol), benzaldehyde (1mmol), and silica gel (2g, 200-300 mesh) were introduced into the domestic microwave oven in an

Volume 29, No. 5, 1997 OPPI BRIEFS

open round-bottomed flask irradiated with microwave at an output of about 120w for 15 min. Then 20 mL methylene dichloride was added into the cooled mixture, the extract was concentrated and chromatographed on silica gel (200 mesh), using 10:1 (v/v) benzene-ethyl acetate as eluent to give the product (150mg, 66%) as a yellow solid.

Acknowledgment. - This research was supported by the NSF of Zhejiang.

REFERENCES

- P. L. Fuchs and T. F. Braish, Chem. Rev., 86, 903 (1986); P. D. Magnus, Tetrahedron, 33, 2019 (1977); L. Field, Synthesis, 713 (1978).
- M. Balasubramanian, V. Baliah and T. Rangarajan, J. Chem. Soc., 1844 (1954); M. Balasubramanian and V. Baliah, ibid., 3296 (1955).
- 3. N. Kamigata, H. Sawada and M. Kobayashi, J. Org. Chem., 48, 3793 (1983).
- 4. W. E. Truceand G. C.Wolf, *ibid.*, **36**, 1727 (1971); C. Correa and M. Fleming, *J. Chem. Soc.*, *Perkin Trans.* 2, 103 (**1987**).
- 5. G. Cardillo, D. Savoia and A. Umani-Ronchi, Synthesis, 453 (1975).
- R. J. Giguere, T. L. Bray, S. M. Duncan and G. Majetich, *Tetrahedron Lett.*, 27, 4945 (1986); R. J. Giguere, A. M. Namen, B. Lopez, A. Arepally, D. E. Romos, G. Majetich and J. Defauw, *ibid.*, 28, 6553 (1987).
- R. J. Gedye, F. Smith, K. Westaway, H. Mi, L. Baldisera, L. Laberge and J. Rousell, *ibid.*, 27, 279 (1986); R. J. Giguere, "Organic Synthesis: Theory and Application, Vol. 1, p.103, JAI Press Inc. Greenwich, CT, 1989.
- 8. C. Xu, G. Chen and X. Huang, *Org. Prep. Proced. Inter.*, **27**, 559 (1995); C. Xu, Chen and X. Huang, *Synth. Commun.*, **25**, 2229 (1995).
- 9. A. J. Speziale and K. W. Ratts, J. Am. Chem. Soc., 87, 5603 (1965).
- 10. V. Baliah and M. Seshapathirao, J. Org. Chem., 24, 867(1959).
- 11. K. Inomata, T. Hirata, H. Suhara, H. Kinoshita, H. Kotake and H. Senda, *Chemistry Lett.*, **204**, 2009 (1988).
