

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

### AN IMPROVED SYNTHESIS OF *tert*-BUTYL $N^{\alpha}$ -*tert*-BUTOXYCARBONYL-L-(S-TRITYL)CYSTEINATE

Elżbeta Masiukiewicz<sup>a</sup>; Barbara Rzeszotarska<sup>a</sup>

<sup>a</sup> Department of Organic Chemistry, University of Opole, Opole, POLAND

**To cite this Article** Masiukiewicz, Elżbeta and Rzeszotarska, Barbara(1999) 'AN IMPROVED SYNTHESIS OF *tert*-BUTYL  $N^{\alpha}$ -*tert*-BUTOXYCARBONYL-L-(S-TRITYL)CYSTEINATE', *Organic Preparations and Procedures International*, 31: 5, 571 – 572

**To link to this Article:** DOI: 10.1080/00304949909355343

**URL:** <http://dx.doi.org/10.1080/00304949909355343>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


**AN IMPROVED SYNTHESIS OF *tert*-BUTYL  
N<sup>α</sup>-*tert*-BUTOXYCARBONYL-L-(S-TRITYL)CYSTEINATE**

Submitted by  
(09/09/99)

Elżbieta Masiukiewicz and Barbara Rzeszotarska

*Department of Organic Chemistry  
University of Opole  
Ul. Oleska 48, 45-052 Opole, POLAND*

The controlled synthesis of peptides requires that functional groups that are not directly involved in the amide bond-forming reaction be blocked. Acidic conditions<sup>1</sup> are preferred for the removal of protective groups since these conditions are less damaging to peptides. Therefore, it may be desirable that all protective groups be acid-sensitive, provided that the selective removal of either *N*- or *C*-protecting group is feasible.<sup>2</sup> Methods useful to this end are elaborated by testing on simple model compounds. One such useful model is the title compound. The literature refers its preparation to general procedures<sup>2</sup> and gives NMR spectra and elemental analysis as the only indicators of purity, while chromatographic criteria would be more appropriate for the compound that is to be used in chromatographic testing the methods of selective deprotection. Herein, we present a complete description of the preparation of Boc-Cys(Trt)-*Ot*Bu<sup>3</sup> in high yield and of high purity.



H-Cys(Trt)-OH was esterified with *tert*-butyl acetate and HClO<sub>4</sub> as a catalyst.<sup>4</sup> The crude product was contaminated with about 5% H-Cys(*t*Bu)-*Ot*Bu<sup>5</sup> and Trt-OH resulting from transesterification. While the former impurity could be removed by extraction with 0.5 M HCl, the latter could not be eliminated. The mixture was thus acylated with Boc<sub>2</sub>O in the presence of triethylamine.<sup>6</sup> Since the crude acylated product contained other contaminants and was very soluble in all solvents, it could not be purified by simple crystallization and a chromatographic column had to be used. This procedure allows Boc-Cys(Trt)-*Ot*Bu to be obtained as well-shaped crystals of 99.7% purity in 80% yield.

### EXPERIMENTAL SECTION

Reactions were monitored and the products checked on silica gel plates (DC Alufolien Kieselgel, 0.25 Merck # 5553) in the following solvent systems (v/v): A = chloroform-methanol-acetic acid (95:5:3), B = chloroform-*n* hexane-acetic acid (10:10:1). HPLC analyses were carried out using a Beckman System Gold chromatograph, a 5 μl loop, an Alltech Alltima, C<sub>18</sub>, 5 μ, 150 x 4.6 mm column, a flow rate of 1 mL/min and detection at 210 nm.

***tert*-Butyl L-(S-Trityl)cysteinate Hydrochloride (HCl·H-Cys(Trt)-*Ot*Bu).**- To a vigorously stirred

suspension of H-Cys(Trt)-OH (10.0 g, 27.2 mmol) in *tert*-butyl acetate (160 mL), 70% HClO<sub>4</sub> (8 mL) was introduced dropwise. Stirring was continued at room temperature for 1 h and ethyl acetate (200 mL) and 1 M aqueous NaHCO<sub>3</sub> to pH 8.0 were added. The precipitated, unreacted H-Cys(Trt)-OH was filtered off (1.55 g, 4.3 mmol) and the aqueous layer was discarded. The organic phase was extracted with 0.5 M HCl (2 x 100 mL) and brine, dried and evaporated to give the product (9.79 g, 88% yield) as a colorless rigid foam. R<sub>f</sub>: A - 0.48. HPLC: 0.1% trifluoroacetic acid-acetonitrile (30:70); HCl·H-Cys(Trt)-*Or*Bu (94%) tR = 2.85 min, Trt-OH (5%) tR = 5.35 min.

***tert*-Butyl N<sup>α</sup>-*tert*-Butoxycarbonyl-L-(S-trityl)cysteinate (Boc-Cys(Trt)-*Or*Bu).**- To a solution of 94% HCl·H-Cys(Trt)-*Or*Bu (5.02 g, 10.3 mmol) in dioxane (30 mL), was added Boc<sub>2</sub>O (3.64 g, 16 mmol) and followed by triethylamine (4.5 mL, 32 mmol), dropwise. After 15 h, triethylamine hydrochloride was filtered off and dioxane was evaporated; the residue was dissolved in ethyl acetate (30 mL) and extracted with 0.5 M HCl and brine and dried. After evaporation of ethyl acetate, the residue was dissolved in *n*-hexane, and applied to a silica gel column (ϕ 5 cm, Merck # 7736, 230 g) conditioned with *n*-hexane. The column was eluted under pressure in sequence with solutions of 1%, 2%, 5%, 10%, 15%, 20%, 25%, 30%, 40% and 50% of chloroform in *n*-hexane (each 50 mL). The appropriate fractions (by TLC) were collected and evaporated; the residue was dissolved in *n*-hexane (10 mL) and left standing in a refrigerator for several days to furnish Boc-Cys(Trt)-*Or*Bu (4.2 g, 80% yield), homogenous by TLC, mp. 80-82° (uncorrected, a Boëtius heating block). R<sub>f</sub>: A - 0.72. HPLC: water-acetonitrile (15:85), tR = 7.38 min, 99.7% purity.

*Anal.* Calcd. for C<sub>31</sub>H<sub>37</sub>NO<sub>4</sub>S: C, 71.64, H, 7.18, N, 2.69. Found: C, 71.38, H, 7.37, N, 2.62

**Acknowledgement.**- The work was financially supported by a grant-in-aid from the Polish State Committee for Scientific Research (KBN). The support deserves our grateful thanks.

## REFERENCES

1. P. Lloyd-Williams, F. Albericio and E. Giralt, *Chemical Approach to the Synthesis of Peptides and Proteins*, CRC Press, Boca Raton – New York, 1997.
2. F. S. Gibson, S. C. Bergmeier and H. Rapoport, *J. Org. Chem.*, **59**, 3216 (1994).
3. Abbreviations used: Boc = *tert*-butoxycarbonyl, Trt = trityl, Cys = L-cysteine, *t*-Bu = *tert*-butyl.
4. Š. Horvat, B. Grgas, N. Raos and Vl. Simeon, *Int. J. Peptide Protein Res.*, **34**, 346 (1989).
5. A. Chimiak, T. Kolasa and J. F. Biernat, *Z. Chem.*, **12**, 264 (1972); *Chem. Abstr.*, **77**, 152541e (1972).
6. E. Ponnusamy, U. Fotadar, A. Spisni and D. Fiat, *Synthesis*, 48 (1986).

\*\*\*\*\*