

This article was downloaded by:

On: 27 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

MICROWAVE-ASSISTED SYNTHESIS OF TRIARYL PHOSPHATES

A. D. Sagar^a; N. A. Shinde^a; B. P. Bandgar^a

^a School of Chemical Sciences, S.R.T.M. University, Nanded-, INDIA

To cite this Article Sagar, A. D. , Shinde, N. A. and Bandgar, B. P.(2000) 'MICROWAVE-ASSISTED SYNTHESIS OF TRIARYL PHOSPHATES', *Organic Preparations and Procedures International*, 32: 3, 269 – 271

To link to this Article: DOI: 10.1080/00304940009355923

URL: <http://dx.doi.org/10.1080/00304940009355923>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

REFERENCES

1. Reviews on bone collagen are: a) N. B. Watts *Clin. Chem.*, **45**, 1359 (1999). b) L. Knott, A. J. Baily *Bone*, **22**, 181 (1998). c) I. T. James, A. J. Walne, D. Perrett, D. *Ann. Clin. Biochem.*, **33**, 397 (1996). d) D. R. Eyre "Principles of Bone Biology", p. 143, J. P. Bilezikian, L. G. Raisz, G. A. Rodan (Eds.), Academic Press, New York, NY, **1996**.
2. D. A. Hanson, D. R. Eyre, *J. Biol. Chem.*, **271**, 26508 (1996).
3. M. Adamczyk, D. D. Johnson, R. E. Reddy, R. E. *Tetrahedron*, **55**, 63 (1999).
4. a) R. Bambal, R. P. Hanzlik, *J. Org. Chem.*, **59**, 730 (1988). b) M. Adamczyk, D. D. Johnson, R. E. Reddy, R. E. *Bioconjugate Chem.*, **11**, 124 (2000).
5. T. W. Green, P. G. M. Wuts, "Protective Groups in Organic Synthesis", 3rd edition, John Wiley & Sons, Inc., New York, NY, **1999**.

MICROWAVE-ASSISTED SYNTHESIS
OF TRIARYL PHOSPHATES


Submitted by A. D. Sagar*, N. A. Shinde and B. P. Bandgar*
(08/30/99)

School of Chemical Sciences, S.R.T.M. University
Nanded-431 606, INDIA
Fax: 0091-2462-29245

Triaryl phosphates are of considerable interest because of their utility as plasticizers, flame retardants, lubricating oil additives and hydraulic fluids.¹ They are synthesized by reaction of phosphoryl chloride with phenols at elevated temperature (160-250°) in the presence of a Lewis acid catalyst,² at room temperature using a phase transfer catalyst³ or by using polymer supported phenoxides.⁴ Patented procedures are also available in the literature^{5,6} for the synthesis of triaryl phosphates. Some of the reported methods have limitations such as (i) drastic reaction conditions,² (ii) liberation of hydrogen chloride which may cause corrosion problems in industrial reactors, (iii) tedious work-up, and (iv) long reaction times. In certain cases, pyridine or dialkylanilines have been used to neutralize the liberated hydrogen chloride. High temperatures often lead to many by-products. Plasticizer grade triaryl phosphates must be of very high purity and the products obtained have to be distilled at low pressure for purification. Distillation at low pressures on an industrial scale raises serious problems.

Therefore, there is need of a rapid and environmentally-friendly general method for the synthesis of triaryl phosphates. In recent years, there has been considerable interest in microwave-induced reactions in organic synthesis.^{7,8} In view of the potential applications of triaryl phosphates and substantial reduction in reaction times under microwaves, we now report a novel method for the phosphorylation of phenols.

Triaryl phosphates have been synthesized in excellent yields by reacting phosphoryl chloride with a mixture of sodium hydroxide, few drops of water, and phenol under focused microwaves. Heating reaction mixture for 2-2.5 min followed by cooling was carried out at intervals of 30 sec. At room temperature, the reaction was very slow and incomplete after 5-6 hrs.



It is important to note that the reactions are carried out without solvent and there is no need of a catalyst as well as an inert atmosphere for this methodology. This method seems to be advanta-

Table. Synthesis of Triaryl Phosphates using Focused Microwaves

Product	Ar	Time (min.)	mp. (bp.) (°C)	lit. ^{1,9} mp (bp.)	Yield ^{a,b} (%)
3a	C ₆ H ₅	2.0	49-50	49-51	90
3b	<i>o</i> -MeC ₆ H ₄	2.5	90-91	90-91	88
3c	<i>m</i> -MeC ₆ H ₄	2.5	(257-262)	(258-263)	87
3d	<i>p</i> -MeC ₆ H ₄	2.5	77	77	88
3e	<i>o</i> -ClC ₆ H ₄	2.5	37	37	89
3f	<i>p</i> -ClC ₆ H ₄	2.5	116	117	89
3g	α-Naphthyl	2.5	148-149	149-150	86
3h	β-Naphthyl	2.0	111	111	88
3i	<i>p</i> -Anisyl	2.0	91	91	88
3j	3-Me-4-ClC ₆ H ₃	2.0	95	96	95

a) Yields are of pure isolated products. b) Products were characterized by IR, ¹H NMR, physical constants, and by comparison with authentic samples.

geous over conventional methods in terms of high yields, pure products, rapid reaction rates and an environmentally friendly procedure.

EXPERIMENTAL SECTION

All chemicals were of analytical grade, and solvents were freshly distilled before use. A microwave oven, Kelvinator T-37 model, was used for microwave irradiation with its 100% power. IR spectra were recorded on Bomem MB 104 FTIR spectrometer.

General Procedure for the Synthesis of Triaryl Phosphates.- A mixture of sodium hydroxide (3.1 mmol), few drops of water, the phenol (3.1 mmol) and phosphoryl chloride (1 mmol) in a 100 mL beaker was covered with watch glass and exposed to microwaves for 2 to 2.5 min. with heating and cooling at the interval of 30 sec. The reaction was monitored by TLC. After completion of the reaction, the product was extracted with chloroform (3 x 10 mL),¹⁰ and the chloroform layer was washed with 2% sodium hydroxide and water (2 x 25 mL each, respectively). Then the chloroform layer was separated from the aqueous layer, dried over anhydrous Na₂SO₄ and evaporated under reduced pressure to give the pure product (checked by TLC using 3 different solvent systems) in excellent yield.

Acknowledgement.- We thank UGC, New Delhi for financial assistance.

REFERENCES

1. E. Cherbuliez, "Organic Phosphorus Compounds" Vol. 6, G. M. Kosolapoff and L. Maier, Eds., Wiley-Interscience, New York p. 454 (1973).
2. W. L. Faith, D. B. Keyes and R. L. Clark, *Industrial Chemicals*, 4th Edn., Wiley- Interscience, New York, p. 849 (1975).
3. (a) V. K. Krishnakumar and M. M. Sharma, *Synthesis*, 558 (1983). (b) V. K. Krishnakumar, *Synth. Commun.*, **14**, 189 (1984).
4. A. D. Sagar, M. T. Thorat, M. M. Salunkhe, R. S. Selukar and P. P. Wadgaonkar, *ibid.*, **24**, 2029 (1994).
5. H. Seebek *et al.*, German Patent 283341 (1980), Bayer AG; *Chem. Abst.*, **93**, 26101 (1981).
6. U. Hucks *et al.*, German Patent 2833342 (1980), Bayer AG; *Chem. Abst.*, **93**, 7839 (1981).
7. R. S. Varma, *Green Chem.*, 43 (1999).
8. A. Loupy, A. Petit, J. Hamelin, F. T. Boulet, P. Jacquault and D. Mathe, *Synthesis*, 1213 (1998).
9. Z. Kucherov, *Zhur. Obshch. Khim.*, **19**, 126 (1949).
10. Ether, ethyl acetate or acetonitrile may be used instead of chloroform.