

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

AN IMPROVED SOLVENT-FREE PREPARATION OF 2-IMIDOYLPHENOLS

Cristina Cimarelli^a; Gianni Palmieri^a; Emanuela Volpini^a

^a Dipartimento di Scienze chimiche, Universita degli Studi di Comerino, Camerino, ITALY

To cite this Article Cimarelli, Cristina , Palmieri, Gianni and Volpini, Emanuela(2001) 'AN IMPROVED SOLVENT-FREE PREPARATION OF 2-IMIDOYLPHENOLS', *Organic Preparations and Procedures International*, 33: 4, 369 — 371

To link to this Article: DOI: 10.1080/00304940109356602

URL: <http://dx.doi.org/10.1080/00304940109356602>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.


The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

AN IMPROVED SOLVENT-FREE PREPARATION OF 2-IMIDOYLPHENOLS

Submitted by Cristina Cimarelli, Gianni Palmieri* and Emanuela Volpini
(10/30/00)

*Universita degli Studi di Camerino
Dipartimento di Scienze chimiche,
Via S. Agostino 1, I-62032 Camerino, ITALY*

2-Imidoylphenols (**3**) are heterofunctionally substituted organic compounds used in several applications and as precursors for some classes of molecules. They serve as ligands¹ in the preparation of rhodium complex catalysts, possess interesting antifungal properties,² and are starting materials for the synthesis of 4H-chromen-4-ylidenamines.³ Moreover, the stereoselective reduction of imidoyl phenols is a convenient synthetic pathway to aminophenols, another class of ligands extremely efficient in catalytic reactions such as the synthesis of optically active secondary alcohols by enantioselective addition of organozinc compounds to aldehydes.⁴



During the course of our studies on aminophenols and 4H-chromen-4-ylidenamines, we needed to prepare imidoylphenols (**3**) in a multigram scale as starting materials. These compounds are commonly prepared by condensation of primary amines with 2-acylphenols (**1**). The reaction protocol reported in the literature⁵ consists of dissolving the starting materials in benzene, toluene or xylene and refluxing the mixture for prolonged time (some hours) with azeotropic removal of water through a Dean-Stark apparatus. In some cases, Brønsted⁶ or Lewis acids (ZnCl₂)⁷ are used as catalysts. Extraction of the mixture affords the crude product in variable yields and these products normally required further purification, either by distillation or crystallization. The main drawbacks of this method are the use of toxic aromatic solvents, and the difficulty to obtain pure liquid products; purification by chromatographic separation is not convenient because of the partial decomposition of the imidoyl phenol on silica gel. We now report an improved route to the preparation of imidoyl phenols (**3**).

This new methodology produces the desired products without any need of heating, solvent, or acid catalyst. When 2-acylphenol (**1**, R¹ = Me, Et, Ph) is mixed with a slight excess of primary aliphatic amine [**2**, R² = Me, Prⁱ, Bn, (*R*)-CHCH₃,Ph] in a 1.1 molar ratio, an immediate yellow coloration of the mixture and evolution of heat are observed. After a few seconds the water produced separates as droplets. The reaction mixture became more viscous during the course of the reaction until an oil or a solid is obtained. This final mixture contains the product in a 91-98% yield and purification is carried out by washing the reaction mixture with water to remove the excess amine.

The speed and efficacy of the reaction is probably due to carbonyl activation by the acidic

phenolic proton; acetophenone does not react with amines under the same conditions. Moreover, the presence of the hydroxy function in the *ortho* position helps stabilize the final imidoxyphenol. For example, when 4-hydroxyacetophenone and isopropylamine are mixed, a bright yellow color develops immediately, but the final product is isolated in poor yield and rapidly loses isopropylamine to give back 4-hydroxyacetophenone. This method is especially suitable for low boiling amines, which can be used as hydrochlorides. In this case, a larger excess of amine is used to circumvent the volatility problem, and the reaction takes place only when 30% aqueous sodium hydroxide is added dropwise at 0° to the mixture of amine hydrochloride and 2-acylphenols. Aniline does not react, probably because it is less nucleophilic than the aliphatic amines.

Table 1. Synthesis of 2-Imidoxyphenols (**3**)

Product	R ¹	R ²	Time (h)	Yield (%)	mp [bp] (°C)	Lit. (°C)
3a	Me	Me	4.00 ^a	98	65-66	68 ^{5a}
3b	Me	Pr ⁱ	0.75	98	[120 (3 mm)]	[123 (3 mm)] ^{5a}
3c	Me	Bn	2.00	97	118-119	118-119 ⁸
3d	Et	Me	5.00 ^a	93	[118 (3 mm)]	- ⁸
3e	Et	Pr ⁱ	1.00	96	[132 (3 mm)]	- ⁸
3f	Me	(<i>R</i>)-CHCH ₃ Ph	2.00	96 ^b	[179 (3 mm)]	- ⁸
3g	Et	(<i>R</i>)-CHCH ₃ Ph	5.00	94	78-80	79-81 ⁸
3h	Ph	(<i>R</i>)-CHCH ₃ Ph	8.00	91	129-131	128-130 ^{4a}

a) Methylamine hydrochloride was used; b) The reaction performed starting from 20 g of acetophenone gave a 94% yield.

In summary, this preparation does not involve any complex procedure, use of toxic aromatic solvents is avoided, as is the need for acid catalysis and heating. The brief work-up procedure affords the products directly in high purity and the reaction can be easily scaled up. The very mild conditions makes this procedure especially suitable for use with low boiling amines, more practically as their hydrochlorides.

EXPERIMENTAL SECTION

¹H and ¹³C-NMR spectra were recorded with a Varian VXR 300 instrument. IR spectra were obtained with a Perkin-Elmer 257 spectrometer. GC-MS analyses were performed with an HP 59970 workstation formed by an HP-5890 gas chromatograph equipped with a methyl silicone capillary column and by an HP-5970 mass detector. All melting points are uncorrected. All reagents and solvents were distilled prior to use or were of commercial quality from freshly opened containers.

Procedure for High-boiling Amines.- To 5 mmol of 2-acylphenols **1** (R¹ = Me, Et, Ph) in a 25 mL round bottom flask, equipped with a stir bar at 0° (ice bath) under inert atmosphere, was added dropwise 5.5 mmol of the amine **2** (R² = Prⁱ, Bn, (*R*)-CHCH₃Ph) (20 min). The reaction mixture immedi-

ately turned to bright yellow and heat was evolved. The reaction mixture was stirred at ambient temperature and monitored by GC or NMR methods until the starting 2-acylphenol disappeared. It was then diluted with *n*-hexane (50 mL) and washed with water (50 mL). The organic layer was dried over Na₂SO₄ and evaporated under reduced pressure. The resulting oil or solid was the pure 2-imidoylphenol (**3**) in 91-98% yields, whose structure was confirmed on the basis of spectroscopic data according to literature.⁸ The purity of imidoylphenol (**3**) (95-98 %) was determined by HPLC (*n*-hexane - ethyl acetate, 95:5).

The reactions of 4-hydroxyacetophenone and acetophenone with isopropylamine were performed in the same ratios and under the same conditions.

Procedure for Low-boiling Amines.- To a mixture of 5 mmol of 2-acylphenols **1** (R¹ = Me, Et) and 15 mmol of the methylamine hydrochloride (R² = Me) in a round bottom flask equipped with a stir bar at 0° (ice bath) under inert atmosphere, was added dropwise 2.5 mL of 30% wt. NaOH aqueous solution. The reaction mixture became bright yellow and developed heat with the first drop of base. The reaction mixture, stirred at 0°, was monitored by GC or NMR methods until the starting 2-acylphenol disappeared. Work-up was performed by washing the reaction mixture diluted in *n*-hexane (50 mL) with water (50 mL). The organic layer was then dried with Na₂SO₄ and evaporated under reduced pressure. The resulting oil or solid was the pure 2-imidoylphenol **3** in 93-98% yields, whose structure was assigned on the basis of spectroscopic data according to literature.⁸

Acknowledgement.- Financial support from Ministero dell'Università e della Ricerca Scientifica e Tecnologica, Rome, and the University of Camerino (National Project "Stereoselezione in Sintesi Organica. Metodologie ed applicazioni") is gratefully acknowledged

REFERENCES

1. W. R. Cullen and E. B. Wickenheiser, *J. Organomet. Chem.*, **370**, 141 (1989).
2. M. R. Manrao and S. Kohli, *J. Indian Chem. Soc.*, 348 (1986).
3. C. Cimarelli and G. Palmieri, *Tetrahedron*, **56**, 475 (2000).
4. a) G. Palmieri, *Eur. J. Org. Chem.*, 805 (1999); b) G. Palmieri, *Tetrahedron: Asymmetry*, **11**, 3361 (2000).
5. a) P. Maroni, L. Cazaux, P. Tisnes and M. Zambeti, *Bull. Soc. Chim. France*, 179 (1980); b) S. J. Joglekar and S. D. Samant, *J. Indian Chem. Soc.*, **65**, 110 (1988).
6. a) A. F. Al-Sayyab, A. Lawson and J. O Stevens, *J. Chem. Soc. C*, 411 (1968); b) K. Hiroi, S. Sato and R. Kitayama, *Chem. Pharm. Bull.*, **31**, 3471 (1983).
7. S. Mal, J. L. Norula and M. S. M Rawat, *Indian J. Chem. Sect. B*, **21**, 120 (1982).
8. C. Cimarelli and G. Palmieri, *Tetrahedron*, **54**, 15711 (1998).