

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

A PRACTICAL LARGE-SCALE PREPARATION OF 5'-BROMOSPIRO(CYCLOHEXANE-1,3-[3H]INDOL)-2'(1'H)-ONE

Bogdan K. Wilk^a; Arkadiy Rubezhov^a; Jean L. Helorn^a; Lisa R. Routel^a; John R. Potoski^a

^a Chemical Development, Wyeth Research, Pearl River, NY

To cite this Article Wilk, Bogdan K. , Rubezhov, Arkadiy , Helorn, Jean L. , Routel, Lisa R. and Potoski, John R.(2005) 'A PRACTICAL LARGE-SCALE PREPARATION OF 5'-BROMOSPIRO(CYCLOHEXANE-1,3-[3H]INDOL)-2'(1'H)-ONE', *Organic Preparations and Procedures International*, 37: 3, 283 – 285

To link to this Article: DOI: 10.1080/00304940509354961

URL: <http://dx.doi.org/10.1080/00304940509354961>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


13. (a) A. R. Hajipour, M. Arbabian and A. E. Ruoho, *J. Org. Chem.*, **67**, 8622 (2002). (b) A. R. Hajipour and A. E. Ruoho, *Org. Prep. Proced. Int.*, **34**, 647 (2002). (c) A. R. Hajipour and H. Adibi, *J. Chem. Research (S)*, 294 (2004).
14. I. Jernej, S. Stojan and Z. Marko, *Synthesis*, 1869 (2004).

**A PRACTICAL LARGE-SCALE PREPARATION OF
5'-BROMOSPIRO(CYCLOHEXANE-1,3'-[3H]INDOL)-2'(1'H)-ONE**

Submitted by Bogdan K. Wilk*, Arkadiy Rubezhov, Jean L. Helom, Lisa R. Routel,
(03/30/05) and John R. Potoski

Wyeth Research, Chemical Development
401 N. Middletown Road, Pearl River, NY 10965
e-mail: wilkb@wyeth.com

We were interested in a large-scale preparation of 5'-bromospiro(cyclohexane-1,3'-[3H]indol)-2'(1'H)-one (**3**), an intermediate in the synthesis of progesterone receptor modulators.¹ Initially, the compound was prepared from oxindole using the Kende procedure^{2a} by alkylation followed by bromination with bromine in acetic acid/chloroform.³ Unfortunately, this sequence gave multiple impurities in the alkylation product.^{1a,4} On large scale, this was a serious problem and we had to resort to chromatography and recrystallization.⁵ The subsequent bromination had to be carefully controlled to avoid over-bromination. In addition, brominated impurities were formed from the side-products generated in the alkylation step.



The regioselectivity and purity problems were overcome by simply changing the order of transformations, *i. e.* introduction of bromine in the oxindole core followed by alkylation of 5-bromoisatin (**2**),^{3,6} which was prepared from inexpensive, technical grade 5-bromoisatin (**1**).⁷ It was found that *n*-butyllithium could be replaced with potassium *tert*-butoxide and 1,5-diiodopentane with the more stable 1,5-dibromopentane at $0^\circ C$ in tetrahydrofuran (THF) in the alkylation step.⁸ In comparison to the initial sequence, the reaction was clean and the product crystallized out upon replacement of THF with acetonitrile.⁹

The new procedure is suitable for scale-up. In fact, six consecutive scale-up batches gave a total of 45 kg of 97.0-99.5% pure product. In addition, the alkylation proceeded with reproducible and consistent yields of 84-89%.

EXPERIMENTAL SECTION

Melting points were determined on a Mel-Temp II apparatus and are uncorrected. HPLC was performed using Phenomenex Luna C8(2), 3 μ m, 150 x 4.60 mm column and acetonitrile - water mobile phase containing ammonium acetate modifier. The reagents were purchased from Aldrich and used without further purification; 5-bromoisatin (**1**) was purchased from Lancaster Synthesis, Inc.

5-Bromo-1,3-dihydroindol-2-one (2).- A 5-L, four-necked flask equipped with a nitrogen inlet, an addition funnel, an overhead mechanical stirrer, temperature controller, condenser, and a heating mantle was charged with 5-bromoisatin (**1**, 300 g, 1.30 mol, 98% pure) and ethylene glycol (2.0 L). Hydrazine hydrate, $N_2H_4 \cdot 1.5 H_2O$ (150 mL) was added from the addition funnel at the rate of 13 mL/min. The orange suspension turned gradually to a brown solution. The mixture was heated to 120°C and stirred for 6 h. The suspension which formed, was cooled to ambient temperature and the solid was collected. The cake was washed with water (4 x 200 mL) and dried in a vacuum oven at 40°C/20 mmHg to give **2** (197 g, 70% yield; 97.0% purity, mp 218-222°C, *lit.*^{3b} mp 220-221°C) as a tan solid. If needed, the product may be recrystallized from ethanol.^{3b}

5'-Bromospiro(cyclohexane-1,3'-indolin)-2'-one (3).- A 5-L, four-necked flask was equipped with mechanical stirrer, condenser with nitrogen adapter, addition funnel, thermometer and cooling bath. Tetrahydrofuran (750 mL) was added to the flask followed by 5-bromo-1,3-dihydroindol-2-one (**2**, 150 g, 0.708 mol) and the mixture was cooled to -15°C. Potassium *tert*-butoxide (251 g, 2.12 mol, 3.0 equiv.) was dissolved in THF (1.5 L) and added dropwise within 30 min., while the temperature was maintained at -15°C. The contents of the addition funnel were rinsed into the flask with THF (250 mL). 1,5-Dibromopentane (119 mL, 199 g, 0.866 mol, 1.2 equiv.) was added dropwise with the temperature being maintained between -10 and -2°C. The contents of the addition funnel were rinsed with THF (100 mL) and the mixture was stirred at 0°C for 1 h. The reaction mixture was quenched with 3.5% hydrochloric acid (1.0 L). After 1 h of stirring, the layers were separated. The dark red, organic layer was washed with brine (250 mL) and transferred to a 5-L distillation flask. THF was distilled off and replaced with acetonitrile. During distillation, acetonitrile (1.0 L) was added gradually until the batch temperature reached 77-80°C and an orange slurry was formed. The heating mantle was replaced with a cooling pan and the slurry was gradually cooled to 0-5°C. The reaction mixture was stirred for an additional 30 min. at 0-5°C. The solid was collected, washed with cold (0-5°C) acetonitrile (3 x 150 mL) and dried in a vacuum oven at 50°C/20 mmHg to give **3** (158 g, 80% yield; 99.5% purity, mp 196-198°C, *lit.*^{1b} mp 196-199°C) as an off-white solid.

Acknowledgments.- We thank J. Brazzillo, D. Boop, S. Devarakonda, M. Hofman, C. Rothrock and R. Zhao for their valuable input during scale-up and the Analytical and Quality Sciences group for their support. We also acknowledge M. Salata, A. Sharma and A. Fensome for their help.

REFERENCES

1. a) A. Fensome, R. H. W. Bender, J. Cohen, M. A. Collins, V. Mackner, L. L. Miller, J. W. Ullrich, R. Winneker, J. E. Wrobel, P. Zhang, Z. Zhang and Y. Zhu, *Bioorg. Med. Chem. Lett.*, **12**, 3487 (2002). M. A. Collins, V. Hudak, R. Bender, A. Fensome, P. Zhang, L. Miller, R. C. Winneker, Z. Zhang, Y. Zhu, J. Cohen, R. J. Unwalla and J. Wrobel, *Bioorg. Med. Chem. Lett.*, **14**, 2185 (2004); b) G. S. Grubb, L. Zhi, T. K. Jones, C. M. Tegley, A. Fensome, L. L. Miller, J. W. Ullrich, R. H. W. Bender, P. Zhang, J. E. Wrobel, J. P. Edwards, PCT Publication No. WO 00/66167; *Chem. Abstr.*, **133**, 350135 (2000).
2. (a) Of the several routes available for the synthesis of spiro(cyclohexane-1,3'-indolin)-2'-one core, Kende method is the shortest. A. S. Kende and J. C. Hodges, *Synth. Commun.*, **12**, 1 (1982). (b) Other methods are less suitable for scale-up. R. F. Moore and S. G. P. Plant, *J. Chem. Soc.*, 3475 (1951). J. Wolff and M. Taddei, *Tetrahedron*, **42**, 4267 (1986). I. Fleming, M. A. Loreto and I. H. M. Wallace, *J. Chem. Soc., Perkin Trans. 1*, 349 (1986). S. Lee and J. F. Hartwig, *J. Org. Chem.*, **66**, 3402 (2001).
3. a) H. R. Henze and C. M. Blair, *J. Am. Chem. Soc.*, **55**, 4621 (1933). b) W. C. Sumpter, M. Miller and L. N. Hendrick, *J. Am. Chem. Soc.*, **67**, 1656 (1945).
4. I. Gruda, *Can. J. Chem.*, **50**, 18 (1972).
5. TMEDA, a complexing agent used in the Kende procedure, was later replaced with LiCl which simplified the work-up. Also, 1,5-dibromopentane, 1,5-ditosylpentane, and a 1:1 mixture of 1,5-dibromo- and 1,5-diiodopentane were tried with no apparent improvement in the yield and the purification procedure.
6. L. Sun, N. Tran, F. Tang, H. App, P. Hirth, G. McMahon and C. Tang, *J. Med. Chem.*, **41**, 2588 (1998). We found this method impractical for a large-scale preparation.
7. C. Crestini and R. Saladino, *Synth. Commun.*, **24**, 2835 (1994).
8. A. Di Malta, L. Foulon, G. Garcia, D. Nisato, R. Roux, C. Serradeil-Legal, G. Valette and J. Wagnon, EP0636608 (1995); *Chem. Abstr.*, **122**, 213926 (1995).
9. Alkylation of oxindole under conditions described in Ref. 8 required extensive chromatographic purification.
