

This article was downloaded by:

On: 26 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t902189982>

A SYNTHESIS OF SELECTED *o,o'*-DISUBSTITUTED DIARYLACETIC ESTERS AND DIARYLMETHANES

Richard A. Bunce^a; James E. Schammerhorn^a

^a Department of Chemistry, Oklahoma State University, Stillwater, Oklahoma

To cite this Article Bunce, Richard A. and Schammerhorn, James E.(2005) 'A SYNTHESIS OF SELECTED *o,o'*-DISUBSTITUTED DIARYLACETIC ESTERS AND DIARYLMETHANES', *Organic Preparations and Procedures International*, 37: 6, 550 – 555

To link to this Article: DOI: 10.1080/00304940509354984

URL: <http://dx.doi.org/10.1080/00304940509354984>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

3. T. Honda, F. G. Favaloro, Jr., T. Janosik, Y. Honda, N. Suh, M. B. Sporn, and G. W. Gribble, *Org. Biomol. Chem.*, **1**, 4384 (2003).
4. J. M. Cassady and M. Suffness, "Anticancer Agents Based on Natural Product Models", p 254, Academic Press, New York, NY, 1980.
5. D. L. Snitman and D. S. Watt, *Synth. Commun.*, **8**, 187 (1978).
6. S. M. Kerwin, A. G. Paul, and C. H. Heathcock, *J. Org. Chem.* **52**, 1686 (1987).
7. D. Caine, *Org. React.*, **23**, 1 (1976) and references cited therein.
8. Overlapped signals which cannot be assigned.


A SYNTHESIS OF SELECTED *o,o'*-DISUBSTITUTED DIARYLACETIC ESTERS AND DIARYLMETHANES

Submitted by: Richard A. Bunce* and James E. Schammerhorn[†]
(7/26/05)

*Department of Chemistry, Oklahoma State University
Stillwater, Oklahoma 74078-3071
E-mail: rab@okstate.edu*

A recent project required access to 2-(2-nitrobenzyl)benzoic acid (**4a**) for a cyclization study to prepare 5,11-dihydro-6*H*-dibenz[*b,e*]azepin-6-one.¹ This compound has been previously synthesized by nitration of 2-benzylbenzoic acid,² but it was not purified from the accompanying C-4 nitration product. We, therefore, sought a method to prepare the C-2 nitrated compound in pure form. Our synthetic plan involved (1) nucleophilic aromatic substitution of the 2-[2-(methoxycarbonyl)phenyl]acetate anion (**2a**) to 2-fluoro-1-nitrobenzene, (2) basic hydrolysis of the two esters, and (3) selective decarboxylation of the doubly benzylic acid group.³ Previous work in this laboratory⁴ and by others⁵ has shown that stabilized anions can be added to 2- and 4-halo-1-nitrobenzenes by nucleophilic aromatic substitution but, to date, additions of methyl phenylacetate derivatives bearing anion-stabilizing electron withdrawing groups at C-2 of the aromatic ring have not been described. We report here our results on the application of this strategy to the synthesis of several *o,o'*-disubstituted diarylacetic esters and diarylmethanes.

The synthesis of the target compounds is illustrated in the *Scheme*. The methyl phenylacetate derivatives **1a-d** were prepared by known methods.⁶⁻⁹ The corresponding anions **2a-d** were generated using potassium *tert*-butoxide as the base; sodium hydride proved less satisfactory. As in our earlier work,⁴ DMPU (1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone) was found to be a convenient and relatively safe solvent for these reactions. Following anion formation, 0.75 equivalents of 2-fluoro-1-nitrobenzene was added dropwise and the mixture was stirred at 23°C for 1 h and at 60°C for 12-24 h. Aqueous ammonium chloride workup gave the substitution products in 60-72% yield.¹⁰ With the exception of **3c**, all of the products were solids and easily purified by trituration with 5% petroleum ether in ether followed by filtration. Product **3c** was purified by flash chromatography to give the addition product along with *ca* 20% of recovered **1c**.



a) X = CO₂CH₃ except in **4** where X = CO₂H; b) X = CN; c) X = PhCO; d) X = NO₂

The hydrolysis-decarboxylation procedure proceeded smoothly for substrates **3a-c** to give **4a-c** in 67-92% yield. Substrate **3d**, however, underwent decomposition when treated with aqueous base in the hydrolysis step. The final product was a complex mixture containing at best only a trace of **4d**. Attempts to demethoxycarbonylate **3d** by an S_N2-type dealkylation¹¹ using lithium chloride in polar aprotic solvents (DMSO and DMPU) also failed and further work on this substrate was discontinued.

In summary, a procedure has been optimized for the nucleophilic aromatic substitution of several *o*-substituted phenylacetic ester anions to 2-fluoro-1-nitrobenzene. The synthesis is efficient, clean and generates products that are easily purified. The method is limited in not permitting the use of esters bearing enolizable groups at C-2, but should find use for the preparation of a modest number of *o,o'*-disubstituted diarylacetic esters and diarylmethanes.

EXPERIMENTAL SECTION

DMPU (1,3-dimethyl-3,4,5,6-tetrahydro-2(1*H*)-pyrimidinone, Aldrich, puriss grade, absolute, over molecular sieves) was stored under nitrogen and transferred by syringe into reactions where it was used. DMF (EM Science, GR grade), from a freshly opened bottle, was dried over 4Å molecular sieves under nitrogen and transferred by syringe into reactions where it was used. Commercial anhydrous potassium carbonate was ground to a fine powder, dried under vacuum at 120°C for 24 h and stored in an oven at 120°C. Methyl 2-[2-(methoxycarbonyl)phenyl]acetate

(**1a**),⁶ methyl (2-cyanophenyl)acetate (**1b**),⁷ methyl (2-benzoylphenyl)acetate (**1c**)⁸ and methyl (2-nitrophenyl)acetate (**1d**)⁹ were prepared by literature methods. All reactions were performed under dry nitrogen in oven-dried glassware. Reactions were monitored by thin layer chromatography on silica gel GF plates. Preparative separations were carried out using flash column chromatography¹² on silica gel (grade 62, 60-200 mesh) mixed with UV active phosphor (Sorbent Technologies UV-5); band elution was monitored using a hand-held UV lamp. Melting points were uncorrected. IR spectra were run as thin films on NaCl disks and were referenced to polystyrene. ¹H and ¹³C NMR spectra were measured in CDCl₃ at 300 MHz and 75 MHz, respectively, and referenced to internal TMS; coupling constants (*J*) are given in Hz. Mass spectra (EI/DP) were obtained at 70 eV.

Representative Procedure for Nucleophilic Aromatic Substitution of *o*-Substituted Methyl Phenylacetates to 2-Fluoro-1-nitrobenzene: Methyl 2-[2-(Methoxycarbonyl)phenyl]-2-(2-nitrophenyl)acetate (3a).

A 50-mL three-necked round-bottomed flask equipped with an addition funnel, a condenser and a magnetic stir bar was charged with 8 mL of DMPU and 0.57 g (5.10 mmol) of potassium *tert*-butoxide. The mixture was stirred to dissolve the potassium *tert*-butoxide and a solution of 1.04 g (5.00 mmol) of **1a** in 2 mL of DMPU was added dropwise during 5 min. The reaction became warm as the bright orange anion was formed. The mixture was stirred for 10 min and 0.53 g (3.75 mmol) of 2-fluoro-1-nitrobenzene was added. The reaction again became warm during the addition. The reaction was stirred at 23°C for 1 h and at 60°C (oil bath) for 12-24 h, then added to 10% aqueous NH₄Cl and extracted with ether (3x). The combined ether extracts were washed with water (2x) and 5% aqueous NaCl (1x), dried (MgSO₄), and concentrated under vacuum to give a yellow oil that crystallized on standing. The product was purified by trituration with 5% petroleum ether in ether to give 0.89 g (72%) of **3a** as light yellow crystals, mp 115-117°C.

IR 1738, 1716, 1527, 1351 cm⁻¹; ¹H NMR δ 8.05 (overlapping dd, 2 H, *J* = 7.9, 1.6), 7.54 (td, 1 H, *J* = 7.6, 1.4), 7.50 (td, 1 H, *J* = 7.6, 1.6), 7.45 (td, 1H, *J* = 7.6, 1.6), 7.40 (dd, 1 H, *J* = 7.6, 1.1), 7.23 (dd, 1 H, *J* = 7.4, 1.1), 7.05 (dd, 1 H, *J* = 7.6, 1.6), 6.56 (s, 1 H), 3.78 (s, 3 H), 3.77 (s, 3 H); ¹³C NMR δ 172.4, 166.8, 148.0, 138.2, 133.8, 133.0, 132.5, 131.7, 131.0, 129.8, 129.5, 128.1, 127.9, 124.9, 52.6, 52.2, 50.7; MS *m/z* 329 (M⁺).

Anal. Calcd for C₁₇H₁₅NO₆: C, 62.01; H, 4.56; N, 4.26. Found: C, 61.89; H, 4.52; N, 4.31

Methyl 2-(2-Cyanophenyl)-2-(2-nitrophenyl)acetate (3b): 0.75 g (68%) as tan crystals purified by trituration with 5% petroleum ether in ether, mp 109-110°C; IR 2225, 1738, 1527, 1349 cm⁻¹; ¹H NMR δ 8.13 (dd, 1 H, *J* = 7.6, 1.6), 7.73 (dm, 1 H, *J* = 7.6), 7.64 (td, 1 H, *J* = 7.6, 1.1), 7.57 (td, 1 H, *J* = 7.6, 1.6), 7.54-7.44 (complex, 2 H), 7.32 (d, 1 H, *J* = 7.9), 7.03 (dd, 1 H, *J* = 7.9, 1.6), 6.11 (s, 1 H), 3.80 (s, 3 H); ¹³C NMR δ 170.7, 148.2, 140.2, 133.6, 133.5, 133.1, 131.8, 131.1, 129.3, 129.0, 128.5, 125.6, 116.9, 113.9, 53.0, 51.9; MS *m/z* 296 (M⁺).

Anal. Calcd for C₁₆H₁₂N₂O₄: C, 64.86; H, 4.05; N, 9.46. Found: C, 64.89; H, 4.09; N, 9.32

Methyl 2-(2-Benzoylphenyl)-2-(2-nitrophenyl)acetate (3c): 0.84 g (60%) as a light yellow oil, purified by flash column chromatography using increasing concentrations of ether in hexane

(0.20 g of the starting keto ester was recovered); IR 1738, 1657, 1527, 1349 cm^{-1} ; ^1H NMR δ 7.99 (dd, 1 H, $J = 8.2, 1.4$), 7.73 (dd, 2 H, $J = 8.2, 1.4$), 7.56-7.48 (complex, 3 H), 7.46- 7.37 (complex, 5 H), 7.28 (dd, 1 H, $J = 7.9, 0.5$), 7.22 (dd, 1 H, $J = 7.9, 1.4$), 6.04 (s, 1 H), 3.73 (s, 3 H); ^{13}C NMR δ 197.5, 171.9, 148.6, 138.4, 137.3, 136.5, 133.6, 133.1 (2), 131.5, 131.0, 130.3, 130.2, 129.8, 128.3 (2), 127.0, 125.2, 52.8, 50.5; MS m/z 375 (M^+).

Anal. Calcd for $\text{C}_{22}\text{H}_{17}\text{NO}_5$: C, 70.40; H, 4.53; N, 3.73. Found: C, 70.17; H, 4.49; N, 3.81

Methyl 2,2-bis(2-Nitrophenyl)acetate (3d): 0.78 g (66%) as tan crystals purified by trituration with 5% petroleum ether in ether, mp 101-102°C; IR 1742, 1528, 1349 cm^{-1} ; ^1H NMR δ 8.11 (dd, 2 H, $J = 8.2, 1.4$), 7.61 (td, 2 H, $J = 7.6, 1.6$), 7.54 (td, 2 H, $J = 7.9, 1.6$), 7.19 (dd, 2 H, $J = 7.6, 1.4$), 6.34 (s, 1 H), 3.77 (s, 3 H); ^{13}C NMR δ 171.0, 148.8, 133.5, 132.0, 130.9, 128.9, 125.6, 52.9, 50.1; MS m/z 316 (M^+).

Anal. Calcd for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{O}_6$: C, 59.96; H, 3.80; N, 8.86. Found: C, 59.69; H, 3.85; N, 8.98

Representative Procedure for Hydrolysis-Decarboxylation: 2-[(2-Nitrophenyl)methyl]benzoic Acid (4a).- The procedure of Bull and co-workers was modified.³ Step 1: A solution of 1.15 g (3.50 mmol) of **3a** in 15 mL of dioxane was treated with 4.2 mL of aqueous 2.0 M NaOH (8.40 mmol) and stirred for 6 h at room temperature. The dioxane was removed under vacuum, the aqueous product was acidified to pH 2 with 1.0 M HCl and extracted with ether (3x). The combined ether extracts were washed with water (2x) and 5% aqueous NaCl (1x), dried (MgSO_4), and concentrated under vacuum. The crude product contained a mixture of the dicarboxylic acid and the monodecarboxylated product, which was carried on without further purification.

Step 2: The above mixture was dissolved in 15 mL of anhydrous DMF, 1.00 g (7.25 mmol) of anhydrous K_2CO_3 was added and the mixture was stirred at 50°C (oil bath) for 1-2 h. The reaction mixture was cautiously added to 1.0 M HCl and extracted with ether (3x). The combined ether extracts were washed with water (2x) and 5% aqueous NaCl (1x), dried (MgSO_4), and concentrated under vacuum. The crude product was purified by trituration from 5% petroleum ether in ether to give 0.60 g (67%) of **4a** as light yellow crystals. An analytical sample was obtained by recrystallization from benzene, mp 143-144°C.

IR 3515-2109, 1688, 1524, 1348 cm^{-1} ; ^1H NMR δ 11.7 (br s, 1 H), 8.13 (dd, 1 H, $J = 7.9, 1.4$), 7.94 (dd, 1 H, $J = 7.9, 1.3$), 7.50 (td, 1 H, $J = 7.4, 1.6$), 7.44 (td, 1 H, $J = 7.6, 1.4$), 7.35 (m, 2 H), 7.13 (dd, 1 H, $J = 7.6, 0.5$), 7.04 (dd, 1 H, $J = 7.9, 1.1$), 4.71 (s, 2 H); ^{13}C NMR δ 172.4, 149.6, 141.5, 135.8, 133.4, 132.8, 132.2, 131.6, 131.5, 128.2, 127.1, 127.0, 124.5, 37.0; MS m/z 257 (M^+).

Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{NO}_4$: C, 65.37; H, 4.28; N, 5.45. Found: C, 65.42; H, 4.31; N, 5.39

2-[(2-Nitrophenyl)methyl]benzonitrile (4b): run on a 1.70 mmol scale; step 1 used 2.0 mL of 2.0 M NaOH (4.00 mmol) in 8 mL of dioxane; step 2 used 0.23 g (1.70 mmol) of K_2CO_3 in 8 mL of DMF; 0.37 g (92%) of **4b** was isolated as a light yellow powder purified by trituration with 2% ether in petroleum ether, mp 107-108°C; IR 2218, 1521, 1344 cm^{-1} ; ^1H NMR δ 8.03 (dd, 1 H, $J = 8.2, 1.4$), 7.68 (dd, 1 H, $J = 7.6, 1.4$), 7.58 (td, 1 H, $J = 7.6, 1.4$), 7.47 (m, 2 H), 7.34 (tm, 1 H, $J = 7.6$), 7.28 (ddd, 1 H, $J = 7.6, 1.4, 0.5$), 7.09 (dm, 1 H, $J = 7.9$), 4.58 (s, 2 H); ^{13}C NMR δ

149.2, 142.5, 133.4, 133.1 (2), 133.0, 132.5, 129.5, 128.2, 127.3, 125.2, 117.7, 113.0. 37.0; MS m/z 238 (M^+).

Anal. Calcd for $C_{14}H_{10}N_2O_2$: C, 70.59; H, 4.20; N, 11.76. Found: C, 70.46; H, 4.14; N, 11.91

2-[(2-Nitrophenyl)methyl]benzophenone (4c): run on a 1.88 mmol scale; step 1 used 2.2 mL of 2.0 M NaOH (4.40 mmol) in 8 mL of dioxane; step 2 used 0.26 g (1.88 mmol) of K_2CO_3 in 8 mL of DMF; 0.54 g (91%) of **4c** was isolated as a tan oil purified by flash column chromatography; IR 1660, 1524, 1349 cm^{-1} ; 1H NMR δ 7.87 (dd, 1 H, $J = 8.2, 1.4$), 7.74 (dm, 2 H, $J = 8.4$), 7.56 (tt, 1 H, $J = 7.4, 1.4$), 7.46-7.40 (complex, 4 H), 7.38-7.26 (complex, 3 H), 7.23 (dm, 1 H, $J = 8.7$), 7.13 (dm, 1 H, $J = 7.9$), 4.42 (s, 2 H); ^{13}C NMR δ 198.2, 149.5, 138.6, 138.1, 137.4, 135.3, 133.2, 133.0, 132.9, 130.6, 130.4, 130.2, 129.1, 128.3, 127.4, 126.0, 124.7, 36.2; MS m/z 317 (M^+).

Anal. Calcd for $C_{20}H_{15}NO_3$: C, 75.71; H, 4.73; N, 4.42. Found: C, 75.50; H, 4.68; N, 4.29

1-Nitro-2-[(2-nitrophenyl)methyl]benzene (4d): Compound **3d** decomposed under the basic hydrolysis conditions to give a complex mixture of products. The trace amount of **4d** produced could not be isolated.

Acknowledgment.- The authors wish to thank the Oklahoma Center for the Advancement of Science and Technology (OCAST, HR01-015) for support of this research. J. E. S. also thanks the Oklahoma State University College of Arts and Sciences for a Niblack Research Scholarship and the Department of Chemistry for a Skinner Scholarship. Funding for the 300 and 400 MHz NMR spectrometers of the Oklahoma Statewide Shared NMR Facility was provided by NSF (BIR-9512269), the Oklahoma State Regents for Higher Education, the W. M. Keck Foundation, and Conoco, Inc. Funding for our mass spectrometer was obtained from NSF and OCAST.

REFERENCES AND NOTES

† Undergraduate research participant, 2004-present.

1. R. A. Bunce and J. E. Schammerhorn, *J. Heterocycl. Chem.*, Submitted.
2. A. J. G. Baxter, J. Dixon, F. Ince, C. N. Manners and S. J. Teague, *J. Med. Chem.*, **36**, 2739 (1993).
3. D. J. Bull, M. J. Fray, M. C. Mackenny and K. A. Malloy, *Synlett*, 647 (1996).
4. R. A. Bunce, M. H. Randall and K. G. Applegate, *Org. Prep. Proced. Int.*, **34**, 493 (2002).
5. (a) R. L. Augustine, A. J. Gustavsen, S. F. Wanat, I. C. Pattison, K. S. Houghton and G. Koletar, *J. Org. Chem.*, **38**, 3004 (1973). (b) M. Perchinunno, A. Guerrato and F. Pregno-lato, *Org. Prep. Proced. Int.*, **9**, 303 (1977).
6. This diester was prepared by dissolving homophthalic acid in methanol containing a catalytic amount of concentrated sulfuric acid and refluxing overnight with removal of water using 3Å molecular sieves according to the procedure given in H. R. Harrison, W. M. Haynes, P. Arthur and E. J. Eisenbraun, *Chem. Ind. (London)*, 1568 (1968).

7. R. A. Bunce and L. B. Johnson, *Org. Prep. Proced. Int.*, **31**, 407 (1999).
8. (2-Benzoylphenyl)acetic acid was prepared according to T. de Paulis, C. R. Ross Betts, H. E. Smith, P. L. Mobley, D. H. Manier and F. Sulser, *J. Med. Chem.*, **24**, 1021 (1981) and esterified using the general method given in ref 6.
9. R. A. Bunce, D. M. Herron, L. B. Johnson and S. V. Kotturi, *J. Org. Chem.*, **66**, 2822 (2001).
10. An attempt was also made to generate products resulting from nucleophilic aromatic substitution of **1a** and **1b** with 4-fluoro-1-nitrobenzene. While the initial addition was successful in each case, the yields were 10-25% lower than reactions with 2-fluoro-1-nitrobenzene. Additionally, the hydrolysis-decarboxylation procedure gave products that were more difficult to purify. Thus, we did not investigate additional substitution patterns.
11. J. E. McMurry, *Org. React.*, **24**, 187 (1976).
12. W. C. Still, M. Kahn and A. Mitra, *J. Org. Chem.*, **43**, 2923 (1978).

A CONVENIENT SYNTHESIS OF 3-ETHOXYCARBONYLAMINO- 2-HYDROXY-4-PHENYLBUTYRIC ACID

Submitted by Li Jun Lei* and Xu Chang He
(06/08/05)

Shanghai Institute of Materia Medica
Shanghai Institutes for Biological Sciences
Chinese Academy of Sciences, 555 Zuchongzhi Road, Shanghai 201203
E-mail: lijunlei@hotmail.com

3-Amino-2-hydroxy-4-phenylbutyric acid (AHPBA) and its derivatives¹ are useful starting materials for the preparation of HIV-1 protease inhibitors such as saquinavir² and amprenavir.³ Various synthetic routes for the preparation of this type of compounds have been published.⁴⁻⁶ We previously reported a novel and facile synthetic route to AHPBA and its derivatives (4*S*,5*S*)-4-benzyl-5-hydroxymethyl oxazolidin-2-one as part of our study of the synthesis of HIV-1 protease inhibitors.⁷ In this paper, *N*-phthaloyl protected *L*-phenylalanine was treated with