

Organoboranes. 29. A convenient synthesis of alkyldibromoboranes and dialkylbromoboranes via hydroboration-redistribution

Herbert C. Brown, D. Basavaiah, and N. G. Bhat

Organometallics, 1983, 2 (10), 1309-1311 • DOI: 10.1021/om50004a007 • Publication Date (Web): 01 May 2002

Downloaded from <http://pubs.acs.org> on April 24, 2009

More About This Article

The permalink <http://dx.doi.org/10.1021/om50004a007> provides access to:

- Links to articles and content related to this article
- Copyright permission to reproduce figures and/or text from this article



ACS Publications
High quality. High impact.

Organoboranes. 29. A Convenient Synthesis of Alkyldibromoboranes and Dialkyldibromoboranes via Hydroboration-Redistribution








Herbert C. Brown,* D. Basavaiah, and N. G. Bhat

Richard B. Wetherill Laboratory, Purdue University, West Lafayette, Indiana 47907

Received March 22, 1983



Trialkylboranes, obtained via hydroboration of terminal and cyclic alkenes with $\text{BH}_3\cdot\text{SMe}_2$, undergo redistribution with boron tribromide at room temperature under the influence of a catalytic amount (7 mol %) of $\text{BH}_3\cdot\text{SMe}_2$ in *n*-pentane to afford the corresponding alkyldibromoboranes or dialkyldibromoboranes, depending upon the stoichiometry, thus providing a convenient method for preparing these valuable derivatives. The reaction of trialkylboranes derived from internal alkenes such as tri-3-hexylborane is much slower. However, these derivatives could be transformed into the corresponding alkyldibromoboranes at 70 °C under neat conditions without significant isomerization. The complete conversion of tri-3-hexylborane into di-3-hexylborane could not be achieved.

The importance of alkyldibromoboranes and dialkyldibromoboranes in organic synthesis has been well documented.¹⁻⁹ Simple procedures are now available for the synthesis of such valuable derivatives. Mono- and dichloroborane etherates ($\text{BH}_2\text{Cl}\cdot\text{OEt}_2$, $\text{BHCl}_2\cdot\text{OEt}_2$), monohaloborane-dimethyl sulfides ($\text{BH}_2\text{X}\cdot\text{SMe}_2$) and dihaloborane-dimethyl sulfides ($\text{BX}_2\cdot\text{SMe}_2$) are the important borane reagents that were developed recently for the synthesis of alkyldibromoboranes (eq 1-5).¹⁰⁻¹⁵




Dibromoborane-dimethyl sulfide ($\text{BHBBr}_2\cdot\text{SMe}_2$) proves to be a more versatile and useful reagent in organic transformations.¹⁻⁴ However, highly hindered alkenes such

as cyclohexene react very sluggishly with this reagent (eq 6).¹⁶ Addition of BBR_3 solves this problem, providing cyclohexyldibromoborane in excellent yields (eq 7).¹⁷



Also, in the case of gaseous alkenes like ethylene, dibromoborane-dimethyl sulfide ($\text{BHBBr}_2\cdot\text{SMe}_2$) fails to provide the desired ethyldibromoborane-dimethyl sulfide ($\text{EtBBR}_2\cdot\text{SMe}_2$) (eq 8). This might be attributed to the nonavailability of ethylene to $\text{BHBBr}_2\cdot\text{SMe}_2$ because ethylene gas escapes. However, the addition of BBR_3 helps in this case also (eq 9).¹⁷



The increasing importance of alkyldibromoboranes in organic synthesis¹⁻⁴ requires a simple method for the preparation of ethyldibromoborane and cyclohexyldibromoborane, as well as other derivatives not readily accessible by direct hydroboration with the dibromoborane.

Earlier methods for the preparation of alkyldibromoboranes involve halogenation of the carbon-boron bond in trialkylboranes by either hydrogen halide or halogen or halogenation of the boron-oxygen bond in trialkylboroxines or alkylboron esters using boron trihalides.^{18,19} Alternatively, alkyldibromoboranes were obtained by the action of trialkylboranes with boron trihalides at high temperatures.^{18,20,21} Zakharkin reported a similar reaction at 125 °C using sodium borohydride as catalyst.²²

- (1) Brown, H. C.; Basavaiah, D. *J. Org. Chem.* **1982**, *47*, 3806.
- (2) Brown, H. C.; Basavaiah, D. *J. Org. Chem.* **1982**, *47*, 5407.
- (3) Brown, H. C.; Basavaiah, D.; Kulkarni, S. U. *J. Org. Chem.* **1982**, *47*, 3808.
- (4) Kulkarni, S. U.; Basavaiah, D.; Zaidlewicz, M.; Brown, H. C. *Organometallics* **1982**, *1*, 212.
- (5) Brown, H. C.; Basavaiah, D. *J. Org. Chem.* **1982**, *47*, 754.
- (6) Brown, H. C.; Basavaiah, D.; Kulkarni, S. U. *J. Org. Chem.* **1982**, *47*, 171.
- (7) Kulkarni, S. U.; Basavaiah, D.; Brown, H. C. *J. Organomet. Chem.* **1982**, *225*, C1.
- (8) Basavaiah, D.; Brown, H. C. *J. Org. Chem.* **1982**, *47*, 1792.
- (9) Basavaiah, D. *Heterocycles* **1982**, *18*, 153.
- (10) Brown, H. C.; Ravindran, N. *J. Am. Chem. Soc.* **1976**, *98*, 1785.
- (11) Brown, H. C.; Ravindran, N. *J. Am. Chem. Soc.* **1976**, *98*, 1798.
- (12) Brown, H. C.; Ravindran, N. *Inorg. Chem.* **1977**, *16*, 2938.
- (13) Brown, H. C.; Ravindran, N.; Kulkarni, S. U. *J. Org. Chem.* **1979**, *44*, 2417.
- (14) Brown, H. C.; Ravindran, N.; Kulkarni, S. U. *J. Org. Chem.* **1980**, *45*, 384.
- (15) Kinberger, K.; Siebert, W. Z. *Naturforsch., B: Anorg. Chem., Org. Chem.* **1975**, *30*, 55.




- (16) Brown, H. C.; Bhat, N. G.; Basavaiah, D., unpublished results.
- (17) Brown, H. C.; Basavaiah, D., unpublished results.
- (18) For a review on "Synthesis of Organohaloboranes", see: Neidenzu, K. *Organomet. Chem. Rev.* **1966**, *1*, 305.
- (19) McCusker, P. A.; Ashby, E. C.; Makowski, H. S. *J. Am. Chem. Soc.* **1957**, *79*, 5182.
- (20) Buls, V. W.; Davis, O. L.; Thomas, R. I. *J. Am. Chem. Soc.* **1957**, *79*, 337.
- (21) Noedritzer, K. "Organometallic Reactions"; Becker, E. I., Tsutsric, M., Eds.; Wiley-Interscience: New York, 1972; Vol. 2, p 1.

Table I. Preparation of Alkylbromoboranes via Redistribution of Trialkylboranes and Boron Tribromide

borane (R ₃ B)	product	yield, ^c %	bp, °C (mm)
triethyl-	ethyl dibromoborane-dimethyl sulfide ^a	77	112-114 (15)
	diethylbromoborane-dimethyl sulfide ^b	80	74-76 (80) [lit. ⁶ 70-72 (75)]
tri- <i>n</i> -propyl-	<i>n</i> -propyl dibromoborane-dimethyl sulfide ^a	83	88-89 (3)
tri- <i>n</i> -hexyl-	<i>n</i> -hexyl dibromoborane ^a	80	69-71 (3) [lit. ¹⁴ 56-58 (0.9)]
	di- <i>n</i> -hexylbromoborane ^b	81	102-103 (4)
tris(2-methyl-1-pentyl)-	2-methyl-1-pentyl dibromoborane ^a	85	60-62 (3.7)
	bis(2-methyl-1-pentyl)bromoborane ^b	81	100-102 (4.5)
tricyclopentyl-	cyclopentyl dibromoborane ^a	82	50-52 (2.8)
	dicyclopentylbromoborane ^b	82	104-105 (6)
tricyclohexyl-	cyclohexyl dibromoborane ^a	78	70-72 (4)
	dicyclohexylbromoborane ^b	84	122-124 (4)
tri-3-hexyl-	3-hexyl dibromoborane ^a	79	60-62 (4.6)

^a All reactions for the preparation of alkyl dibromoboranes were carried out on a 20-mmol (R₃B) scale. ^b All reactions for the preparation of dialkylbromoboranes were carried out on a 30-mmol (R₃B) scale. ^c Yields of isolated products.



Köster and Grassberger have elegantly established that the redistribution of trialkylboranes with boron trihalides, to provide alkyl dihaloboranes and dialkylhaloboranes, is strongly catalyzed by tetraalkylboranes (eq 10 and 11).^{23,24} The redistribution of trialkylboranes with boron trichloride, catalyzed by added diborane, provides a convenient synthetic route to RBCl₂ (eq 12).²⁵



The requirement of cyclohexyldibromoborane and ethyldibromoborane in our studies led us to explore the Köster procedure for the synthesis of alkylbromoboranes. Consequently, we examined the redistribution of representative trialkylboranes with boron tribromide under the influence of a catalytic amount of BH₃·SMe₂ (BMS), a more convenient means of introducing catalytic diborane into the reaction mixture.

Results and Discussion

First, we studied the redistribution of tri-*n*-hexylborane with boron tribromide. Tri-*n*-hexylborane [free from solvent THF and SMe₂, obtained by the hydroboration of 1-hexene with BH₃·SMe₂ (BMS)], reacts readily at room temperature with boron tribromide in *n*-pentane in the presence of 7 mol % BH₃·SMe₂ (BMS). The product is either *n*-hexyldibromoborane (eq 13) or di-*n*-hexylbromoborane (eq 14), depending upon the amount of boron





tribromide used in the redistribution reaction. Thus, treatment of 1 molar equiv of tri-*n*-hexylborane, with 2 molar equiv of boron tribromide, affords *n*-hexyldibromoborane in 82% yield. The ¹¹B NMR spectrum showed a clean peak at δ 64, indicative of an alkyl di-



bromoborane. The ¹¹B NMR spectrum of the methanolized product showed a clean signal at δ 32, indicative of alkyl boronate esters. Similarly, treatment of tri-*n*-hexylborane with boron tribromide, in a 2:1 molar ratio, affords di-*n*-hexylbromoborane (¹¹B NMR δ 80, after methanolysis, δ 54) in 80% yield.

Alkenes of intermediate steric requirements such as 2-methyl-1-pentene and cyclopentene could be converted under the same conditions into the corresponding alkyl dibromoboranes and dialkylbromoboranes in excellent yields (Table I).

More hindered alkenes such as cyclohexene could also be transformed into the corresponding alkyl dibromoboranes and dialkylbromoboranes in very high yields (eq 15, 16) (Table I).




Trialkylboranes derived from gaseous alkenes such as ethylene could be readily converted into the corresponding alkyl dibromoborane or dialkylbromoborane, depending upon the molar ratio of the reactants (eq 17, 18). It should



be mentioned that the preparation of EtBBr₂ by direct hydroboration of the gaseous alkene with BHBr₂·SMe₂ offers difficulties. Thus, the present procedure surmounts this problem, providing a simple and clean synthesis of EtBBr₂.

We did encounter difficulties with the 3-hexyl derivative. Redistribution was very sluggish at room temperature. Attempts at intermediate temperatures such as in refluxing methylene chloride or in refluxing *n*-hexane were unsuccessful. Fortunately, the redistribution proceeded satisfactorily at 70 °C by using the neat reagents, affording 3-hexyldibromoborane in 80% yield (eq 19). No signifi-



cant isomerization occurred, as evidenced by the formation of >99% pure 3-hexanol in the oxidation of the product by alkaline hydrogen peroxide. It was previously noted that the redistribution of tri-3-hexylborane with boron

(22) Zakharkin, L. I.; Kouredov, A. I. *Izv. Akad. Nauk SSSR* 1962, 12, 2247.

(23) Köster, R.; Grassberger, M. A. *Liebigs Ann. Chem.* 1968, 719, 169.

(24) Köster, R.; Binger, P. *Inorg. Synth.* 1974, 15, 149.

(25) Brown, H. C.; Levy, A. B. *J. Organomet. Chem.* 1972, 44, 233. For other reports on redistribution of trialkylboranes with boron trihalides, see refs. 27 and 28.

trichloride produces products in which the boron atom has migrated in part from its original position.²⁵

Unfortunately, the further formation into di-3-hexylbromoborane was very slow, even at still higher temperatures (85 °C), requiring 48 h for only 30% completion. We did not pursue this redistribution further since di-3-hexylbromoborane can be easily prepared by hydroboration of 3-hexene with $\text{BH}_2\text{Br}\cdot\text{SMe}_2$.

Thus the present procedure supplements the direct synthesis of alkylidibromo- and dialkylbromoboranes via hydroboration of alkenes with $\text{BHBr}_2\cdot\text{SMe}_2$ and $\text{BH}_2\text{Br}\cdot\text{SMe}_2$. Particularly, this method demonstrates its utility for the synthesis of ethyldibromoborane and cyclohexyldibromoborane, two derivatives which cannot be obtained easily from $\text{BHBr}_2\cdot\text{SMe}_2$. Internal alkenes such as 3-hexene work well for the synthesis of the alkylidibromoboranes but fail to provide the corresponding dialkylbromoboranes. Presently we are exploring the applications of these alkylbromoboranes for organic transformations.

Experimental Section

All glassware used for the experiments were dried in an oven at 140 °C for several hours, assembled hot, and cooled under a stream of nitrogen. The alkenes used for the study were obtained from Aldrich Chemical Co. or Chemical Samples Co. and were purified by distillation over a small quantity of LiAlH_4 . Triethylborane was obtained from Callery Chemical Co. Reagent grade methanol was used after being stored over 3-Å molecular sieves and *n*-pentane was dried over molecular sieves, type 5 Å. The special experimental techniques used in handling air and moisture sensitive materials are described elsewhere.²⁶ ^1H NMR and ^{11}B NMR spectra were recorded on Varian T-60 and Varian FT-80A spectrometers, respectively. ^{11}B NMR chemical shifts are with reference to $\text{BF}_3\cdot\text{OEt}_2$ (δ 0), and the resonances upfield from the standard are assigned negative signs. GC analyses were carried out on a Varian 1200 FID gas chromatograph (column 12 ft \times 1/8 in. packed with 10% Carbowax 1540 on Chromosorb W 100/120).

Preparation of *n*-Hexyldibromoborane. Tri-*n*-hexylborane (20 mmol) was prepared from 1-hexene and $\text{BH}_3\cdot\text{SMe}_2$ in the usual

way.²⁶ Solvent THF and SMe_2 were removed under reduced pressure. The residual tri-*n*-hexylborane was dissolved in 30 mL of *n*-pentane. Boron tribromide (40 mmol, 3.80 mL) was added, followed by a slow addition of 1.4 mmol of $\text{BH}_3\cdot\text{SMe}_2$ (BMS) in pentane (1.75 mL, 0.8 M) at 0 °C. The reaction mixture was stirred at room temperature for 4 h. ^{11}B NMR of the crude product showed a single peak at δ 64, which is indicative of *n*-hexyldibromoborane. Solvent *n*-pentane was removed, and distillation afforded 12.3 g (80%) of pure *n*-hexyldibromoborane: bp 69–71 °C (3 mm) [lit.¹⁴ 56–58 °C (0.9 mm)]; ^{11}B NMR ($\text{BF}_3\cdot\text{OEt}_2$) δ 64 ((*n*-Hex) BBr_2). The methanolized product showed a single peak at δ 32 ((*n*-Hex) $\text{B}(\text{OMe})_2$). Ethyl-, *n*-propyl-, 2-methyl-1-pentyl-, cyclopentyl-, and cyclohexyldibromoboranes were prepared in identical fashion to the above procedure.

Preparation of Di-*n*-hexylbromoborane. Tri-*n*-hexylborane (30 mmol) was treated with boron tribromide (15 mmol, 1.42 mL) under the same conditions as in the preparation of *n*-hexyldibromoborane to afford 9.52 g (81%) of di-*n*-hexylbromoborane: bp 102–103 °C (4 mm); ^{11}B NMR ($\text{BF}_3\cdot\text{OEt}_2$) δ 82 ((*n*-Hex) $_2\text{BBr}$). Diethyl-, bis(2-methyl-1-pentyl)-, dicyclopentyl-, and dicyclohexyldibromoboranes were prepared in an identical manner.

Preparation of 3-Hexyldibromoborane. To 20 mmol of neat tri-3-hexylborane was added 40 mmol of BBr_3 (3.80 mL) at 0 °C, followed by a slow addition of 0.16 mL of neat $\text{BH}_3\cdot\text{SMe}_2$ (8.8 M). The reaction mixture was stirred at 70 °C for 20 h. Distillation furnished 12.20 g (79%) of 3-hexyldibromoborane: bp 60–62 °C (4.5 mm); ^{11}B NMR ($\text{BF}_3\cdot\text{OEt}_2$) δ 67 ((3-Hex) BBr_2). The methanolized product showed a single peak at δ 31, indicative of (3-Hex) $\text{B}(\text{OMe})_2$. Oxidation of 3-hexyldibromoborane was carried out in the usual way²⁶ by using NaOH and H_2O_2 . The alcohol obtained was found to be >99% pure 3-hexanol on GC analysis.

Acknowledgment. We wish to thank the National Institutes of Health (Grant GM 10937) and the National Science Foundation (Grant CHE 79-18881) for their financial support of this work.

Registry No. $\text{RBBR}_2\cdot\text{SMe}_2$ (R = ethyl), 83967-42-4; $\text{R}_2\text{BBR}\cdot\text{SMe}_2$ (R = ethyl), 86646-22-2; $\text{RBB}_2\cdot\text{SMe}_2$ (R = *n*-propyl), 86646-23-3; RBBR_2 (R = *n*-hexyl), 64770-03-2; R_2BBR (R = *n*-hexyl), 57476-26-3; RBBR_2 (R = 2-methyl-1-pentyl), 72205-96-0; R_2BBR (R = 2-methyl-1-pentyl), 86646-21-1; RBBR_2 (R = cyclopentyl), 64770-09-8; R_2BBR (R = cyclopentyl), 57476-09-2; RBBR_2 (R = cyclohexyl), 6783-09-1; R_2BBR (R = cyclohexyl), 22086-59-5; RBBR_2 (R = 3-hexyl), 64770-05-4; R_3B (R = ethyl), 97-94-9; R_3B (R = *n*-propyl), 1116-61-6; R_3B (R = *n*-hexyl), 1188-92-7; R_3B (R = 2-methyl-1-pentyl), 1188-50-7; R_3B (R = cyclopentyl), 23985-40-2; R_3B (R = cyclohexyl), 1088-01-3; R_3B (R = 3-hexyl), 1883-34-7; BBr_3 , 10294-33-4.

(26) Brown, H. C.; Kramer, G. W.; Levy, A. B.; Midland, M. M. "Organic Syntheses via Boranes"; Wiley-Interscience: New York, 1975.

(27) Costes, J. P.; Cros, G.; Laurent, J. P. *Can. J. Chem.* 1976, 54, 2996.

(28) Krishnamurthy, S. S.; Lappert, M. F.; Pedley, J. B. *J. Chem. Soc., Dalton Trans.* 1975, 1214.

Organoboranes. 30. Convenient Procedures for the Synthesis of Alkyl- and Alkenylboronic Acids and Esters

Herbert C. Brown,* N. G. Bhat,^{1a} and Vishwanatha Somayaji^{1b}

Richard B. Wetherill Laboratory, Purdue University, West Lafayette, Indiana 47907

Received March 22, 1983

Alkyl- and alkenyldibromoborane-dimethyl sulfide complexes, readily obtained by the hydroboration of alkenes and alkynes with dibromoborane-dimethyl sulfide ($\text{HBBR}_2\cdot\text{SMe}_2$), react with water, giving the corresponding boronic acids, and with alcohols and glycols to give the corresponding esters. Various procedures have been developed for the preparation of boronic esters with primary and secondary alcohols, glycols, and tertiary alcohols. Boronic acids react with primary and secondary alcohols reversibly to form the corresponding esters. The equilibrium may be conveniently displaced in favor of ester by carrying out the reaction in pentane, from which the water component separates. This procedure does away with the necessity of azeotrope distillation of a ternary mixture, extensively used previously for the esterification of boronic acids.

Alkyl- and alkenylboronic acids and their esters have found extensive applications in organic synthesis in the

last decade.²⁻⁸ Recently, a number of chiral boronic esters have been utilized for the synthesis of optically active