

tendency to bend inward toward the opposite metal atom, an effect shown also by all the equatorial ligands in 5 (Ru'-Ru-C in the range 85.0 (2)-88.7 (2)°), but only by the carbonyl ligands in 2 (Ru'-Ru-C(1) = 85.1 (3)°), and by an equatorial isocyanide ligand cis to each carbonyl group (Ru'-Ru-C(2) = 86.0 (3)°). The other equatorial isocyanide ligands in 2 bend away from the opposite ruthenium atom (Ru'-Ru-C(3) = 90.5 (3) and Ru'-Ru-C(4) = 92.4 (3)°). Another interesting effect is observed for 5 where a pair of trans radial isocyanides have the phenyl groups bending inward toward the center of the dinuclear unit, while the phenyl groups of the other ligand pair are bent away from the center of the molecule. In this regard, it is noteworthy that the smallest dihedral angles 38.2 (5) and 39.2 (5)° each involve ligands on opposite ruthenium atoms for which the phenyl groups bend away from the center of the dimer. It is thus apparent that in both 2 and 5, the ligand arrangement reflects reduced steric interactions not only within each half of the molecules but also between each half of the dimers. An examination of the nonbonded contacts for 2 and also of space-filling models further suggests that the ligand arrangement observed crystallographically for 2 represents sterically the best possible arrangement and that the dinuclear unit should be considered to be sterically saturated. It is thus conceivable that the low-temperature ¹H NMR effects can be attributed to nonequivalent methyl groups in individual *tert*-butyl isocyanide ligands, while the introduction of further *tert*-butyl isocyanide ligands is to be considered unlikely due to prohibitive steric interactions, thus possibly explaining the lack of further substitution of 2.

In conclusion, it is clear that the complex *catena*-[Ru(CO)₂(O₂CCH₃)] under suitable experimental conditions provides a most unusual synthetic precursor in ruthenium(I) chemistry. We are actively pursuing this further.

Registry No. 1, 26317-70-4; 2, 101377-32-6; 3, 101377-27-9; 4, 101377-29-1; 5, 101377-30-4; 6, 101402-18-0; Me₃CNC, 7188-38-7; PhCH₂NC, 10340-91-7; 2,6-Me₂C₆H₃NC, 2769-71-3; Ru, 7440-18-8.

Supplementary Material Available: Tables of bond lengths, bond angles, atomic coordinates, and thermal parameters for 2 and 5 and tables of analytical and spectroscopic data for 2-6 (43 pages). Ordering information is given on any current masthead page.

Electrochemical Generation and Reactivity of Bis(triphenylphosphine)platinum(0): An Electrosynthesis of Platinum-Acetylene Complexes

Julian A. Davies,* Cassandra T. Eagle,
Deborah E. Otis,[†] and Uma Venkataraman
Department of Chemistry, University of Toledo
Toledo, Ohio 43606

Received August 15, 1985

Summary: Controlled potential bulk reductive electrolysis of *cis*-[PtCl₂(PPh₃)₂] results in the generation of [Pt(PPh₃)₂] in solution. This 2-coordinate, 14-electron compound is efficiently trapped by acetylenes, RC≡CR (R = Ph, COOMe), to produce [Pt(PPh₃)₂(RC≡CR)] quantitatively and by oxidative addition addenda (e.g., C₆H₅COCl, MeI) to produce the corresponding oxidative addition products in high yield. No products arising through ligand disproportionation, Pt(0)/Pt(II) electron transfer, or ligand P-C bond cleavage are produced.

[†] Permanent address: Department of Chemistry, Adrian College, Adrian, Michigan 49221.

Stable, 2-coordinate, 14-electron bis(tertiary phosphine)platinum(0) complexes [PtL₂] are well-known where L is sterically demanding, e.g., P(*t*-Bu)₃,¹ P(*t*-Bu)₂Me,² P(*t*-Bu)₂Ph,¹ P(*c*-Hx)₃,^{1,3} or P(*i*-Pr)₃,¹ with chemical reactivity appearing to increase with decreasing steric demand. The unhindered complex [Pt(PEt₃)₂] has been generated photochemically *in situ* from the oxalate [Pt(C₂O₄)(PEt₃)₂] and shown to be extremely reactive toward coordinating and oxidative addition addenda.^{4,5} It is reported⁶ that platinum(0) complexes can be similarly generated from [Pt(C₂O₄)(PPh₃)₂], but the characterization of several of the reaction products has been questioned.⁷ The generation of [Pt(PPh₃)₂] is problematic because of the tendency for ligand P-C bond cleavage to occur in coordinatively unsaturated complexes containing arylphosphine ligands.⁸ Thus, passing nitrogen through solutions of [Pt(C₂H₄)(PPh₃)₂] results in not only loss of ethylene but also formation of benzene and generation of the ortho-metalated phosphido-bridged cluster [Pt₂(μ-PPh₃)₂(PPh₂C₆H₄)₂].⁹ Similarly, thermolysis¹⁰ of [Pt(PPh₃)₄] produces [Pt₂(μ-PPh₂)₂(PPh₃)₂] and [Pt₃(μ-PPh₂)₃(PPh₃)₂Ph] whereas photolysis¹¹ of [Pt(C₂O₄)(PPh₃)₂] under hydrogen does not produce [PtH₂(PPh₃)₂] (cf. the PEt₃ analogue)^{4,5} but rather, after precipitation with NaBF₄, the cluster [Pt₃(μ-PPh₂)₂(μ-H)(PPh₃)₃].[†]

Use of electrochemical methods to generate low-valent intermediates, such as [Pt(PPh₃)₂], has received scant attention in the past, although such methods may show distinct advantages over thermal and photochemical techniques. In the past it has been reported that controlled potential bulk reductive electrolysis of [PtCl₂(PPh₃)₂], followed by addition of HCl gas,¹² yields a hydrido complex, although no details of characterization were given. Interestingly, electrochemical reduction of [PtCl₂(PPh₃)₂], followed by the addition of water, also yields a hydride,¹³ characterized by an IR absorption at 2232 cm⁻¹, whereas electrochemical reduction in the presence of excess triphenylphosphine is reported¹³ to produce [Pt(PPh₃)₃], characterized by elemental analysis only.

Here we report on the controlled potential bulk reductive electrolysis of *cis*-[PtCl₂(PPh₃)₂]¹⁴ to generate [Pt(PPh₃)₂] and trapping reactions with coordinating and oxidative addition addenda. This method provides a versatile synthesis of a diverse range of platinum(0) and platinum(II) complexes from a common precursor. The electrosynthesis is described in detail for the preparation of [Pt(PPh₃)₂(RC≡CR)] complexes on a ca. 50-mg scale.¹⁵

(1) Otsuka, S.; Yoshida, T.; Matsumoto, M.; Nakatsu, K. *J. Am. Chem. Soc.* 1976, 98, 5850.

(2) Fornies, J.; Green, M.; Spencer, J. L.; Stone, F. G. A. *J. Chem. Soc., Dalton Trans.* 1977, 1006.

(3) Anderson, G. K.; Clark, H. C.; Davies, J. A. *Organometallics* 1982, 1, 550.

(4) Paonessa, R. S.; Prignano, A. L.; Trogler, W. C. *Organometallics* 1985, 4, 647.

(5) Paonessa, R. S.; Trogler, W. C. *Organometallics* 1982, 1, 768.

(6) Blake, D. M.; Nyman, C. J. *J. Am. Chem. Soc.* 1970, 92, 5359.

(7) Blake, D. M.; Leung, L. M. *Inorg. Chem.* 1972, 11, 2879.

(8) Garrou, P. E. *Chem. Rev.* 1985, 85, 171.

(9) Glockling, F.; McBride, T.; Pollock, R. J. *J. Chem. Soc., Chem. Commun.* 1973, 650.


(10) Taylor, N. J.; Chieh, P. C.; Carty, A. J. *J. Chem. Soc., Chem. Commun.* 1975, 448.

(11) Bellon, P. L.; Ceriotti, A.; Demartin, F.; Longoni, G.; Heaton, B. T. *J. Chem. Soc., Dalton Trans.* 1982, 1671.

(12) Mazzocchin, G.; Bontempelli, G.; Nicolini, M.; Crociani, B. *Inorg. Chim. Acta* 1976, 18, 159.


(13) Gavrilova, I. V.; Gel'fman, M. I.; Kiselera, V. M.; Ragumovskii, V. V. *Zh. Neorg. Khim.* 1974, 19, 3045.

(14) Prior cyclic voltammetric studies have been reported. See ref 12 and: Davies, J. A.; Uma, V. *Inorg. Chim. Acta* 1983, 76, L305; *J. Electroanal. Chem.* 1983, 158, 13.




^a Estimated yield by peak heights from ³¹P{¹H} NMR spectrum of reaction product in situ (two values indicate results from two separate preparations). ^b Yield of isolated, recrystallized product. ^c Undergoes slow decarbonylation in solution to produce *trans*-[Pt(CO)(C₆H₅)-(PPh₃)₂][Cl].

The controlled potential bulk electrolysis experiments are typically performed in a three-compartment cell, equipped with a mercury pool working electrode, a platinum spiral wire counterelectrode, and a Ag/AgCl reference electrode (-35 mV vs. SCE). *cis*-[PtCl₂(PPh₃)₂] is dissolved in a preelectrolyzed acetonitrile/benzene (5/2) solvent system containing tetra-*n*-butylammonium perchlorate (TBAP) and suspended activated alumina. Electrolysis at a potential of -1.60 V vs. Ag/AgCl, under an argon atmosphere, leads to the formation of a clear, bright orange solution. Coulometry indicates a two-electron reduction (eq 1). The reduced solution is extremely



air-sensitive, and trace amounts of air lead to formation of substantial quantities of [PtO₂(PPh₃)₂], [Pt(CO)₃(PPh₃)₂], and OPPh₃, which have been identified in solutions after exposure to air by ³¹P{¹H} NMR spectroscopy.

In the absence of air, or any added substrates, a slow disproportionation to [Pt(PPh₃)₃] and platinum metal occurs¹⁶ (eq 2). [Pt(PPh₃)₃] has been identified by ³¹P{¹H}




NMR spectroscopy, and discoloration indicates the formation of platinum metal. The time scale of this disproportionation is sufficiently slow that it does not interfere with trapping reactions of the 2-coordinate intermediate. This is particularly significant since it is known¹⁶ that

(15) Experimental details for the preparation of the [Pt(PPh₃)₂-(PhC≡CPh)] complex are as follows. A rigorously dried three-compartment cell equipped with a mercury pool working electrode (28 cm²) and a Ag/AgCl reference electrode, separated from a platinum spiral counterelectrode by two medium-porosity frits, is charged with 3.0 g of alumina (previously activated at 110 °C for 24 h); acetonitrile (50 mL, previously dried by stirring over CaH₂ followed by successive distillation from CaH₂, P₂O₅, and CaH₂ under nitrogen), benzene (20 mL, previously dried by distillation from sodium under nitrogen), and TBAP (2.3941 g, 7.01 × 10⁻³ mol, recrystallized five times from ethanol and dried in vacuo) and purged with argon for 1 h. The solution is then stirred and preelectrolyzed at -1.60 V to zero current. *cis*-[PtCl₂(PPh₃)₂] (0.0608 g, 7.7 × 10⁻⁵ mol) is added to the working electrode compartment, and the solution is purged with argon for 1 h. The solution is then electrolyzed to zero current at -1.60 V. Coulometry indicates a two-electron reduction with integrations typically in the range of 1.9–2.1 electrons. In this specific case the integration indicated an uptake of 2.09 electrons. The solution is then filtered under argon into a Schlenk tube and the acetylene (1.14 equiv) added. Following reduction to dryness, the solid mass is extracted with benzene (10 mL), filtered, and concentrated to small volume. The solution is then either transferred to an NMR tube under argon and sealed in vacuo for NMR studies or transferred to a short alumina column and chromatographed with benzene/ethanol. Slow evaporation of the eluent allows isolation and collection of the products in 77% yield. Anal. Calcd: C, 66.9; H, 4.5. Found: C, 67.7; H, 5.0. The corresponding complex of MeOOC≡CCOOMe was prepared in 78% yield in an entirely analogous manner.

(16) Abis, L.; Sen, A.; Halpern, J. *J. Am. Chem. Soc.* 1978, 100, 2915.

generation of [Pt(PPh₃)₂] by reductive elimination of methane from the corresponding hydrido methyl complex in toluene solution leads to disproportionation at temperatures above -25 °C. In the present case it seems likely that the solvent system stabilizes the low-valent intermediate. Such stabilization by coordination of benzene⁵ and by coordination of hard donor solvents¹⁷ has been proposed previously.

Trapping reactions have generally been performed by the addition of ca. 1.1 equiv of the substrate to the solution after the reduction is complete, as indicated by coulometry. Yields estimated for products in situ and measured after isolation and purification (Scheme I) are sufficiently high that it is clear that disproportionation (eq 2), electron transfer (eq 3), or other destructive processes (e.g., ligand



P–C bond cleavage,⁸ vide supra) are not significant. Trapping reactions with coordinating and oxidative addition addenda are summarized in Scheme I. In all cases, reaction workup¹⁵ is performed by routine inert-atmosphere techniques, involving separation of the product from the alumina desiccant and the TBAP background electrolyte, followed by in situ examination by ³¹P{¹H} NMR spectroscopy. Yields are estimated from peak heights with deviations from 100% caused by trace quantities of starting material and/or triphenylphosphine oxide. In the case of the acetylene complexes, products were isolated and recrystallized with yields of 77% (R = Ph) and 78% (R = COOMe). These yields are comparable with most traditional syntheses of platinum acetylene complexes¹⁸ and are an improvement on many methods.

The reactivity of electrochemically generated [PtL₂] complexes leads to a number of diverse syntheses which are possible from a common precursor. Since substrates may be added after the electrolysis is complete, the preparative method may be employed with substrates which are themselves electroactive in the potential range of interest. Unlike photochemical^{4,5} or thermal methods,¹⁰ the stability of the trapped product is not a controlling factor with this methodology.

The use of controlled potential electrolysis to generate [PtL₂] intermediates is not restricted to the case where L = PPh₃. We have also examined cases where L = PEt₃ and where L₂ = dppe.¹⁹ In both cases we see enhanced reactivity in comparison with the triphenylphosphine system.

(17) Siedle, A. R.; Newmark, R. A.; Gleason, W. B. *J. Am. Chem. Soc.* 1986, 108, 767.

(18) Lewandos, G. S. In "The Chemistry of the Metal–Carbon Bond"; Hartley, F. R.; Patai, S., Eds.; Wiley: Chichester, 1982; Vol. 1, p 287.

(19) Davies, J. A.; Eagle, C.; Uma, V., unreported results.

Where $L = \text{PEt}_3$, we observe reproducible reaction chemistry with the C-H and/or C-C bonds of acetonitrile and the tetra-*n*-butylammonium cation, depending upon the experimental conditions. A separate report will describe this chemistry in detail.¹⁹

Acknowledgment. Thanks are expressed to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for support of this work. We acknowledge generous loans of platinum metals from Johnson-Matthey, Inc., and Kigre, Inc.

Registry No. *cis*-[PtCl₂(PPh₃)₂], 15604-36-1; [PtO₂(PPh₃)₂], 29894-57-3; [Pt(CO₃)(PPh₃)₂], 17030-86-3; OPPh₃, 791-28-6; [Pt(PPh₃)₃], 13517-35-6; [Pt(PEt₃)₂], 66916-63-0; [Pt(dppe)], 101519-43-1; [Pt(PPh₃)₂(PhC≡CPh)], 15308-61-9; PhC≡CPh, 501-65-5; [Pt(PPh₃)₂(MeOCC≡CCOMe)], 22853-55-0; MeOCC≡CCOMe, 762-42-5; *trans*-[Pt(Me)(PPh₃)₂], 28850-19-3; *trans*-[PtCOC₆H₅(Cl)(PPh₃)₂], 18421-48-2; *trans*-[Pt(CO)(C₆H₅)(PPh₃)₂][Cl], 101519-44-2.

Synthesis of Some Arsinogallanes and the Novel Rearrangement of a Dimeric Bis(arsino)gallane, Bis[bis(trimethylsilyl)methyl]arsino]chlorogallane}


Colin G. Pitt,* Andrew P. Purdy, Kelvin T. Higa, and Richard L. Wells*

Department of Chemistry
Paul M. Gross Chemical Laboratory, Duke University
Durham, North Carolina 27706

Received February 28, 1986




Summary: Silylarsines, $\text{R}_2\text{AsSiMe}_3$ ($\text{R} = \text{Me}_3\text{SiCH}_2$ and mesityl), have been used to synthesize (arsino)gallanes. Dynamic NMR spectroscopy studies of one of these, $\{[(\text{Me}_3\text{SiCH}_2)_2\text{As}]_2\text{GaCl}_2\}_2$, show it has fluxional properties.

Despite the importance of gallium arsenide in emerging semiconductor technology,¹ the chemistry of this elemental combination had, until recently,² been restricted to the studies of Coates et al.³ in the 1960s. They prepared two mono(arsino)gallanes by cleavage of one Ga-C bond in Me_3Ga with secondary arsines, resulting in the elimination of methane (eq 1). In our hands, this method became



impractical, and eventually failed, as the steric bulk of the substituents was increased;⁴ it was also found to be limited

to the cleavage of a single Ga-C bond, and bis- and tris-(arsino)gallanes, $[(\text{R}_2\text{As})_{3-x}\text{GaR}_x]_n$ ($x = 0, 1$), were inaccessible by this route.⁵ A number of reports of the synthesis of main-group compounds by metathetical elimination of trimethylchlorosilane⁶ prompted us to evaluate this route to (arsino)gallanes. Here we report its successful application (eq 2-4), as well as the fluxional properties of a dimeric bis(arsino)gallane.



Silylarsines⁷ are readily available from lithium arsenides and trimethylchlorosilane, and their reaction with trichlorogallane proceeds stepwise in hydrocarbon solvents.⁸

(5) For example, when Ph_2AsH and Me_3Ga (2:1 molar ratio) were heated at 80-100 °C for 2 days, no evidence for any product other than CH_4 , $(\text{Ph}_2\text{AsGaMe}_2)_n$, and unchanged Ph_2AsH was obtained.

(6) See, for example: (a) Abel, E. W.; Armitage, D. A.; Willey, G. R. *J. Chem. Soc.* 1965, 57. (b) Wells, R. L.; Collins, A. L. *Inorg. Chem.* 1966, 5, 1327. (c) Abel, E. W.; Illingsworth, S. M. *J. Chem. Soc. A* 1969, 1094. (d) Harman, J. S.; McCartney, M. E.; Sharp, D. W. A. *J. Chem. Soc. A* 1971, 1547. (e) Goetze, R.; Noeth, H. *Z. naturforsch., B: Anorg. Chem., Org. Chem., Biochem., Biophys., Biol.* 1975, 30B, 875. (f) Nutt, W. R.; Stimson, R. E.; Leopold, M. F.; Rubin, B. H. *Inorg. Chem.* 1982, 21, 1909. (g) Hoffmann, G. G. *J. Organomet. Chem.* 1984, 273, 187.

(7) See for example: (a) Abel, E. W.; Honigschmidt-Grossich, R.; Illingsworth, S. M. *J. Chem. Soc. A* 1968, 2623. (b) Abel, E. W.; Crow, J. P. *J. Organomet. Chem.* 1969, 17, 337. (c) Mislow, K.; Senkler, G. H.; Cook, R. J.; Casey, J. P.; Baechler, R. D. *J. Am. Chem. Soc.* 1972, 94, 2859. The new silylarsines used in this work were prepared under strict inert atmospheric conditions as follows. $(\text{Me}_3\text{SiCH}_2)_2\text{AsSiMe}_3$. A mixture of $(\text{Me}_3\text{SiCH}_2)_2\text{AsH}^{2c}$ (6.0 g, 24 mmol) and *n*-BuLi (10 mL, 2.4 M in hexane) was stirred at 60 °C for 1 day under nitrogen pressure and then cooled to 0 °C, before excess Me_3SiCl (4 mL in 20 mL pentane) was added dropwise. After refluxing for 0.5 h, addition of 10 mL of Et_2O , and standing at room temperature for 1 day, the mixture was filtered; following solvent removal, distillation afforded the silylarsine as a colorless liquid (5.8 g, 75% yield), bp 40 °C (0.04-0.35 torr). Anal. Calcd (Found) for $\text{C}_{11}\text{H}_{31}\text{AsSi}_3$: C, 40.96 (40.83); H, 9.69 (9.82). NMR: ¹H (80 MHz, C_6D_6) δ 0.80 and 0.46 [AB pattern (²J_{HH} = 13.6 Hz), 4 H, CH₂], 0.18 (s, 9 H, AsSiMe₃), 0.15 (s, 18 H, CSiMe₃). ¹³C (22.5 MHz, C_6D_6) δ 5.7 (s, CH₂), -1.5 (s, AsSiMe₃), 0.37 (s, CSiMe₃). $\text{Mes}_2\text{AsSiMe}_3$. (a) Mesitylmagnesium bromide (274 mmol) in THF (200 mL) was added over a 45-min period, with stirring, to AsCl_3 (25.0 g, 138 mmol) in THF; after 12 h, degassed water (125 mL) and concentrated HCl (60 mL) were added, and the mixture was stirred for 1 day. Following extraction with benzene, the volatiles were removed by distillation leaving crude, solid Mes_2AsCl (45.1 g, 94% yield). NMR: ¹H (80 MHz, C_6D_6) δ 6.60 (s, 4 H, ring CH), 2.35 (s, 12 H, Me), 2.02 (s, 6 H Me). (b) The crude Mes_2AsCl was dissolved in THF (500 mL), and LiAlH_4 (10.0 g, 263 mmol) was added. After the solution was stirred for 46 h, addition of water (220 mL), extraction with benzene, and removal of volatiles, distillation afforded Mes_2AsH as a colorless liquid (14.6 g, 58% yield), bp 145-150 °C (0.01 torr). Anal. Calcd (Found) for $\text{C}_{18}\text{H}_{23}\text{As}$: C, 68.79 (68.66); H, 7.38 (7.55). NMR: ¹H (80 MHz, C_6D_6) δ 6.70 (s, 4 H, ring CH), 4.83 (s, 1 H, AsH), 2.27 (s, 12 H, Me), 2.07 (s, 6 H, Me). IR (Nujol): 2130 cm⁻¹ (s, AsH). (c) A mixture of Mes_2AsH (14.62 g, 46.42 mmol), *n*-BuLi (20 mL, 2.4 M in hexane), and ligroin (150 mL) was heated at 50 °C for 20 min. Filtration gave solid, yellow Mes_2AsLi which was dried under vacuum (14.41 g, 96.7% yield). Anal. Calcd (Found) for $\text{C}_{18}\text{H}_{22}\text{AsLi}$: C, 67.51 (67.15); H, 7.82 (7.95). NMR: ¹H (80 MHz, C_6D_6 + 1 drop THF) δ 6.93 (s, 4 H, ring CH), 2.61 (s, 12 H, Me) 2.29 (s, 6 H, Me). (d) A mixture of Mes_2AsLi (3.79 g, 11.8 mmol), Me_3SiCl (10 mL), and ligroin (100 mL) was heated to reflux, stirred at room temperature for 12 h, and filtered. The filtrate was concentrated in vacuo to an oil which eventually crystallized as colorless $\text{Mes}_2\text{AsSiMe}_3$ (4.46 g, 98% yield), mp 62-66 °C. Anal. Calcd (Found) for $\text{C}_{21}\text{H}_{31}\text{AsSi}$: C, 65.26 (65.53); H, 8.08 (8.24). NMR: ¹H (80 MHz, C_6D_6) δ 6.74 (s, 4 H, ring CH), 2.31 (s, 12 H, Me), 2.10 (s, 6 H, Me), 0.28 (s, 9 H, SiMe₃).

(1) Keyes, R. W. *Science (Washington, D.C.)* 1985, 230, 138.

(2) (a) Wells, R. L.; Purdy, A. P.; McPhail, A. T.; Pitt, C. G. *Abstracts of Papers, 189th National Meeting of the American Chemical Society, Miami Beach, FL; American Chemical Society: Washington, DC, 1985; INOR 26.* (b) Wells, R. L.; Purdy, A. P.; McPhail, A. T.; Pitt, C. G. *J. Chem. Soc., Chem. Commun.* 1986, 487. (c) Wells, R. L.; Purdy, A. P.; McPhail, A. T.; Pitt, C. G. *J. Organomet. Chem.*, in press. (d) Pitt, C. G.; Higa, K. T.; McPhail, A. T.; Wells, R. L., submitted for publication in *Inorg. Chem.*

(3) (a) Coates, G. E.; Graham, J. J. *J. Chem. Soc.* 1963, 233. (b) Beachley, O. T.; Coates, G. E. *J. Chem. Soc.* 1965, 3241.

(4) For example, $[(\text{Me}_3\text{SiCH}_2)_2\text{AsGaPh}_2]_2$ was prepared by this method;^{2c} however, heating Ph_2AsH and $(\text{Me}_3\text{SiCH}_2)_2\text{Ga}$ (1:1 molar ratio) at 60 °C for 2 days afforded impure $[(\text{Ph}_2\text{AsGa}(\text{CH}_2\text{SiMe}_3)_2)_n, \text{Ph}_2\text{AsAsPh}_2, \text{Me}_3\text{Si}, \text{H}_2]$, and unreacted $(\text{Me}_3\text{SiCH}_2)_2\text{Ga}$; no evidence for the desired product was obtained when $(\text{Me}_3\text{SiCH}_2)_2\text{AsH}$ and $(\text{Me}_3\text{SiCH}_2)_2\text{Ga}$ (1:1 molar ratio) were heated at 100-120 °C for 1 month.