

# Substituent Effects on the Rates of CO Substitution of $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$

Minsek Cheong and Fred Basolo\*

Department of Chemistry, Northwestern University, Evanston, Illinois 60208

Received April 7, 1988

The syntheses and characterization of the new compounds  $(\eta^5\text{-C}_5\text{H}_4\text{CF}_3)\text{Rh}(\text{CO})_2$  and  $(\eta^5\text{-C}_5\text{H}_4\text{N}(\text{CH}_3)_2)\text{Rh}(\text{CO})_2$  are reported. The compounds  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$  and  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})\text{PPh}_3$  (where X = H, NO<sub>2</sub>, CF<sub>3</sub>, Cl, CH<sub>3</sub>, or N(CH<sub>3</sub>)<sub>2</sub>) show a good linear correlation between the values of CO stretching frequencies,  $\nu_{\text{CO}}$ , and the Hammett  $\sigma$  values of the X ring substituents. This may be the first time such a correlation has been reported, and it is believed to mirror the electron density on the metal which in turn results in greater or lesser  $\pi$ -back-bonding, M $\rightarrow$ CO. In spite of this the rates of CO substitution of these compounds show strong deviations from linear correlation with Hammett  $\sigma$  values. This is rationalized in terms of resonance stabilization of the transition state for reaction by certain ring substituents.

## Introduction

Investigations of associative ligand substitution reactions of 18-electron transition-metal organometallic compounds that were reported 2 decades ago<sup>1</sup> have been revisited, and a rule formulated<sup>1d</sup> that states "substitution reactions of 18-electron metal complexes may proceed by an associative mechanism provided the metal complex can delocalize a pair of electrons onto one of its ligands and make available a vacant low-energy orbital on the metal to permit nucleophilic attack on the metal."

Recent studies<sup>2</sup> show that this process of localizing a pair of electrons on the cyclopentadienyl ligand of its metal complex to allow associative substitution reactions is accelerated by having electron-withdrawing substituents on the cyclopentadienyl ligand. The substituent effects on the reactivity of aromatic molecules have been studied by organic chemists for decades. They have shown that substituent effects can be treated quantitatively by use of a linear free energy relationship, and the most commonly used variant of such an empirical treatment involves use of the Hammett equation (eq 1).<sup>3</sup> In this classical equation

$$\log(k_x/k_0) = \rho\sigma_x \quad (1)$$

tion,  $k_x$  is the rate constant for a compound substituted by a group x,  $k_0$  is the rate constant for the unsubstituted compound,  $\sigma_x$  is a substituent constant, and  $\rho$  is the proportionality factor called reaction constant. Data for literally hundreds of reactions have been successfully correlated by means of this empirical treatment. However, these have been little exploited by the organometallic chemists, although the few examples reported have shown it to be applicable to inorganic reactions.<sup>4</sup>

Equilibrium constants can be treated in a similar way, and even physical properties such as chemical shifts in NMR, frequencies in IR spectra, and polarographic half-wave potentials<sup>5</sup> have been successfully correlated by using the Hammett equation. Egger and Nikiforov<sup>6</sup> have shown

nice linear relationships between the  $\alpha$  and ortho and between the  $\beta$  and meta, respectively, substituents on the cyclopentadienylmanganese tricarbonyls and benzene ring by using <sup>1</sup>H NMR spectra.

No example in the literature of the application of Hammett relationship to CO substitution reactions of substituted cyclopentadienyl metal carbonyls was found. In order to test the applicability of the Hammett relationship on these reactions, we decided to investigate the effects of substituents on the rates of CO substitution of the compounds  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$ .

Kinetic studies on the unsubstituted compound  $(\eta^5\text{-C}_5\text{H}_5)\text{Rh}(\text{CO})_2$ <sup>1a</sup> and the substituted compound  $(\eta^5\text{-C}_5\text{H}_4\text{NO}_2)\text{Rh}(\text{CO})_2$ <sup>2</sup> ( $\sigma_p = 0.78$ ) have been reported. Considering a wide range of Hammett  $\sigma$  values and spacings between them, we decided to study trifluoromethyl (0.54), chloro (0.23), methyl (-0.17), and dimethylamino (-0.60) compounds  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$  ( $\sigma_p$  values<sup>7</sup> are in parentheses). Unfortunately  $(\eta^5\text{-C}_5\text{H}_5)\text{Rh}(\text{CO})_2$  does not undergo aromatic substitution reactions to yield functionally substituted derivatives.<sup>8</sup> As a consequence the syntheses and subsequent chemistry of functionally substituted  $(\eta^5\text{-cyclopentadienyl})\text{rhodium}$  compounds have been severely limited in scope. Alternative methods are necessary for the formation of functionally substituted cyclopentadienylmetal compounds, and these generally involve the reaction of substituted cyclopentadienyl reagents such as  $\text{TiC}_5\text{H}_4\text{X}$ ,  $\text{NaC}_5\text{H}_4\text{X}$ , or  $\text{LiC}_5\text{H}_4\text{X}$  with metal halide fragments. Fortunately the compounds  $(\eta^5\text{-C}_5\text{H}_4\text{Cl})\text{Rh}(\text{CO})_2$ <sup>9</sup> and  $(\eta^5\text{-C}_5\text{H}_4\text{CH}_3)\text{Rh}(\text{CO})_2$ <sup>10</sup> had been synthesized earlier, and thallium (trifluoromethyl)cyclopentadienide<sup>11</sup> and lithium (dimethylamino)cyclopentadienide<sup>12</sup> are known. We report here a new synthesis of the (chlorocyclopentadienyl)rhodium carbonyl (I) and what we believe are the first reported syntheses of II and IV. All four compounds were studied kinetically for CO substitution reactions, and the Hammett relationship observed is discussed for the mechanisms proposed.

(1) (a) Schuster-Woldan, H. G.; Basolo, F. *J. Am. Chem. Soc.* **1966**, *88*, 1657. (b) Thorsteinson, E. M.; Basolo, F. *J. Am. Chem. Soc.* **1966**, *88*, 3927. (c) Morris, D. E.; Basolo, F. *J. Am. Chem. Soc.* **1968**, *90*, 251. (d) Basolo, F. *Inorg. Chim. Acta* **1985**, *100*, 33.

(2) Rerek, M. E.; Basolo, F. *Organometallics* **1983**, *2*, 372; *J. Am. Chem. Soc.* **1984**, *106*, 5908.

(3) Hammett, L. P. *J. Am. Chem. Soc.* **1937**, *59*, 96.

(4) (a) Aprile, F.; Cagliotti, V.; Illuminati, G. *J. Inorg. Nucl. Chem.* **1961**, *21*, 325. (b) Jones, W. E.; Thomas, J. D. R. *J. Chem. Soc. A* **1966**, 1481.

(5) Nesmeyanov, A. N.; Denisovich, L. I.; Gubin, S. P.; Vol'kenau, N. A.; Sirotkina, E. I.; Bolesova, I. N. *J. Organomet. Chem.* **1969**, *20*, 169.

(6) Egger, E.; Nikiforov, A. *Monatsh. Chem.* **1969**, *100*, 483.

(7) Wells, P. R. *Chem. Rev.* **1963**, *63*, 171.


(8) Hart, W. P.; Macomber, D. W.; Rausch, M. D. *J. Am. Chem. Soc.* **1980**, *102*, 1196.

(9) Conway, B. G.; Rausch, M. D. *Organometallics* **1985**, *4*, 688.

(10) Arthurs, M.; Nelson, S. M.; Drew, M. G. B. *J. Chem. Soc., Dalton Trans.* **1977**, 779.

(11) (a) Olsson, T.; Wennerström, O. *Acta Chem. Scand., Ser. B* **1978**, *B32*, 293. (b) Gassman, P. G.; Winter, C. H. *J. Am. Chem. Soc.* **1986**, *108*, 4228.

(12) Bernheim, M.; Boche, G. *Angew. Chem., Int. Ed. Engl.* **1980**, *19*, 1010.



### Experimental Section

**Compounds and Solvents.** All manipulations were routinely carried out by using standard Schlenk techniques under an atmosphere of  $N_2$ , excluding oxygen and moisture. Hexane, tetrahydrofuran, and Decalin were dried and distilled over Na in the presence of benzophenone under  $N_2$  atmosphere. The hexane and Decalin were stored over concentrated  $H_2SO_4$  and washed with KOH before distillation. Chlorodicarbonylrhodium(I) dimer<sup>13</sup> and 5-diazo-1,3-cyclopentadiene<sup>14</sup> were prepared by published procedures.  $PPh_3$  (Aldrich) was recrystallized from methanol prior to use.

**( $\eta^5$ -Chlorocyclopentadienyl)dicarbonylrhodium(I), ( $\eta^5$ - $C_5H_4Cl$ )Rh(CO)<sub>2</sub> (I).** To a solution of  $[RhCl(CO)_2]_2$  (0.17 g) in THF (30 mL) in a nitrogen-filled 50-mL Schlenk flask connected to a nitrogen line equipped with a bubbler was added a portion of 1.8 M pentane solution of 5-diazo-1,3-cyclopentadiene (0.5 mL). Bubbles of gas were evolved. The reaction was allowed to proceed overnight, and then the solvent was removed under vacuum. The product was dissolved in 20 mL of hexane and filtered. The solvent was removed from the filtrate to yield 0.07 g (31% yield) of a yellow oil: IR (Decalin)  $\nu(CO)$  2053, 1991; IR (hexane)  $\nu(CO)$  2055, 1994  $cm^{-1}$ ;  $^1H$  NMR ( $C_6D_6$ )  $\delta$  5.42 (t, 2 H), 4.82 (t, 2 H); mass spectrum,  $m/e$ ,  $M^+$  258. The  $^1H$  NMR and mass spectra agree with literature results.<sup>9</sup>

**[ $\eta^5$ -(Trifluoromethyl)cyclopentadienyl]dicarbonylrhodium(I), ( $\eta^5$ - $C_5H_4CF_3$ )Rh(CO)<sub>2</sub> (II).** In a 50-mL Schlenk flask was placed 1.9 g of  $TiCl_3H_4CF_3$ ,<sup>11</sup> 0.5 g of  $[RhCl(CO)_2]_2$ , and 30 mL of THF. The resultant mixture was stirred for 2 h. The solvent was removed under vacuum, 40 mL of hexane added, and the mixture suction filtered through a medium frit. Hexane was removed from the filtrate to yield 0.19 g (32% yield) of a yellow oil: IR (Decalin)  $\nu(CO)$  2060, 1999  $cm^{-1}$ ;  $^1H$  NMR ( $CHCl_3-d$ )  $\delta$  5.51 (s, 2 H), 5.76 (s, 2 H); mass spectrum,  $m/e$  (relative intensity)  $M^+$  292 (33.9), (M-CO)<sup>+</sup> 264 (45.3), (M-2CO)<sup>+</sup> 236 (100).

**( $\eta^5$ -Methylcyclopentadienyl)dicarbonylrhodium(I), ( $\eta^5$ - $C_5H_4CH_3$ )Rh(CO)<sub>2</sub> (III).** This compound was prepared in a similar manner to II, using  $NaC_5H_4CH_3$  to give the product as a yellow oil: IR (Decalin)  $\nu(CO)$  2041, 1978; IR (THF)  $\nu(CO)$  2036, 1970  $cm^{-1}$ ;  $^1H$  NMR ( $CDCl_3-d$ )  $\delta$  2.03 (s, 3 H), 5.20 (t, 2 H), 5.39 (t, 2 H); mass spectrum,  $m/e$  (relative intensity)  $M^+$  238 (45.7), (M-CO)<sup>+</sup> 210 (52.1), (M-2CO)<sup>+</sup> 182 (100).

**[ $\eta^5$ -(Dimethylamino)cyclopentadienyl]dicarbonylrhodium(I), ( $\eta^5$ - $C_5H_4N(CH_3)_2$ )Rh(CO)<sub>2</sub> (IV).** *N,N*-Dimethyl-1,3-cyclopentadienylamine was prepared by reacting  $NaC_5H_5$  with *N,N*-dimethyl-*O*-(toluenesulfonyl)hydroxylamine in THF at  $-20^\circ C$ . Compound IV was synthesized in manners analogous to those used for II. The orange crystalline product was purified by sublimation under vacuum (ca.  $40^\circ C$ ): IR (Decalin)  $\nu(CO)$  2030, 1965; IR (THF) 2020, 1955  $cm^{-1}$ ;  $^1H$  NMR ( $CH_2Cl_2-d$ )  $\delta$  2.41 (s, 6 H), 4.96 (t, 2 H), 5.10 (t, 2 H); mass spectrum,  $m/e$  (relative intensity)  $M^+$  267 (42.2), (M-CO)<sup>+</sup> 239 (48.3), (M-2CO)<sup>+</sup> 211 (100). Anal. Calcd for  $C_9H_{11}NO_2Rh$ : C, 40.47; H, 3.77; N, 5.24. Found: C, 40.55; H, 3.71; N, 5.26.

**[( $\eta^5$ -Trifluoromethyl)cyclopentadienyl]carbonyl(triphenylphosphine)rhodium(I), ( $\eta^5$ - $C_5H_4CF_3$ )Rh(CO)PPh<sub>3</sub> (V).** To a 5-mL hexane solution of II (0.095 g) was added  $PPh_3$  (0.085 g) (1:1) by ampule transfer. The solution was stirred at room temperature for ca. 3 days, during which time a yellow precipitate formed. This was filtered and dried to afford the product as a yellow powder: IR (hexane)  $\nu(CO)$  1972  $cm^{-1}$ ; mass spectrum,  $m/e$  (relative intensity)  $M^+$  526 (13.4), (M-CO)<sup>+</sup> 498 (100), (M-CO-PPh<sub>3</sub>)<sup>+</sup> 236 (23.7). Anal. Calcd: C, 57.05; H, 3.64. Found: C, 57.12; H, 3.84.

$PPh_3$  derivatives for the (chlorocyclopentadienyl)-, (methylcyclopentadienyl)-, or [(dimethylamino)cyclopentadienyl]rhodium

**Table I. Rate Constants and Activation Parameters for the CO Substitution Reactions of ( $\eta^5$ - $C_5H_4X$ )Rh(CO)<sub>2</sub> with  $PPh_3$  in Decalin (Eq 2)**

X	T, °C	$k_{CO}$ , $M^{-1} s^{-1}$	$\Delta H^\ddagger$ , kcal/mol	$\Delta S^\ddagger$ , eu
NO <sub>2</sub> <sup>a</sup>	25	1.26	+12.0 (±0.2)	-17.9 (±0.6)
	37	2.58		
	49	6.14		
CF <sub>3</sub>	15	$6.16 \times 10^{-3}$	+10.2 (± 1.3)	-30 (±3)
	25	$1.58 \times 10^{-2}$		
	35	$2.17 \times 10^{-2}$		
Cl	25	$5.18 \times 10^{-2}$		
	25	$1.3 \times 10^{-4c}$	+15 <sup>d</sup>	-26 <sup>d</sup>
CH <sub>3</sub>	25	$9.98 \times 10^{-5}$	+15.6 (±2.0)	-27 (±4)
	35	$1.89 \times 10^{-4}$		
	45	$5.20 \times 10^{-4}$		
	45	$3.10 \times 10^{-3}$		
N(CH <sub>3</sub> ) <sub>2</sub>	25	$1.05 \times 10^{-3}$	+9.5 (±1.4)	-40 (±3)
	35	$1.73 \times 10^{-3}$		
	45	$3.10 \times 10^{-3}$		

<sup>a</sup> Reference 2. <sup>b</sup> Reference 1a. <sup>c</sup> Estimated from rate of reaction with  $PPh_3$  at  $40^\circ C$  and activation parameters of reaction with  $P(n-OC_4H_9)_3$ . <sup>d</sup> For the reaction with  $P(n-OC_4H_9)_3$ .

**Table II. Carbonyl Stretching Frequencies of the ( $\eta^5$ - $C_5H_4X$ )Rh(CO)<sub>2</sub> and ( $\eta^5$ - $C_5H_4X$ )Rh(CO)PPh<sub>3</sub> in Decalin**

R	dicarbonyl	$\nu_{CO}$ , $cm^{-1}$	monocarbonyl	$\nu_{CO}$ , $cm^{-1}$
NO <sub>2</sub>	1	2067, 2011	7	1980 <sup>a</sup>
CF <sub>3</sub>	2	2060, 1999	8	1969
Cl	3	2053, 1991	9	1963
H	4	2051, 1987 <sup>b</sup>	10	1957 <sup>a</sup>
CH <sub>3</sub>	5	2041, 1978	11	1950
N(CH <sub>3</sub> ) <sub>2</sub>	6	2030, 1965	12	1941

<sup>a</sup> Hexane. <sup>b</sup> Cyclohexane.



carbonyls were synthesized in manners analogous to those used for V. The IR carbonyl stretches of all products agree closely with those obtained in kinetic measurements.

**Instrumentation.**  $^1H$  NMR spectra were recorded on a Varian EM-390 spectrometer. Deuterated solvents were dried and degassed. IR spectra were recorded on a Nicolet 7199 FT-IR spectrometer using either a Specac P/N 20.500 variable-temperature IR cell which contained a 0.5-mm AgCl cell or 0.2-mm CaF<sub>2</sub> cells. Mass spectra were obtained by Dr. D. Hung of Northwestern University Analytical Services Laboratory on a HP 5985 A spectrometer using 70 eV ionization. Elemental analyses for solid samples were carried out by Galbraith Laboratories, Inc. (Knoxville, TN).

**Kinetic Measurements.** Rate data for the disappearance of rhodium dicarbonyls were obtained by monitoring spectral changes in the  $\nu(CO)$  region of the IR. In a typical experiment, the solutions were approximately 5 mM in rhodium dicarbonyls, and the reactions were carried out under pseudo-first-order conditions with at least a 10-fold excess of  $PPh_3$ . The IR cells were sealed with rubber septa and flushed with  $N_2$  prior to use. Decalin solutions were placed in the constant temperature bath. Constant temperature was maintained by an external circulating bath (Neslab RTE-8). After thermal equilibrium the two solutions were mixed and placed into the IR cell. During the reaction, temperature was maintained within  $\pm 0.05^\circ C$ . IR spectra changes for the disappearance of the highest carbonyl absorption band were monitored. Plots of  $-\ln A$  vs time for the disappearance of reactant rhodium dicarbonyl and plots of  $-\ln(A_\infty - A)$  vs time for the appearance of product rhodium monocarbonyl were linear over 3-4 half-lives (linear correlation coefficient >0.99). The slopes of these lines yielded  $k_{obsd}$ . Both methods for monitoring the reaction rates yielded similar  $k_{obsd}$  values of a given reaction.


### Results

The reaction of I, II, III, or IV with  $PPh_3$  yields a monosubstituted product as shown in eq 2. Typical IR




(13) McCleverty, J. A.; Wilkinson, G. *Inorg. Synth.* 1966, 8, 211.

(14) (a) Weil, T.; Cais, M. *J. Org. Chem.* 1963, 28, 2472. (b) Reimer, K. J.; Shaver, A. *Inorg. Synth.* 1980, 20, 188.



**Figure 1.** Infrared spectral changes for the reaction  $(\eta^5\text{-C}_5\text{H}_4\text{CF}_3)\text{Rh}(\text{CO})_2 + \text{PPh}_3 \rightarrow (\eta^5\text{-C}_5\text{H}_4\text{CF}_3)\text{Rh}(\text{CO})\text{PPh}_3 + \text{CO}$  in Decalin at 25.0 °C.




**Figure 2.** Plot of  $k_{\text{obsd}}$  vs  $\text{PPh}_3$  for the reaction  $(\eta^5\text{-C}_5\text{H}_4\text{CF}_3)\text{Rh}(\text{CO})_2 + \text{PPh}_3 \rightarrow (\eta^5\text{-C}_5\text{H}_4\text{CF}_3)\text{Rh}(\text{CO})\text{PPh}_3 + \text{CO}$  in Decalin at various temperatures.

spectral changes in the carbonyl region for the CO substitution reaction of II with  $\text{PPh}_3$  in Decalin at 25 °C are shown in Figure 1. Rate constants and activation parameters for CO substitution reactions of I, II, III, or IV with  $\text{PPh}_3$  appear in Table I. Kinetic results for the CO substitution reactions support an associative mechanism. The rates of reactions are first order in complex and first order in  $\text{PPh}_3$  for all compounds, with activation parameters of  $\Delta H^\ddagger < 20$  kcal/mol and  $\Delta S^\ddagger < -15$  eu. No evidence for a dissociative pathway was obtained. Figure 2 shows plots of observed CO substitution rate constants at different temperatures vs  $\text{PPh}_3$  concentrations for the reaction of II with  $\text{PPh}_3$ . Table II contains the carbonyl stretches of the various substrates and their substituted products.

### Discussion

Most of the substituted cyclopentadienylrhodium carbonyls synthesized are compounds with electron-withdrawing substituents, such as  $\text{CH}=\text{CH}_2$ ,<sup>15</sup>  $\text{CHO}$ ,<sup>8</sup>  $\text{COCH}_3$ ,<sup>8</sup>



**Figure 3.** Plot of Hammett  $\sigma_p$  values vs  $\nu_{\text{CO}}$ 's for the cyclopentadienylrhodium carbonyls (see Table II for identity of specific compounds): O,  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$ ;  $\Delta$ ,  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})\text{PPh}_3$ .

or  $\text{COOCH}_3$ .<sup>8</sup> Cyclopentadienylrhodium carbonyls substituted with electron-donating groups are compounds with pentamethyl<sup>16</sup> or pentaphenyl<sup>17</sup> substituents. Although it is known<sup>2</sup> that the electron-withdrawing group  $\text{NO}_2$  on the cyclopentadienyl ring of  $(\eta^5\text{-C}_5\text{H}_5)\text{Rh}(\text{CO})_2$  increases its rate of CO substitution reaction by  $10^4$ , no systematic study of the effect of ring substituents on the rates of CO substitution of such metal carbonyls has been reported. The effect of electron-donating groups is limited to  $(\eta^5\text{-C}_5(\text{CH}_3)_5)\text{Rh}(\text{CO})_2$  which decreases the rate of CO substitution by  $10^{-2}$  relative to the parent compound, but some of this decrease is surely due to steric factors.

For associative reactions of 18-electron organometallic compounds, an  $\eta^5$  to  $\eta^3$  rearrangement mechanism has been proposed<sup>1,18</sup> (see eq 3). By putting electron-donating as well as electron-withdrawing substituents on the cyclopentadienyl ring, we now are able to consider the  $\eta^5 \rightarrow \eta^3 \rightarrow \eta^5$  ring slippage in greater detail. Figure 3 shows the dependence of the CO stretching frequencies  $\nu_{\text{CO}}$  for the cyclopentadienylrhodium carbonyls on the Hammett  $\sigma_p$  values. The plot is reasonably linear, with no marked deviation for any substituent. This suggests consistent effects for substituents of the cyclopentadienyl ring on the CO frequencies of rhodium carbonyls, be they the reactant carbonyls  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})_2$  or the product carbonyls  $(\eta^5\text{-C}_5\text{H}_4\text{X})\text{Rh}(\text{CO})\text{PPh}_3$ . It is well-known<sup>19</sup> that for such systems the values of  $\nu_{\text{CO}}$  increase when the electron density on the metal decreases and they decrease when the electron density on the metal increases. That is exactly what happens here; for positive values of  $\sigma_p$  (electron-withdrawing groups) the  $\nu_{\text{CO}}$  values increase, whereas for negative values of  $\sigma_p$  (electron-donating groups) the  $\nu_{\text{CO}}$  values decrease. This may be the first time such a good linear correlation between values of  $\nu_{\text{CO}}$  and Hammett  $\sigma$  values for metal carbonyls have been reported.

On the basis of such a good correlation between  $\nu_{\text{CO}}$  and  $\sigma$  values, perhaps reflecting variations in electron densities


(15) Macomber, D. W.; Hart, W. P.; Rausch, M. D.; Priester, R. D.; Pittman, C. U., Jr. *J. Am. Chem. Soc.* **1982**, *104*, 884.

(16) Kang, J. W.; Maitlis, P. M. *J. Organomet. Chem.* **1971**, *26*, 393.

(17) Connelly, N. G.; Raven, S. J. *J. Chem. Soc., Dalton Trans.* **1986**, 1613.

(18) Cramer, R.; Seiwel, L. P. *J. Organomet. Chem.* **1975**, *92*, 245.


(19) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 4th ed.; Wiley: New York, 1980; Chapter 25.



**Figure 4.** Plot of Hammett  $\sigma_p$  values vs  $\log k_{CO}$  for CO substitution reactions of cyclopentadienylrhodium carbonyls in Decalin at 25 °C.


on the metal, it was expected that a similar correlation would be found between the Hammett  $\sigma$  values of the substituents and the rates of CO substitution of the metal carbonyls. This is expected on the basis of the associative  $\eta^5 \rightarrow \eta^3 \rightarrow \eta^5$  mechanism<sup>1,2</sup> which involves delocalization of electron density from the metal to the ring ligand. However, Figure 4 shows there are strong deviations from linearity of a plot of  $\log k_{CO}$  vs Hammett  $\sigma_p$  values. Four of the substituents (H,  $\text{CH}_3$ ,  $\text{CF}_3$ , and  $\text{NO}_2$ ) give a satisfactory correlation, but two of the substituents ( $\text{N(CH}_3)_2$  and Cl) on this basis give compounds that react much faster than expected.

One difference between the groups  $\text{N(CH}_3)_2$  and Cl and the other four ring substituents is that these contain one or more nonbonding electron pairs that may get involved to help stabilize the transition state for reaction. The associative  $\eta^5 \rightarrow \eta^3 \rightarrow \eta^5$  mechanism for CO substitution (eq 3) is also supported by X-ray<sup>20a</sup> and core and valence



ionization studies<sup>20b</sup> that show allyl-ene distortion in such

ring-substituted cyclopentadienyl compounds of the cobalt triad of elements. The  $\eta^3$ -allyl-ene ring structure is believed to be the structure of the transition state or active intermediate for CO substitution reactions, and it may be that this structure can be resonance stabilized by ring substituents with nonbonding pairs of electrons, as shown by (4). Such resonance stabilization of the transition state



for reaction may account for the electron donor group  $\text{N(CH}_3)_2$  increasing the rate of substitution by 8 times over the unsubstituted parent compound, when the mechanism proposed (eq 3) suggests it should decrease the rate of reaction. The fact that in the ground-state good linearity of the Hammett plot (Figure 3) is observed for all substituents and the failure to see a similar linearity of the Hammett plot for rates of reaction (Figure 4) resides in how substituents affect the transition state. Also a poor fit of  $\log k$  with  $\sigma^+$  or  $\sigma^-$  values indicates the substituent does not enter into direct resonance interaction with the reaction site in the transition state. Instead, stabilization of the transition state comes from stabilization of the localized double bond away from the reaction site of the transition state.

The  $\rho$  value ( $= +4.0$ ) for this reaction series, excluding Cl and  $\text{N(CH}_3)_2$ , is quite large. The magnitude of  $\rho$  is a measure of the amount of charge developed at the reaction center and the extent to which the substituents are able to interact with developing charge. Therefore, very large  $\rho$  values for CO substitution of cyclopentadienylrhodium carbonyls indicate that substituents on the cyclopentadienyl ring can interact extensively with an electron pair from the 18-electron rhodium center, stabilizing or destabilizing the electron-rich transition state.

**Acknowledgment.** This research was supported by a National Science Foundation Grant (CHE-8514366). We wish to thank Professor Paul G. Gassman for helpful information on the synthesis of  $\text{TiC}_5\text{H}_4\text{CF}_3$ . We are grateful to the Johnson-Matthey Corp. for its loan of the Rh used in this study.

**Registry No.** 1, 75862-04-3; 2, 115406-90-1; 3, 94890-74-1; 4, 12192-97-1; 5, 63544-85-4; 6, 115406-91-2; 7, 91842-67-0; 8, 115406-92-3; 9, 115406-93-4; 10, 12203-88-2; 11, 115406-94-5; 12, 115406-95-6;  $[\text{RhCl(CO)}_2]_2$ , 14523-22-9;  $\text{TiC}_5\text{H}_4\text{CF}_3$ , 102649-16-1;  $\text{NaC}_5\text{H}_4\text{CH}_3$ , 55562-83-9;  $\text{NaC}_5\text{H}_5$ , 4984-82-1; 5-diazo-1,3-cyclopentadiene, 1192-27-4; *N,N*-dimethyl-1,3-cyclopentadienylamine, 75812-62-3; *N,N*-dimethyl-*O*-(toluenesulfonyl)hydroxylamine, 32884-91-6.

(20) (a) Byers, L. R.; Dahl, L. F. *Inorg. Chem.* **1980**, *19*, 277. (b) Lichtenberger, D. L.; Calabro, D. C.; Kellogg, G. E. *Organometallics* **1984**, *3*, 1623.