

Competition between Intramolecular and Intermolecular C-H Activation in Iridium Ethene Complexes

Tanachat W. Bell, Sally-Ann Brough, Martin G. Partridge, Robin N. Perutz,* and A. Denise Rooney*


Department of Chemistry, University of York, York YO1 5DD, U.K.

Received February 25, 1993

The ethene complexes $\text{CpIr(L)(C}_2\text{H}_4)$ ($\text{Cp} = \eta^5\text{-C}_5\text{H}_5$, $\text{L} = \text{PPh}_3, \text{CO}$) undergo two competing photochemical reactions in solution: (i) isomerization to the vinyl hydride $\text{CpIr(L)(C}_2\text{H}_3)\text{H}$ and (ii) dissociation of ethene and insertion into solvent C-H bonds to form CpIr(L)(R)H . The vinyl hydride is favored over the intermolecular product by longer wavelength photolysis, lower temperature, and a more rigid surrounding medium and by $\text{L} = \text{PPh}_3$ compared to $\text{L} = \text{CO}$. The vinyl hydride isomer is the exclusive product of UV irradiation of $\text{CpIr(CO)(C}_2\text{H}_4)$ in solid toluene (77 K) or solid argon (12 K) but competes with formation of CpIr(CO)_2 in CO-doped Ar matrices and $\text{CpIr(CO)(CH}_3)\text{H}$ in methane matrices. The latter is also formed by secondary photolysis of the vinyl hydride. The barrier for the thermal conversion of $\text{CpIr(L)(C}_2\text{H}_3)\text{H}$ to the parent ethene complex $\text{CpIr(L)(C}_2\text{H}_4)$ increases in the following order: $\text{L} = \text{C}_2\text{H}_4$, $\Delta H^\ddagger = 67 \pm 5 \text{ kJ mol}^{-1}$; $\text{L} = \text{CO}$, $\Delta H^\ddagger = 95 \pm 4 \text{ kJ mol}^{-1}$; $\text{L} = \text{PPh}_3$, $\Delta G^\ddagger = 135 \pm 3 \text{ kJ mol}^{-1}$ at 318 K. These findings are consistent with our proposal that C-H bond activation occurs via a "cage-complex" intermediate. However, the marked influence of temperature on the reaction indicates that intermolecular C-H bond activation involves a thermally activated step. The photochemical reaction of $\text{CpIr(C}_2\text{H}_4)_2$ with PPh_3 in CD_3CN generates the vinyl hydride $\text{CpIr(PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ via $\text{CpIr(PPh}_3)(\text{C}_2\text{H}_4)$. The corresponding reaction occurs in benzene with the additional formation of the solvent activation product $\text{CpIr(PPh}_3)(\text{Ph})\text{H}$.


Introduction

The photoinduced isomerization of metal ethene to metal vinyl hydride complexes has proved to be a widespread phenomenon, especially among complexes of second- and third-row transition metals.¹⁻⁸ The metal vinyl hydrides are usually thermally unstable with respect to the precursor complexes (eq 1), but the temperature



required for this back-reaction varies according to the metal and ancillary ligands. The span of temperatures necessary to induce the back-reaction is surprisingly large, varying from -20°C for $[\text{M}] = \text{CpRh(PMe}_3)$ to $+180^\circ\text{C}$ for $[\text{M}] = \text{Cp}^*\text{Ir(PMe}_3)$ ($\text{Cp} = \eta^5\text{-C}_5\text{H}_5$, $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$). For some tris(pyrazolyl)borate complexes, the relative stabilities of the ethene complex and the vinyl hydride complex are reversed, so that the forward reaction of eq 1 may be achieved thermally as well as photochemically.^{9,10} It has been postulated that the increased thermal stability of these vinyl hydride complexes is associated with

Scheme I. Cage-Complex Mechanism for the Isomerization of the Ethene Complexes



conversion of the tris(pyrazolyl)borate ligand from η^2 - to η^3 -coordination.

In a previous paper, we proposed that the photoisomerization reactions proceed via cage complexes between metal and expelled ethene (Scheme I).⁵ The extent of interaction within the cage is postulated to govern the proportion of photoisomerization compared to reaction with solvents or added ligands. When the interaction in the cage is strong, photoisomerization is dominant and may be observed at room temperature in preference to solvent activation (e.g. for $\text{CpIr(PMe}_3)(\text{C}_2\text{H}_4)$). When the interaction is weak, as for $\text{CpRh(PMe}_3)(\text{C}_2\text{H}_4)$ or $\text{CpIr(C}_2\text{H}_4)_2$, photoisomerization is only significant in solid matrices such as toluene or argon at low temperature. The isomerization of $(\eta^6\text{-mesitylene})\text{Os(C}_2\text{H}_4)\text{CO}$ represents an intermediate case.⁶ Thus there is a gradation of "intramolecularity" between one complex and another. The order of increasing intramolecularity correlates with the order of stability of the vinyl hydride complexes with respect to metal ethene complexes. Our postulate of the cage complex is equivalent to Bergman's " σ -complex" formed as an intermediate in the thermal reaction of $\text{Cp}^*\text{Ir(PMe}_3)(\text{cyclohexyl})\text{H}$ with ethene.¹ There is one observation of such a σ -complex,

(1) Stoutland, P. O.; Bergman, R. G. *J. Am. Chem. Soc.* 1988, 110, 5732.

(2) Haddleton, D. M.; Perutz, R. N. *J. Chem. Soc., Chem. Commun.* 1986, 1734.

(3) Baker, M. V.; Field, L. D. *J. Am. Chem. Soc.* 1986, 108, 7433, 7436.

(4) Wenzel, T. T.; Bergman, R. G. *J. Am. Chem. Soc.* 1986, 108, 4856.

(5) Bell, T. W.; Haddleton, D. M.; McCamley, A.; Partridge, M. G.; Perutz, R. N.; Willner, H. *J. Am. Chem. Soc.* 1990, 112, 9212.

(6) McCamley, A.; Perutz, R. N.; Stahl, S.; Werner, H. *Angew. Chem., Int. Ed. Engl.* 1989, 28, 1690.

(7) Tanke, R. S.; Crabtree, R. H. *Inorg. Chem.* 1989, 28, 3444.


(8) Schulz, M.; Werner, H. *Organometallics* 1992, 11, 2790.

(9) Ghosh, C. K.; Hoyano, J. K.; Krentz, R.; Graham, W. A. G. *J. Am. Chem. Soc.* 1989, 111, 5480.

(10) Perez, P. J.; Poveda, M. L.; Carmona, E. *J. Chem. Soc., Chem. Commun.* 1992, 8.

viz. $\text{Fe}(\eta^1\text{-C}_2\text{H}_4)$ observed in low temperature matrices; this complex does indeed isomerize to $\text{Fe}(\text{C}_2\text{H}_3)\text{H}$, albeit photochemically.¹¹

An alternative mechanism for the isomerization of the $\text{CpM}(\text{L})(\text{C}_2\text{H}_4)$ complexes is that proposed by Tanke and Crabtree for (tris(pyrazolyl)borato)bis(ethylene)iridium-(I).⁷ They postulated that the isomerization of this complex is initiated by expulsion of one of the ethene ligands. A more general version of this mechanism applied to the compounds studied here is given in eq 2.



In this paper, we examine a series of reactions of $(\eta^5\text{-cyclopentadienyl})(\text{ethene})\text{iridium complexes}$, $\text{CpIr}(\text{C}_2\text{H}_4)\text{L}$ ($\text{L} = \text{CO}, \text{PPh}_3, \text{C}_2\text{H}_4$), in which there is competition between photoisomerization and attack on other ligands or on the solvent/matrix. Since the thermal stability of the vinyl hydride complexes is of particular importance in understanding the reactions, we have also measured the activation parameters for the conversion of two of them to the corresponding ethene complexes.

Experimental Section

General Methods and Materials. The compounds were synthesized and manipulated under argon using standard Schlenk and high vacuum techniques or in a nitrogen purged glovebox. The solvents were dried by refluxing over sodium/benzophenone and then distilling under an argon atmosphere. The deuterated solvents were purchased from Goss Scientific Instruments Ltd. They were all stored under vacuum in the presence of 4-Å molecular sieves, apart from benzene- d_6 which was dried by stirring over potassium/benzophenone and then distilled. The gases used for the matrices ($\text{Ar}, \text{N}_2, \text{CO}, \text{CH}_4$) were BOC research grade (99.999% purity). The ethene gas used in the syntheses was BOC standard grade (99.9% purity). Iridium trichloride hydrate, triphenylcarbenium tetrafluoroborate, bromoform, and triphenylphosphine were obtained from Aldrich. Hexadecane was obtained from BDH and purified by standard procedures prior to use.

The irradiation of the compounds was carried out using either an ILC Technology Ltd. ILC302UV xenon lamp equipped with a UV mirror giving an output range of 240–390 nm or a Philips HPK 125-W medium pressure mercury lamp. Excess heat was removed with a water filter; wavelengths were selected with cutoff or interference filters.

Solutions were irradiated in Pyrex ampules fitted with a ptfe stopcock or in sealed NMR tubes. Photolysis of the samples in frozen toluene- d_8 was carried out by immersing a sealed NMR tube in a partially silvered dewar containing liquid nitrogen. After photolysis the samples were thawed before obtaining the NMR spectra. Photolysis between 188 and 273 K was carried out in NMR tubes cooled by a stream of cold N_2 gas. The temperature was adjusted with a JEOL controller.

Spectroscopy. The NMR spectra were recorded with a Bruker MSL300 spectrometer unless otherwise stated. The ^1H NMR chemical shifts were referenced to residual protiated solvent as follows: benzene- d_6 $\delta = 7.13$, toluene- d_7 $\delta = 2.1$, dms o-d_6 $\delta = 2.5$, cyclohexane- d_{11} $\delta = 1.38$ ppm. The ^{13}C NMR chemical shifts

were referenced to the solvent signals as follows: toluene- d_8 $\delta = 20.1$, CDCl_3 $\delta = 77$ ppm. ^{13}C spectra were obtained with a standard DEPT pulse sequence. The ^{31}P NMR chemical shifts were referenced externally to H_3PO_4 at $\delta 0$. The spectra were recorded with broad band ^1H decoupling.

The IR spectra were recorded on either a Mattson Sirius or a Mattson Research Series FTIR spectrometer with a TGS detector and KBr beam splitter (4000–450 cm^{-1}). The spectrometer was continuously purged with CO_2 -free dry air. The spectra were recorded as the average of 128 scans with 1- cm^{-1} resolution (25K data points, 64K transform points).

The matrix isolation apparatus at York has been described in detail previously.¹² $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ was deposited on a CsI window which was cooled to 12–20 K by an Air Products CS202 closed-cycle Displex refrigerator. The complex was sublimed at 268 K onto the window from a right-angled tube concurrent with the gas stream that entered the vacuum shroud from a separate inlet.

Syntheses. $\text{CpIr}(\text{C}_2\text{H}_4)_2$ was prepared by a modification of the Mayer and Calabrese synthesis.^{5,13} $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ was prepared by the literature procedure.¹⁴ $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$ was prepared via $[\text{CpIrCl}_2]_x$, $\text{CpIrCl}_2(\text{PPh}_3)$, $\text{CpIr}(\text{CH}_3)_2(\text{PPh}_3)$, and $[\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)\text{H}]\text{PF}_6$.

$[\text{CpIrCl}_2]_x$. $\text{CpIr}(\text{C}_2\text{H}_4)_2$ (230 mg) was dissolved in thf (15 cm^3). The solution was degassed and hydrogen chloride gas admitted with vigorous stirring. The solution immediately changed color to dark orange, and a red precipitate formed. The precipitate was collected and washed with thf and hexane (225 mg, 94%). Due to the insolubility of $[\text{CpIrCl}_2]_x$ analysis by NMR was precluded. Anal. Calcd for $[\text{CpIrCl}_2]_x$: C, 18.30; H, 1.54; Cl, 21.60. Found: C, 18.73; H, 1.76; Cl, 21.19.

$\text{CpIrCl}_2(\text{PPh}_3)$. $[\text{CpIrCl}_2]_x$ (80 mg) and PPh_3 (64 mg, 2.44×10^{-4} mol) in thf (10 cm^3) were stirred at room temperature overnight. The suspension changed color from orange to yellow. The thf was removed under vacuum, and the yellow solid was crystallized from CH_2Cl_2 /hexane (134 mg, 93%). ^1H NMR (80 MHz, CDCl_3): δ 7.51 (15 H, m, $3 \times \text{C}_6\text{H}_5$), 5.38 (5 H, d, $J(\text{HP}) = 1.5$ Hz, C_5H_5). MS (peaks are quoted for ^{193}Ir and ^{36}Cl isotopomers): m/z 590 (7%, M^+), 262 (100%, PPh_3).

$\text{CpIr}(\text{CH}_3)_2(\text{PPh}_3)$. $\text{CpIrCl}_2(\text{PPh}_3)$ (250 mg, 4.22×10^{-4} mol) was suspended in diethyl ether and cooled to 0 °C. MeMgBr (diluted to 1 mol dm^{-3}) was slowly added. The yellow solid slowly changed to a yellow resinous material. After overnight stirring, a pale yellow precipitate formed. The solution was filtered and concentrated, and the $\text{CpIr}(\text{CH}_3)_2(\text{PPh}_3)$ (pale yellow solid) was extracted with hexane and purified by passing the solution through a Celite column: yield 138 mg (58%). NMR (C_7D_8): ^1H δ 7.27 (15 H, m, $3 \times \text{C}_6\text{H}_5$), 4.8 (5 H, d, $J(\text{HP}) = 1.8$ Hz, C_5H_5), 0.96 (6 H, d, $J(\text{HP}) = 5.1$ Hz, $2 \times \text{CH}_3$); $^{31}\text{P}\{^1\text{H}\}$ δ 10.46 (s). MS (peaks are quoted for ^{193}Ir isotopomers): m/z 550 (18%, M^+), 519 (100%). Accurate mass: calcd 548.1378, found 548.1389.

$[\text{CpIr}(\text{C}_2\text{H}_4)\text{H}(\text{PPh}_3)]\text{PF}_6$. $\text{CpIr}(\text{CH}_3)_2(\text{PPh}_3)$ (130 mg, 2.36×10^{-4} mol) was dissolved in CH_2Cl_2 (5 cm^3). The yellow solution was cooled to -78 °C, and $[\text{Ph}_3\text{C}]\text{PF}_6$ (90 mg, 2.3×10^{-4} mol) in CH_2Cl_2 (2 cm^3) was added in portions. The reaction mixture was allowed to warm to room temperature slowly (>1 h). Et_2O (2 cm^3) was added and the solution cooled to 0 °C. A precipitate was formed which was washed with hexane and thf to give a white solid: yield 85 mg (67%). NMR (CD_2Cl_2): ^1H (90 MHz) δ 7.39 (15 H, m, $3 \times \text{C}_6\text{H}_5$), 5.72 (5 H, d, $J(\text{HP}) = 1$ Hz, C_5H_5), 3.00 (2 H, m, C_2H_4), 2.40 (2 H, m, C_2H_4), -15.60 (1 H, d, $J(\text{HP}) = 27$ Hz, IrH); $^{13}\text{C}\{^1\text{H}\}$ δ 133.5 (d, $J(\text{CP}) = 9$ Hz), 132.5 (s), 129.5 (d, $J(\text{CP}) = 12$ Hz), 129.3 (s) ($3 \times \text{C}_6\text{H}_5$), 89.7 (s, C_5H_5), 29.3 (s, C_2H_4).

$\text{CpIr}(\text{C}_2\text{H}_4)(\text{PPh}_3)$. $[\text{CpIr}(\text{C}_2\text{H}_4)\text{H}(\text{PPh}_3)]\text{PF}_6$ (80 mg, 1.15×10^{-4} mol) was suspended in thf (10 cm^3). Excess NaH suspended in thf was added dropwise with vigorous stirring. The reaction

(12) Haddleton, D. M.; McCamley, A.; Perutz, R. N. *J. Am. Chem. Soc.* 1988, 110, 1810.

(13) Mayer, J. M.; Calabrese J. C. *Organometallics* 1984, 3, 1292.

(14) Szajek, L. P.; Lawson, R. J.; Shapley, J. R. *Organometallics* 1991, 10, 357.

(11) Kafafi, Z. H.; Hauge, R. H.; Margrave, J. L. *J. Am. Chem. Soc.* 1985, 107, 7550.


mixture was heated to 35–40 °C. Addition of NaH was continued until the evolution of gas subsided (ca. 30 min). The yellow solution was filtered and pumped to dryness. The solid was extracted with 3 × 10 cm³ of hexane and filtered. The filtrate was pumped to dryness to give a yellow solid: yield (45 mg, 71%). NMR: ¹H (dmso) δ 7.45–7.37 (15 H, m, 3 × C₆H₅), 5.05 (5 H, d, *J*(HP) = 1.1 Hz, C₆H₅), 1.84 (2 H, m, C₂H₄), 0.196 (2 H, m, C₂H₄); ¹³C{¹H} (C₇D₈) δ 138.4, 135.6, 135.4, 128.5 (3 × C₆H₅), 82.52 (C₅H₅), 7.96 (CH₂=CH₂); ³¹P{¹H} (90 MHz, C₆D₆) δ 15.9 (s). Accurate mass: calcd 546.120, found 546.1221.

CpIr(PPh₃)(C₆H₅)Br and CpIr(PPh₃)(C₂H₃)Br. CpIr-(C₂H₄)₂ (50 mg, 1.6 × 10⁻⁴ mol) was irradiated (λ > 365 nm, 84 h) in benzene (1.5 cm³) in the presence of PPh₃ (45 mg, 1.7 × 10⁻⁴ mol). The products were CpIr(PPh₃)(C₆H₅)H, CpIr(PPh₃)(C₂H₃)H, and CpIr(PPh₃)(C₂H₄). The solvent was removed and bromination of the hydrides was carried out according to the literature procedure.¹⁵ The crude mixture was dissolved in thf (10 cm³) and cooled to -78 °C. CHBr₃ (45 mg, 1.8 × 10⁻⁴ mol) diluted in thf (4 cm³) was added slowly. The mixture was left for 15 min and then allowed to warm to room temperature, and the solvent was removed. CpIr(PPh₃)(C₆H₅)Br and CpIr(PPh₃)(C₂H₃)Br were separated on a silica column using 60% toluene/40% hexane as the eluant.

CpIr(PPh₃)(C₆H₅)Br: ¹H NMR (CDCl₃) δ 7.5 (1 H, d, *J*(H_cH_m) = 11.5 Hz, *o*-phenyl hydrogen), 7.33 (15 H, m, 3 × C₆H₅), 6.81 (3 H, m, *p*- and *m*-phenyl hydrogens), 5.33 (5 H, d, *J*(HP) = 2 Hz, C₆H₅), one *o*-phenyl hydrogen signal is masked by the phosphine phenyl signals; ¹³C{¹H} NMR (CDCl₃) δ 144.1 (d, *J*(CP) = 2.8 Hz), 134.9 (d, *J*(CP) = 2.8 Hz), 128.6 (d, *J*(CP) = 2.8 Hz, 3 × C₆H₅), 132.7 (d, *J*(CP) = 3 Hz, *i*-C₆H₅), 131.0 (d, *J*(CP) = 2.4 Hz, *o*-C₆H₅), 127.7 (s, *m*-C₆H₅), 123.1 (s, *p*-C₆H₅), 86.7 (d, *J*(CP) = 3.1 Hz, C₅H₅); ³¹P{¹H} NMR (CDCl₃) δ 0.7; MS (peaks are quoted for ¹⁹³Ir, ⁸¹Br isotopomers) *m/z* 678 (5%, M⁺), 262 (100%, PPh₃). Accurate mass: calcd 678.0486, found 678.0491.

CpIr(PPh₃)(C₂H₃)Br: ¹H NMR (90 MHz, CDCl₃) (the annotation for the vinyl protons is the same as defined for the vinyl hydride complexes in the diagram below) δ 7.96 (1 H, ddd, *J*(H_cH_a) = 17 Hz, *J*(H_cH_b) = 10 Hz, *J*(H_cP) = 4.2 Hz, H_c), 7.5 (15 H, m, 3 × C₆H₅), 6.06 (1 H, ddd, *J*(H_bH_c) = 10 Hz, *J*(H_bP) = 3.2 Hz, *J*(H_bH_a) = 1.8 Hz, H_b), 5.28 (1 H, ddd, *J*(H_aH_c) = 17 Hz, *J*(H_aH_b) = 1.8 Hz, *J*(HP) = 1.8 Hz, H_a), 5.14 (5 H, d, *J*(HP) = 1.5 Hz, C₆H₅); ¹³C{¹H} NMR (CDCl₃) δ 134.9 (d, *J*(CP) = 10 Hz), 131.7 (d, *J*(CP) = 3 Hz), 128.7 (d, *J*(CP) = 11 Hz, P(C₆H₅)₃), 127.2 (d, *J*(CP) = 15 Hz, CH=CH₂), 124.7 (d, *J*(CP) = 4 Hz, CH=CH₂), 86.9 (d, *J*(CP) = 3 Hz, C₅H₅); ³¹P{¹H} NMR (CDCl₃) δ 1.0 (s); MS (peaks are quoted for ¹⁹³Ir, ⁸¹Br isotopomers) *m/z* 628 (6%, M⁺), 262 (100%, PPh₃).

NMR data for CpIr(L)(R)H (L = CO, PPh₃, R = vinyl, alkyl, aryl) are as follows. The vinyl protons are assigned according to



NMR Data for Iridium Hydride Complexes. **CpIr(CO)-(C₂H₃)H.** NMR: ¹H (C₆D₅CD₃) δ 7.51 (1 H, dd, *J*(H_cH_a) = 17.2 Hz, *J*(H_cH_b) = 9.8 Hz, H_c), 6.49 (1 H, dd, *J*(H_bH_c) = 9.8 Hz, *J*(H_bH_a) = 2.3 Hz, H_b), 5.74 (1 H, dd, *J*(H_aH_c) = 17.2 Hz, *J*(H_aH_b) = 2.3 Hz, H_a), 4.77 (5 H, s, C₅H₅), -15.57 (1 H, s, IrH); ¹³C{¹H} (C₆D₅CD₃) δ 170.4 (CO), 128.4 (CH₂), 103.4 (CH), 87.7 (C₅H₅).

CpIr(CO)(C₆D₅)H. ¹H NMR (C₆D₆): δ 4.72 (5 H, s, C₅H₅), -15.34 (1 H, s, IrH).

CpIr(CO)(C₆D₁₁)H. ¹H NMR (C₆D₁₂): δ 5.26 (5 H, s, C₅H₅), -16.64 (1 H, s, IrH).

CpIr(PPh₃)(C₂H₃)H. NMR: ¹H (C₆D₆) δ 7.85 (1 H, dddd, *J*(H_cH_a) = 17 Hz, *J*(H_cH_b) = 10 Hz, *J*(H_cP) = 7 Hz, *J*(H_cH_d) = 1 Hz, H_c), 7.6–7.0 (15 H, m, 3 × C₆H₅), 6.58 (1 H, ddd, *J*(H_bH_a) = 3 Hz, *J*(H_bH_c) = 10 Hz, *J*(H_bP) = 3 Hz, H_b), 5.79 (1 H, ddd,

J(H_aP) = 1.5 Hz, *J*(H_aH_b) = 3 Hz, *J*(H_aH_c) = 17 Hz, H_a), 4.95 (5 H, d, *J*(HP) = 1 Hz, C₅H₅), -16.05 (1 H, d, *J*(HP) = 33.1 Hz, IrH); ³¹P{¹H} (C₆D₆) δ 17.32 (s).

CpIr(PPh₃)(C₆H₅)H. NMR: ¹H (C₆D₆) δ 7.43–7.0 (15 H, m, 3 × C₆H₅), 6.8 (3 H, t, *m*- and *p*-phenyl hydrogens), 4.94 (5 H, d, *J*(HP) = 1.3 Hz, C₅H₅), -16.28 (1 H, d, *J*(HP) = 35.5 Hz, IrH), *o*-phenyl hydrogen signal masked by the phosphine phenyl signals; ³¹P{¹H} (C₆D₆) δ 15.46 (s).

Results

(a) CpIr(PPh₃)(C₂H₄). CpIr(PPh₃)(C₂H₄) was synthesized via hydride abstraction from CpIr(CH₃)₂(PPh₃) with triphenylcarbenium tetrafluoroborate, as described previously for CpIr(PMe₃)(C₂H₄) and related compounds.^{5,16}

(i) Photoproduct Studies in Solution. Photolysis of CpIr(PPh₃)(C₂H₄) has the potential of causing three different C–H activation processes: (i) insertion into the ethene ligand to form CpIr(PPh₃)(C₂H₃)H, (ii) expulsion of ethene and cyclometalation of the triphenylphosphine to form CpIr(PPh₂C₆H₄)H, (iii) expulsion of the ethene and insertion into the C–H bonds of the solvent, XH, yielding CpIr(PPh₃)(X)H. Irradiation of CpIr(PPh₃)(C₂H₄) in cold toluene solution (198 K, λ > 295 nm) resulted in conversion to a single product readily identified from the NMR spectra as CpIr(PPh₃)(C₂H₃)H. A 66% conversion was achieved after 5 h of irradiation. When photolysis of CpIr(PPh₃)(C₂H₄) was carried out in benzene at room temperature (60 min, λ > 290 nm), three products were formed in the ratio 14:2:1, of which the major product was the vinyl hydride complex, the product formed in intermediate yield was identified as CpIr(PPh₃)(Ph)H, and the minor product remains identified only as a cyclopentadienyliridium hydride. When CpIr(PPh₃)(C₂H₄) was irradiated in C₆D₆, analogous products were obtained, but the additional complication of exchange reactions was signaled by the formation of CpIr(PPh₃)(C₆D₅)H rather than CpIr(PPh₃)(C₆D₅)D. Identification of the C–H activation products was confirmed by conversion to the air-stable complexes CpIr(PPh₃)(C₆D₅)Br and CpIr(PPh₃)(C₂H₃)Br by reaction with CHBr₃.

(ii) Thermal Stability of CpIr(PPh₃)(C₂H₃)H. A sample of CpIr(PPh₃)(C₂H₃)H in dmso-*d*₆ was heated at 391 ± 2 K. The thermal isomerization of CpIr(PPh₃)(C₂H₃)H to CpIr(PPh₃)(C₂H₄) was monitored as a function of time by ¹H NMR. In each spectrum the hydride signal was normalized to the residual proton signal of the dmso. We did not use the integration of the Cp resonance because the very long relaxation time (typically 30 s) makes it necessary to use extraordinarily long delays between pulses in order to obtain accurate peak integrations.¹⁷ Typically, the relaxation times of metal hydrides are less than 1 s.¹⁸ The rate constant and Δ*G*[‡] for the isomerization were determined as (7.0 ± 1.4) × 10⁻⁶ s⁻¹ and 135 ± 3 kJ mol⁻¹, respectively, at this temperature.

(b) CpIr(CO)(C₂H₄). **(i) Photoproduct Studies in Cryogenic Solid Matrices.** A sample of CpIr(CO)(C₂H₄) in solid toluene was irradiated at 77 K (7 h, λ > 290 nm) and subsequently thawed. Room temperature ¹H and ¹³C NMR spectra demonstrated 60% conversion to the oxidative addition product CpIr(CO)(C₂H₃)H.

(16) Werner, H.; Feser, R. *J. Organomet. Chem.* **1982**, *232*, 351.

(17) Partridge, M. G. D. Phil. thesis, University of York, 1992.

(18) Hamilton, D. G.; Crabtree, R. H. *J. Am. Chem. Soc.* **1988**, *110*, 4127.

Table I. Principal IR Bands (cm^{-1}) of the Photoproducts of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in Solid Matrices

matrix type				assignment	
Ar	CH_4	2% CO 98% Ar	N_2		
2028	2015	2020	2022	$\nu(\text{CO})$	$\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$
2166	2193		2199	$\nu(\text{Ir}-\text{H})$	
1577	1570	1574	1576	$\nu(\text{CH}=\text{CH}_2)$	$\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$
1375		1379	1385	$\delta(\text{CH}_2)$	
1259		1259	1263	$\delta(\text{CH})$	$\text{CpIr}(\text{CO})_2$
994	995	995	996	$\rho(\text{CH})$	
	2005			$\nu(\text{CO})$	$\text{CpIr}(\text{CO})(\text{N}_2)$
	2193			$\nu(\text{Ir}-\text{H})$	
		2040		$\nu(\text{CO})_{\text{sym}}$	$\text{CpIr}(\text{CO})_2$
		1972		$\nu(\text{CO})_{\text{asym}}$	
			2162	$\nu(\text{N}_2)$	$\text{CpIr}(\text{CO})(\text{N}_2)$

The products from the irradiation of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in low temperature gas matrices are summarized in Table I. When $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ was deposited in an argon matrix at 20 K, it exhibited a CO-stretching band at 1987 cm^{-1} . The electronic absorption spectrum showed a band at 243 nm with a shoulder at 265 nm. After irradiation at 12 K (1 h, $\lambda > 230\text{ nm}$ and 1 h, $\lambda > 290\text{ nm}$) there was a ca. 64% depletion of the band at 1987 cm^{-1} . Product bands were observed which were assigned to $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ as follows: $2166\text{ cm}^{-1}\nu(\text{Ir}-\text{H})$, $2028\text{ cm}^{-1}\nu(\text{CO})$ (Figure 1a). In a further experiment a more highly concentrated matrix was irradiated (12 K, 2 h, $\lambda > 290\text{ nm}$). This enabled the lower intensity IR bands of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ to be observed at $1577\text{ cm}^{-1}\nu(\text{C}=\text{C}, \text{vinyl})$, $1375\text{ cm}^{-1}\delta(\text{CH}_2)$, $1259\text{ cm}^{-1}\delta(\text{CH})$, and $995\text{ cm}^{-1}\rho(\text{CH})$. These vinyl bands correspond closely to bands of other vinyl hydrides formed in matrices.⁵

Upon deposition into a methane matrix at 20 K the $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ complex exhibited a $\nu(\text{CO})$ band at 1973 cm^{-1} . UV photolysis at 12 K resulted in both intra- and intermolecular C-H activation. Bands at $2015\text{ cm}^{-1}\nu(\text{CO})$, $1569\text{ cm}^{-1}\nu(\text{C}=\text{C}, \text{vinyl})$, and $995\text{ cm}^{-1}\rho(\text{CH})$ were assigned to $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$, while the CO stretch of the methyl hydride was observed at 2005 cm^{-1} . A band at 2193 cm^{-1} was assigned to the Ir-H stretching modes of both $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$. The ratio of the products shows a pronounced wavelength dependence. Irradiation at $\lambda > 320\text{ nm}$ for 30 min resulted in a 30% depletion of the starting material and formation of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$ in a ratio of 5.7:1 (based on the absorbances of $\nu(\text{CO})$ bands). However, further photolysis of the matrix at $\lambda > 290\text{ nm}$ for 1 h resulted in 90% conversion of the starting material with $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$ formed in a ratio of 1.7:1 (Figure 1b). In a separate experiment irradiation of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in a methane matrix at $\lambda > 365\text{ nm}$ for 21 h effected 95% conversion to $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ with almost no methane activation. Further photolysis was then carried out with shorter wavelengths of light. Irradiation with $\lambda > 316\text{ nm}$ (25 min) resulted in no change in the IR spectrum, whereas the methyl hydride was formed with $\lambda > 290\text{ nm}$ (2 h). The yield of this product indicated that it could not have originated solely from the remaining $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$.

$\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ exhibited a band at 1974 cm^{-1} when it was deposited in a 98% Ar/2% CO matrix at 20 K. Irradiation ($\lambda > 320\text{ nm}$, 60 min) effected 40% depletion of the starting material and conversion to $\text{CpIr}(\text{CO})$ -




Figure 1. (a) IR spectrum showing (i) the spectrum of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in an argon matrix at 12 K and (ii) the spectrum after 1 h, $\lambda > 320\text{ nm}$ and 1 h, $\lambda > 290\text{ nm}$ showing the formation of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$. (b) (i) IR spectrum of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in a methane matrix at 12 K and (ii) the spectrum after 30 min, $\lambda > 320\text{ nm}$ and 1 h, $\lambda > 290\text{ nm}$ showing the formation of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$.

(C_2H_3)H and $\text{CpIr}(\text{CO})_2$ (Table I) in a ratio of 5:1.¹⁹ Further photolysis of the matrix at $\lambda > 290\text{ nm}$ for 60 min resulted in 60% conversion of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ to products in the ratio of $[\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}]:[\text{CpIr}(\text{CO})_2] = 1.5:1$. Again, the product ratios are based on the $\nu(\text{CO})$ band absorbances. No evidence was observed for the existence of any cyclopentadienyl "ring slip" species.

The complex was deposited in a nitrogen matrix at 20 K in order to determine the extent of CO loss from $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ upon photolysis in a matrix. Irradiation at $\lambda > 320\text{ nm}$ for 1 h followed by irradiation at $\lambda > 295\text{ nm}$ for 2 h resulted in formation of only a trace of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{N}_2)$.

(ii) Photoproduct Studies in Solution. Photolysis of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in benzene- d_6 (300 K, 1 h, $\lambda > 320\text{ nm}$) resulted in the formation of both $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{CO})(\text{C}_6\text{D}_5)\text{H}$ with ca. 60% conversion. $\text{CpIr}(\text{CO})$ -

(19) The $\nu(\text{CO})$ bands of $\text{CpIr}(\text{CO})_2$ are assigned according to: Rest, A. J.; Whitwell, I.; Graham, W. A. G.; Hoyano, J. K.; McMaster, A. D. *J. Chem. Soc., Chem. Commun.* 1984, 624.

(C₆D₅)H was observed instead of the expected CpIr(CO)-(C₆D₅)D due to a rapid H/D exchange which resulted in complete conversion to the hydride. As in the matrix isolation studies reported above, the ratio of intra- to intermolecular C-H activation exhibited a distinct wavelength dependence. When the sample was irradiated at $\lambda > 320$ nm, the products were formed in a 1:1 ratio, whereas intermolecular C-H activation was favored ([CpIr(CO)-(C₆D₅)H]:[CpIr(CO)(C₂H₃)H] = 4:1) when the sample was irradiated with $\lambda > 290$ nm (300 K, 1 h). Irradiation of CpIr(CO)(C₂H₄) in cyclohexane-*d*₁₂ (300 K, 1 h, $\lambda > 320$ nm) resulted in the formation of CpIr(CO)(C₂H₃)H and CpIr(CO)(C₆D₁₁)H in a 1:1 ratio.

Preliminary time-resolved IR experiments (carried out at Nottingham) on CpIr(CO)(C₂H₄) in heptane with 308-nm pulsed excitation show that the heptyl hydride is formed within the response time of the setup (1 μ s). This indicates that intermolecular C-H bond activation arises directly from photolysis of CpIr(CO)(C₂H₄). However, it was necessary to investigate if secondary photolysis of CpIr(CO)(C₂H₃)H was also a route to the formation of intermolecular C-H bond activation in solution. Therefore, the relative rates of formation of both vinyl and phenyl hydride upon photolysis of CpIr(CO)(C₂H₄) in benzene-*d*₆ were examined. The sample was irradiated with $\lambda > 290$ nm at 10 °C, and ¹H NMR spectra were recorded at 10-min intervals over a 1-h photolysis time. The ratio of the height of the hydride signal due to CpIr(CO)(C₂H₃)H to that of CpIr(CO)(C₆D₅)H remained constant with time within experimental error. In a further experiment CpIr(CO)(C₂H₄) was photolyzed in frozen toluene-*d*₈ (77 K, $\lambda > 290$ nm, 48 h), effecting 80% conversion to the vinyl hydride. The solution was then irradiated at 283 K ($\lambda > 290$ nm, 1 h). ¹H NMR showed that the vinyl hydride was depleted by 50% with 25% conversion to the tolyl hydrides. Thus, secondary photolysis of CpIr(CO)(C₂H₃)H is significant, but only when it is the dominant species.

(iii) **Determination of the Effect of Solvent Viscosity and Temperature on the Product Ratios.** According to the in-cage mechanism of isomerization, the viscosity of the solvent should influence the proportion of intra- to intermolecular activation. Therefore, we investigated the photolysis of CpIr(CO)(C₂H₄) in toluene-*d*₈ at three temperatures, 188, 213, and 273 K (1 h at $\lambda > 290$ nm), the lowest being only 11 deg above the freezing point. Four hydride signals were observed in the ¹H NMR spectra at δ -15.40, -15.49, -15.57, and -16.19. Previous experiments in toluene-*d*₈ showed that the high-field signal of the vinyl hydride occurs at -15.57 ppm (see Experimental Section). The other hydride signals are due to intermolecular C-D activation, followed by H/D exchange. Three different isomers are formed by reaction at different positions around the aryl ring and at the methyl group, but no specific assignment is attempted. The formation of the vinyl hydride was favored greatly by the lower temperatures and more viscous medium (Table II). At 188 K the ratio of vinyl hydride to intermolecular C-H activated products was 5.4:1, whereas at 273 K it was only 0.9:1. In order to study the effect of solvent viscosity over a narrower range of temperatures, the solvent was changed to hexadecane. CpIr(CO)(C₂H₄) was photolyzed (15 min, $\lambda > 328$ nm) in hexadecane solution at 22 °C and in a hexadecane solid matrix at 11 °C. The product ratios were monitored by IR spectroscopy. The ratio of vinyl

Table II. Temperature Dependence of the Ratio of Intramolecular to Intermolecular C-H Activation from the Photolysis ($\lambda > 290$ nm) of CpIr(CO)(C₂H₄) in Toluene-*d*₈

temp/K	product distribution		viscosity of toluene/cP ^a
	vinyl hydride	tolyl and benzyl hydrides	
77	1	0	
188	1	0.2	4.25
213	1	0.5	2.23
273	1	1.1	0.77

^a Viscosity values are determined by mathematical interpolation of data given in: *CRC Handbook of Chemistry and Physics*, 58th ed., West, R. C., Ed.; CRC Press: Boca Raton, FL, 1977.




Figure 2. First order plot of $\ln([\text{vinyl hydride}]/[\text{phenyl hydride}])$ vs time for the isomerization of CpIr(CO)(C₂H₃)H to CpIr(CO)(C₂H₄) at 284 K in benzene-*d*₆. The symbols represent the experimental points obtained by ¹H NMR. The best straight line is shown as a full line.

Table III. Rate Constants for the Thermal Isomerization of CpIr(CO)(C₂H₃)H to CpIr(CO)(C₂H₄) in Benzene-*d*₆

rate constant/s ⁻¹	temp/K
$(6.8 \pm 0.8) \times 10^{-7}$	284
$(7.5 \pm 0.8) \times 10^{-6}$	298
$(2.4 \pm 0.8) \times 10^{-5}$	307
$(5.4 \pm 0.8) \times 10^{-5}$	315
$(1.18 \pm 0.08) \times 10^{-4}$	323.5
$(2.41 \pm 0.42) \times 10^{-4}$	328
$(3.32 \pm 1.03) \times 10^{-4}$	332

hydride ($\nu(\text{CO})$ 2019 cm⁻¹):hexadecyl hydride ($\nu(\text{CO})$ 2001 cm⁻¹) in solution was 1.7:1 and in the matrix was 2.4:1.

(iv) **Thermal Stability of CpIr(CO)(C₂H₃)H and the Rate of Isomerization.** CpIr(CO)(C₂H₃)H underwent near complete conversion back to CpIr(CO)(C₂H₄) over an 8-day period at 293 K. The rate of the isomerization was monitored at a range of temperatures in benzene-*d*₆, by measuring the intensity of the CpIr(CO)(C₂H₃)H hydride resonance in the ¹H NMR as a function of time. The measurement was standardized by comparison with the intensity of the CpIr(CO)(C₆H₅)H hydride resonance, which was constant at the range of temperatures used. A first order plot for the decay of vinyl hydride, taken at 284 K, is shown in Figure 2. The rates of reaction (Table III) were used to determine the activation parameters for the thermal isomerization of CpIr(CO)(C₂H₃)H to CpIr(CO)(C₂H₄) using the Eyring equation (Figure 3a), $\Delta H^\ddagger = 95 \pm 4$ kJ mol⁻¹ and $\Delta S^\ddagger = -25 \pm 20$ J mol⁻¹ K⁻¹.

(c) **CpIr(C₂H₄)₂.** It has previously been observed that CpIr(C₂H₄)₂ was isomerized to CpIr(C₂H₄)(C₂H₃)H on irradiation with UV light in an argon matrix at 12 K.⁵




Figure 3. Eyring plots for the temperature dependence of the first order rate constant for the isomerization of (a) $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ to $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in benzene- d_6 ($\Delta H^\ddagger = 95 \pm 4 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -25 \pm 20 \text{ J mol}^{-1} \text{ K}^{-1}$) and (b) $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ to $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in toluene- d_8 ($\Delta H^\ddagger = 67 \pm 5 \text{ kJ mol}^{-1}$ and $\Delta S^\ddagger = -60 \pm 19 \text{ J mol}^{-1} \text{ K}^{-1}$).

When potential ligands such as N_2 or CO were present in the matrix, photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ resulted in the formation of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ and the substitution products $\text{CpIr}(\text{C}_2\text{H}_4)(\text{N}_2)$ and $\text{CpIr}(\text{C}_2\text{H}_4)(\text{CO})$. The vinyl hydride was itself photoactive and upon irradiation in an argon matrix underwent further C-H activation to form the vinylidene complex $\text{CpIr}(\text{CCH}_2)(\text{H})_2$. When $\text{CpIr}(\text{C}_2\text{H}_4)_2$ is irradiated in solid toluene at 77 K, only vinyl hydride formation is observed.⁵

(i) Thermal Stability of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ and the Rate of Isomerization. $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ has the lowest thermal stability of the vinyl hydrides studied here; the complex showed near complete isomerization to $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in only about 1.2 h at 273 K.

The vinyl hydride was generated by irradiation of frozen samples of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ dissolved in toluene- d_8 at 77 K. The samples were then thawed, and the rates of isomerization were measured using ^1H NMR at a range of temperatures (Table IV). The measurements were carried out in a manner similar to that used for $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$, except that the hydride resonance was standardized relative to a grease peak. In order to determine that the vinyl hydride was decaying directly to the η^2 -ethene complex, the re-formation of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ was monitored at one of the temperatures, 269 K. The rate of re-formation of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ was determined as $(3.08 \pm 0.04) \times 10^{-4} \text{ s}^{-1}$, while that for the loss of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ was $(3.19 \pm$

Table IV. Rate Constants for the Thermal Isomerization of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)(\text{H})$ to $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in Toluene- d_8

rate constant/ s^{-1}	temp/K
$(2.35 \pm 0.11) \times 10^{-5}$	247.4
$(5.03 \pm 0.28) \times 10^{-5}$	257.1
$(2.12 \pm 0.08) \times 10^{-4}$	264
$(2.59 \pm 0.14) \times 10^{-4}$	269.5
$(8.22 \pm 0.28) \times 10^{-4}$	275.7
$(9.49 \pm 0.70) \times 10^{-4}$	279.3
$(1.04 \pm 0.08) \times 10^{-3}$	280.7




Figure 4. (a) First order plot of $\ln(A_\infty - A_t)$ vs time (A is the integration ratio [ethene complex]/[grease]) for the isomerization of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ to $\text{CpIr}(\text{C}_2\text{H}_4)_2$ at 269 K in toluene- d_8 . (b) First order plot of $\ln B$ vs time (B is the integration ratio [vinyl hydride]/[grease]) for the isomerization of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ to $\text{CpIr}(\text{C}_2\text{H}_4)_2$ at 269 K in toluene- d_8 .

$0.09) \times 10^{-4} \text{ s}^{-1}$ (Figure 4). These values are the same within experimental error, confirming that $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ was isomerizing back to $\text{CpIr}(\text{C}_2\text{H}_4)_2$. An Eyring plot for the isomerization of the $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ is shown in Figure 3b. From this plot ΔH^\ddagger and ΔS^\ddagger for the isomerization are determined as $67 \pm 5 \text{ kJ mol}^{-1}$ and $-60 \pm 19 \text{ J mol}^{-1} \text{ K}^{-1}$.

(ii) Photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ and PPh_3 . Irradiation ($\lambda > 315 \text{ nm}$) of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in the presence of a slight excess of PPh_3 in CD_3CN solution caused slow release of ethene and formation of $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ as the principal product. In addition, three minor products are formed: $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$, a complex tentatively identified as $\text{CpIr}(\text{PPh}_3)_2$, and a complex formed in $<5\%$ yield which could only be characterized as an (η^5 -cyclopentadienyl)iridium hydride.²⁰ The reaction was not affected by replacement of CD_3CN by CH_3CN or by carrying out the photolysis under 1 atm of ethene.

When $\text{CpIr}(\text{C}_2\text{H}_4)_2$ was irradiated in the presence of PPh_3 in benzene solution rather than in acetonitrile, two major products were formed: $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ and another hydride together with small amounts of $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$ (preliminary results were reported earlier).² The crude reaction mixture was brominated with CHBr_3 at -78°C . Following chromatographic purification, two compounds were isolated which were readily identified as $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{Br}$ and $\text{CpIr}(\text{PPh}_3)(\text{Ph})\text{Br}$.

The photolysis reaction was examined using NMR in C_6D_6 as a function of time and photolysis wavelength (Table V). At $\lambda > 295 \text{ nm}$, the solution darkened and the proportion of phenyl hydride to vinyl hydride increased with time. Irradiation with $\lambda > 315 \text{ nm}$ resulted in less decomposition (i.e. the solution darkened much more slowly), so the reaction could be taken to higher conversion. The ratio of the two hydride products remained steady at

(20) $\text{CpIr}(\text{PPh}_3)_2$ does not appear to have been characterized previously, but its η^5 -indenyl analogue is known. Merola, J. S.; Kamacik, R. T.; Engen, D. V. *J. Am. Chem. Soc.* 1986, 108, 329.

Table V. Product Distribution of the Reaction of CpIr(C₂H₄)₂ and PPh₃ in Benzene as a Function of Photolysis Wavelength and Time

photolysis wavelength/nm	photolysis time/h	% conversion to products	ratio of CpIr(PPh ₃)(C ₆ D ₅)(H) to CpIr(PPh ₃)(C ₂ H ₃)(H)
>295	1	26	1.2
	2.5	44	1.3
	3	45	2.0
	5	59	2.2
>315	6	60	1.0
	24	65	1.0
	32	79	1.0
	55	78	1.0
	79	89	1.0
>385	4	5	0.60
	20	33	0.63
	30	45	0.63
	84	56	0.71

1:1. With $\lambda > 385$ nm the rate of conversion was inevitably reduced because of low absorption, but the ratio of phenyl hydride to vinyl hydride had now switched to 1:1.6.

A notable feature of this reaction was the occurrence of H/D exchange, which could be monitored by using the cyclopentadienyl protons as an internal standard.²¹ Thus photolysis of $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\text{C}_2\text{H}_4)_2$ with $\text{P}(\text{C}_6\text{H}_5)_3$ in C_6D_6 yielded $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\text{PPh}_3)(\text{C}_6\text{D}_5)\text{H}$ (integration of Cp: hydride = 5:1). The exchange was not suppressed by use of $\text{P}(\text{C}_6\text{D}_5)_3$, but reaction of $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\text{C}_2\text{D}_4)_2$ with $\text{P}(\text{C}_6\text{H}_5)_3$ in C_6D_6 did yield a substantial proportion of deuteride $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\text{PPh}_3)(\text{C}_6\text{D}_5)\text{D}$ (Ir(Ph)D:Ir(Ph)H 1:1). However, a similar experiment with $(\eta^5\text{-C}_5\text{H}_5)\text{Ir}(\text{C}_2\text{D}_4)_2 + \text{P}(\text{C}_6\text{H}_5)_3$ in C_6H_6 yielded no evidence for deuterium in the phenyl hydride product. Thus, hydridic protons in the phenyl hydride product are derived both from benzene and from the original coordinated ethene.

(iii) Thermal Reaction of CpIr(C₂H₄)₂ with PPh₃. A control experiment was performed to examine the thermal reaction of CpIr(C₂H₄)₂ with PPh₃. At room temperature no reaction took place. However, when CpIr(C₂H₄)₂ was heated at 150 ± 10 °C for 7 h in dms-*d*₆ in the presence of PPh₃ (70% excess), 28% of the starting material was converted to CpIr(PPh₃)(C₂H₄). After longer periods of heating, the compound tentatively identified as CpIr(PPh₃)₂ was also formed.


(iv) Thermal Reaction of CpIr(C₂H₄)(C₂H₃)H with PPh₃. A further control experiment was performed to test the lability of CpIr(C₂H₄)(C₂H₃)H. This vinyl hydride was generated as described above; after photolysis the solution was warmed to -40 °C, the seal was broken under argon, and a cold solution of PPh₃ in toluene-*d*₈ was added. The sample was resealed and warmed to room temperature. NMR revealed the presence of CpIr(C₂H₄)₂ but no other CpIr compounds, showing that CpIr(C₂H₄)(C₂H₃)H had isomerized back to starting material rather than reacting with PPh₃.

Discussion


(a) Competition between Intramolecular and Intermolecular C-H Activation. This study shows that each of the complexes CpIr(C₂H₄)(L) (L = PPh₃, CO, C₂H₄) undergoes photoinduced isomerization to the vinyl hydride

(21) Strictly, these experiments measure H/D exchange of the hydride relative to H/D exchange of the cyclopentadienyl protons. An experiment with $(\eta^5\text{-C}_5\text{D}_5)\text{Ir}(\text{C}_2\text{H}_4)_2 + \text{P}(\text{C}_6\text{H}_5)_3$ in C_6D_6 provides some evidence that such exchange may also contribute.

Scheme II. Matrix Photochemistry of CpIr(CO)(C₂H₄)



Scheme III. Solution Photochemistry of CpIr(CO)(C₂H₄)



which, in turn, is converted back to the starting ethene complex thermally (eq 1). In the first two cases, we have also observed that intermolecular C-H activation of the solvent competes with intramolecular reaction of coordinated ethene. The reactions of CpIr(C₂H₄)(CO) are summarized in Schemes II and III. It might have been expected that CpIr(C₂H₄)(PPh₃) would react to form the orthometalated CpIr(PPh₂C₆H₄)H in preference to insertion into coordinated ethene. We find that little if any orthometalation occurs; the hydride which is tentatively assigned to an orthometalation species accounts for less than 6% of the product.

The product ratios for intra- and intermolecular C-H activation of CpIr(C₂H₄)(L) (L = PPh₃, CO, C₂H₄) can be controlled by altering the nature of the surrounding media, the excitation wavelength, the temperature, and the ancillary ligand (L). In previous studies on the photoisomerization of metal ethene complexes reported from this laboratory^{2,5,6} we proposed that the reaction mechanism involves a cage-complex intermediate. We argued that the probability of vinyl hydride formation depends on the strength of the metal-ethene interaction in the cage complex and the rigidity of the matrix/solvent cage.

Our present experiments produce much evidence to indicate that the mechanism for C-H activation does involve a cage complex. The photoproduct distribution in these complexes can be altered by changing the rigidity of the surrounding medium, the excitation wavelength, and the ancillary ligand. Moreover, when CpIr(CO)(C₂H₄)


is irradiated in a methane matrix it forms $\text{CpIr}(\text{CO})\text{-(CH}_3\text{)H}$, but irradiation in an argon matrix does not yield $\text{CpIr}(\text{CO})$. Thus intermolecular C–H bond activation can take place without the complete dissociation of the ethene ligand. However, it appears that the mechanism shown in Scheme I is only a partial explanation. Of critical importance are the effects of viscosity and temperature. On the basis of the cage mechanism we had expected that increased viscosity/rigidity of the medium would shift the product distribution toward vinyl hydride. The importance of the rigidity of the surrounding medium is highlighted by the following example: $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ is only formed in appreciable amounts when photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ takes place within a rigid matrix.

In order to study the effect of the viscosity of the surrounding medium in a systematic manner $\text{CpIr}(\text{CO})\text{-(C}_2\text{H}_4)$ was irradiated under the same conditions in toluene at a range of temperatures. The results of this experiment are shown in Table II, and they clearly show that vinyl hydride formation is favored at lower temperature/more viscous solvent. In order to separate the effects of temperature and viscosity, we examined the photolysis of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ in liquid and solid hexadecane. The above results show that temperature is more important than viscosity in controlling product distribution, although both play a part. It follows that formation of intermolecular product is a thermally activated process, but it is not clear whether thermal excitation occurs in the ground or excited electronic states.²² Thermal excitation of an electronically excited state would only be feasible if that state was long-lived. Therefore, we anticipate that forthcoming TRIR experiments on $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ should enable us to distinguish between these two mechanisms.

The nature of the kinetically favored product is also determined by the wavelength of the exciting light, intermolecular C–H activation being favored at shorter wavelengths. This behavior is exhibited by $\text{CpIr}(\text{CO})\text{-(C}_2\text{H}_4)$ both in solution and in a reactive matrix and is consistent with a cage-complex intermediate. Shorter wavelength excitation will result in a vibrationally excited cage complex, from which ethene loss is more favored. This explanation of wavelength dependence is not unique: alternatively, two excited states may be involved. A systematic study of the effect of irradiation wavelength is complicated by the secondary photolysis of the vinyl hydride at shorter wavelengths ($\lambda > 290$ nm). For comparison, a study on the closely related species $\text{CpRh}(\text{CO})_2$ and $\text{Cp}^*\text{Rh}(\text{CO})_2$ showed that ligand substitution can occur through two mechanisms depending on excitation wavelength, one associative (ring-slip) and one dissociative.²³ It was proposed that the two mechanisms arose from the population of different excited states.

A change of the ancillary ligand in the metal ethene complex provides another means of controlling the kinetically favored product. This study investigates the influence of the ligands PPh_3 , CO , and C_2H_4 . When $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$ or $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ is irradiated in benzene- d_6 at room temperature, the ratios of vinyl hydride to phenyl hydride are 3.8:1 and 1:1, respectively. No quantitative measurements have been carried out on the ratio

Scheme IV. Reaction Mechanism for the Photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ and PPh_3 in CH_3CN



of products from the photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in solution because of the extensive decomposition. Even when $\text{CpIr}(\text{C}_2\text{H}_4)_2$ is irradiated at temperatures sufficiently low for the vinyl hydride to be stable, very little vinyl hydride formation is observed.⁵ At the other extreme, we have observed previously that irradiation of $\text{CpIr}(\text{PMe}_3)(\text{C}_2\text{H}_4)$ in solution resulted in the vinyl hydride accounting for greater than 90% of the products formed. Thus the ancillary ligand is important in determining the stability of the cage complex and hence the proportion of vinyl hydride. In the cases discussed, the strength of the metal-ethene interaction increases in the order $\text{C}_2\text{H}_4 < \text{CO} < \text{PPh}_3 < \text{PMe}_3$.



(b) Thermal Reversal: Vinyl Hydride to Ethene Complex. The nature of the ancillary ligand plays an important role in determining the size of the thermal barrier to the isomerization of vinyl hydride back to the η^2 -ethene complex. The complex $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ has the largest barrier to thermal reconversion, with a rate constant of $(7.0 \pm 0.5) \times 10^{-6} \text{ s}^{-1}$ at $391 \pm 2 \text{ K}$. $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ are more thermally labile and have rates of isomerization estimated to be ca. 8.2×10^{-2} and $1.2 \times 10^{-1} \text{ s}^{-1}$ at 392 K, respectively. The enthalpies of activation for the isomerization of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ are calculated to be 95 ± 3 and $67 \pm 5 \text{ kJ mol}^{-1}$, respectively. The barrier to isomerization increases with ancillary ligand in the order $\text{C}_2\text{H}_4 < \text{CO} < \text{PPh}_3$. The enthalpy of activation for the thermal isomerization of $\text{Cp}^*\text{Ir}(\text{PMe}_3)(\text{C}_2\text{H}_3)\text{H}$ to $\text{Cp}^*\text{Ir}(\text{PMe}_3)(\text{C}_2\text{H}_4)$ in cyclohexane- d_{12} is even higher: $145 \pm 5 \text{ kJ mol}^{-1}$.

(c) Reaction of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ with PPh_3 : Combined Substitution and Oxidative Addition. The major product of the photochemical reaction of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ with PPh_3 in CH_3CN is $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$. In addition, small amounts of $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$ are formed. This reaction provides a much simpler route to $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ than isomerization of $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$, since the latter requires a laborious synthesis. A control experiment shows that there is no thermal reaction of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ and PPh_3 until a high temperature is attained. Two plausible routes for the formation of the vinyl hydride may be envisaged (Scheme IV). In the first (1), substitution by PPh_3 precedes isomerization to the vinyl hydride. In the second (2), isomerization of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ precedes substitution.²⁴ We know already that $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$ is isomerized to the vinyl hydride photochemically, so the first route is fully consistent with our observations. A control exper-

(22) In an attempt to see if $\text{CpIr}(\text{CO})$ could react with methane thermally at low temperature we formed this species by photolysis of $\text{CpIr}(\text{CO})_2$ in a 6% methane/Xe matrix. However annealing the matrix up to 70 K resulted in no formation of $\text{CpIr}(\text{CO})(\text{CH}_3)\text{H}$.

(23) Drolet, D. P.; Lees, A. J. *J. Am. Chem. Soc.* 1992, 114, 4186.

(24) Since we do not observe $\text{CpIr}(\text{CH}_3\text{CN})(\text{C}_2\text{H}_4)$ (in contrast to the rhodium analogue), we do not consider it as a possible intermediate.

Scheme V. Reaction Mechanism for the Photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ and PPh_3 in Benzene**Scheme VI. Cage-Complex Reaction Mechanism Incorporating a Thermal Step to Intermolecular C-H Bond Activation**

iment demonstrates that $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ does not undergo thermal substitution by PPh_3 , so both routes require two photons. However, we have also established that $\text{CpIr}(\text{C}_2\text{H}_4)(\text{C}_2\text{H}_3)\text{H}$ is not formed in significant quantities at room temperature and is short-lived above 5 °C. We conclude, therefore, that route 1 must be dominant.

When the reaction between $\text{CpIr}(\text{C}_2\text{H}_4)_2$ and PPh_3 is performed in benzene, $\text{CpIr}(\text{PPh}_3)(\text{Ph})\text{H}$ is formed as an additional product. Again, there are two reasonable routes for its formation (Scheme V): either substitution followed by solvent oxidative addition (1) or vice versa (2). Since we observe $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_4)$, we know that route 1 can contribute. However, we have examined the photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)(\text{PPh}_3)$ and find that we obtain drastically less phenyl hydride compared to vinyl hydride than when we irradiate $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in benzene with PPh_3 . It follows that route 2 must act as the major source of $\text{CpIr}(\text{PPh}_3)(\text{Ph})\text{H}$. At present, we have not sought to establish whether the second step of route 2 is photochemically or thermally induced. When the reaction mixture is irradiated with a short wavelength cutoff ($\lambda > 295$ nm), secondary photolysis of $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ may also alter the product distri-

bution. In addition to the steps of Scheme IV and V, these reactions are complicated by H/D exchange processes involving the solvent, the coordinated ethene, and possibly the cyclopentadienyl protons. Such exchange is also evident in the photoinduced reaction of $\text{CpIr}(\text{CO})(\text{C}_2\text{H}_4)$ with C_6D_6 . Catalytic H/D exchange has been studied in detail elsewhere.²⁵ The mechanism of the process observed here has not been established.

Conclusions

(1) Intramolecular C-H activation of ethene competes with intermolecular activation of solvent in the photochemistry of $\text{CpIr}(\text{L})(\text{C}_2\text{H}_4)$ ($\text{L} = \text{PPh}_3, \text{CO}, \text{C}_2\text{H}_4$) complexes.

(2) The ancillary ligand has a major effect in determining (i) the product ratio and (ii) the kinetic stability of the vinyl hydride with respect to the ethene complexes.

(3) The product ratios are also dependent on the temperature, the rigidity of the surrounding medium, and the photolysis wavelength.

(4) The product distributions are consistent with the formation of an intermediate cage complex $[\text{CpIr}(\text{L}) \cdots \text{C}_2\text{H}_4]$. Insertion of ethene competes with displacement by solvent. The temperature dependence of the product distribution probably arises from the barrier to displacement of the ethene in the cage by the solvent (Scheme VI). This thermal step was overlooked in our previously proposed mechanism (Scheme I). However, thermal excitation of a long-lived electronically excited state has not yet been rigorously excluded. Changes in the ancillary ligand, L, favor ethene insertion if the in-cage σ -interaction is strengthened. The strength of the metal-vinyl σ bond of the product is probably influenced by a change in L in the same way.

(5) Photolysis of $\text{CpIr}(\text{C}_2\text{H}_4)_2$ in the presence of PPh_3 exploits these reactions to give a "one-pot" route to $\text{CpIr}(\text{PPh}_3)(\text{C}_2\text{H}_3)\text{H}$ and $\text{CpIr}(\text{PPh}_3)(\text{Ph})\text{H}$.

Acknowledgment. We are pleased to acknowledge the support of The Royal Society, The European Commission, NATO, SERC, and British Gas. We are grateful to Prof. M. Poliakoff and Dr. M. George for carrying out the TRIR experiments and to Prof. J. Shapley for revealing results prior to publication.

OM930116M

(25) Grebenik, P. D.; Green, M. L. H.; Izquierdo, A.; Mtetwa, V. S. B.; Prout, K. *J. Chem. Soc. Dalton Trans.* 1987, 9.