

Preparation and Reactivity of the Platinum(0) Anion [(Me₂NCS₂)Pt(PEt₃)]⁻

Daniel L. Reger* and Yan Ding


Department of Chemistry and Biochemistry, University of South Carolina, Columbia, South Carolina 29208

Received October 8, 1993*

Summary: The first monometallic anionic complex of platinum(0), [(Me₂NCS₂)Pt(PEt₃)]⁻, has been prepared from the reaction of butyllithium or lithium diisopropylamide and (Me₂NCS₂)Pt(PEt₃)H. The anion decomposes in solutions warmed above 0 °C, but its ³¹P NMR spectrum can be observed at -41 °C. By proper manipulation of the reaction conditions, the anion reacts successfully with triphenyltin chloride, methyl iodide, or iodine to yield (Me₂NCS₂)Pt(PEt₃)SnPh₃, (Me₂NCS₂)Pt(PEt₃)Me, and (Me₂NCS₂)Pt(PEt₃)I, respectively.

Introduction

We are interested in the syntheses of alkylmetal complexes of the type (Me₂NCS₂)Pt(PEt₃)(alkyl).¹ These complexes are important because they can be used to study isomerization reactions of the alkyl ligands, reactions of considerable importance in many processes catalyzed by transition metals.² For example, heating xylene solutions of either the *n*-propyl or isopropyl derivative at 120 °C for 1 week leads to the establishment of a 9/1 ratio of the *n*-propyl/isopropyl complexes (eq 1).^{3,4}



The preparations of the alkylplatinum complexes have been carried out by the reaction of (Me₂NCS₂)Pt(PEt₃)Cl and the appropriate alkylolithium or Grignard reagents.¹ An alternate route to alkylmetal complexes is the reaction of organometallic anions with alkyl halides.⁵ One method to prepare these anions is the deprotonation of metal hydride complexes.⁶ We report here the in situ generation of [(Me₂NCS₂)Pt(PEt₃)]⁻ from the stable hydride complex

* Abstract published in *Advance ACS Abstracts*, February 1, 1994.
(1) Reger, D. L.; Baxter, J. C.; Garza, D. G. *Organometallics* 1990, 9, 16.

(2) (a) Cotton, F. A.; Wilkinson, G. *Advanced Inorganic Chemistry*, 4th ed.; Interscience: New York, 1980; Chapter 30. (b) James, B. R. *Homogeneous Hydrogenation*; John Wiley and Sons: New York, 1973. (c) Tolman, C. A. In *Transition Metal Hydrides*; Muetterties, E. L., Ed.; Marcel Dekker: New York, 1971; pp 271-312. (d) Thomas, M. G.; Pretzer, W. R.; Beier, B. F.; Hirsckorn, F. J.; Muetterties, E. L. *J. Am. Chem. Soc.* 1977, 99, 743 and references therein. (e) Collman, J. P.; Hegedus, L.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Chemistry*, 2nd ed.; University Science Books: Mill Valley, CA, 1987. (f) Parshall, G. W. *Homogeneous Catalysis*; Wiley-Interscience: New York, 1980; Chapters 2-4. (g) Masters, C. *Homogeneous Transition-Metal Catalysis*; Chapman and Hall: London, 1981.

(3) Reger, D. L.; Garza, D. G.; Baxter, J. C. *Organometallics* 1990, 9, 873.

(4) Reger, D. L.; Ding, Y.; Garza, D. G.; Lebioda, L. *J. Organomet. Chem.* 1993, 452, 263.

(5) (a) Collman, J. P.; Hegedus, L.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Chemistry*, 2nd ed.; University Science Books: Mill Valley, CA, 1987; p 98. (b) King, R. B. *Acc. Chem. Res.* 1970, 3, 417.

(6) Crocco, G. L.; Gladysz, J. A. *J. Am. Chem. Soc.* 1988, 110, 6110 and references therein.

(Me₂NCS₂)Pt(PEt₃)H and its reactions with a variety of electrophiles.


Experimental Section

General Procedure. All operations were carried out under a nitrogen atmosphere using either standard Schlenk techniques or a Vacuum Atmosphere HE-493 drybox. All solvents were dried, degassed, and distilled prior to use. ¹H NMR chemical shifts are reported in ppm versus TMS. All ³¹P NMR spectra were run with proton decoupling, and ³¹P NMR chemical shifts are reported in ppm versus H₃PO₄. Elemental analyses were performed by Robertson Laboratory, Inc. All chemicals were purchased from Aldrich Chemical Co. *N,N,N',N'*-Tetramethylethylenediamine (TMEDA) and HN[CH(CH₃)₂]₂ were distilled from CaH₂. Methyl iodide was distilled from P₂O₅. Butyllithium was used as received. (Me₂NCS₂)Pt(PEt₃)H was prepared by using the published method.¹

(Dimethyldithiocarbamato)(triethylphosphine)(triphenylstannyl)platinum(II) (Me₂NCS₂)Pt(PEt₃)(SnPh₃). (Me₂NCS₂)Pt(PEt₃)H (0.080 g, 0.18 mmol) was dissolved in THF (10 mL) and cooled to -78 °C. Butyllithium (0.14 mL, 1.6 M, 0.22 mmol) was added. The reaction mixture was stirred for 2 h. A THF (10 mL) solution of Ph₃SnCl (0.11 g, 0.28 mmol) was added at -78 °C. The mixture was allowed to warm to room temperature (3 h) and stirred for an additional 2 h. The solvent was removed under vacuum. The residue was extracted with benzene (10 mL), the extract was filtered, and the benzene was evaporated. Recrystallization from toluene/hexane yielded the title compound (0.091 g, 0.12 mmol, 67%). ¹H NMR (C₆D₆): δ 8.10 (m, 6, *m*-C₆H₅); 7.31-7.19 (m, 9, *o*-, *p*-C₆H₅); 2.34, 2.11 (s, s; 3, 3; N(CH₃)₂); 1.45 (d of q, 6, PCH₂CH₃, J_{HH} = 7.6 Hz, J_{HP} = 25 Hz); 0.84 (d of t, 9, PCH₂CH₃, J_{HH} = 7.6 Hz, J_{HP} = 17 Hz). ³¹P NMR (C₆D₆): δ 7.03 (J_{PPt} = 3498 Hz, J_{PSn} = 174 Hz). Anal. Calcd for C₂₇H₃₆N₂PPtS₂Sn: C, 41.39; H, 4.63. Found: C, 41.24; H, 4.50.

(Dimethyldithiocarbamato)iodo(triethylphosphine)platinum(II) (Me₂NCS₂)Pt(PEt₃)I. (Me₂NCS₂)Pt(PEt₃)H (0.10 g, 0.23 mmol) was dissolved in THF (8 mL) and cooled to -78 °C. Lithium diisopropylamide (LDA) (0.23 mmol, freshly prepared from HN[CH(CH₃)₂]₂ and butyllithium in 5 mL of THF) was added dropwise at -78 °C. The mixture was stirred for 30 min. I₂ (0.070 g, 0.28 mmol) was added as a THF solution (10 mL, -78 °C). The mixture was warmed to room temperature (2 h) and stirred overnight. The solvent was removed under vacuum. The residue was extracted with benzene (10 mL), and the extract was filtered out and washed with dilute Na₂S₂O₃(aq) solution. The benzene was removed under vacuum, and the resulting solid recrystallized from toluene/hexane to yield a yellow solid (0.080 g, 0.14 mmol, 61%). ¹H NMR (CDCl₃): δ 3.21, 3.17 (s, s; 3, 3; N(CH₃)₂); 1.94 (d of q, 6, PCH₂CH₃, J_{HH} = 7.6 Hz, J_{HP} = 25 Hz); 1.13 (d of t, 9, PCH₂CH₃, J_{HH} = 7.6 Hz, J_{HP} = 17 Hz). ³¹P NMR (CDCl₃): δ 3.95 (J_{PPt} = 3427 Hz). Anal. Calcd for C₉H₂₁INPPtS₂: C, 19.29; H, 3.78. Found: C, 19.00; H, 3.60.

(Dimethyldithiocarbamato)methyl(triethylphosphine)platinum(II) (Me₂NCS₂)Pt(PEt₃)CH₃. (Me₂NCS₂)Pt(PEt₃)H (0.10 g, 0.23 mmol) was dissolved in THF (8 mL) and cooled to -78 °C. TMEDA (0.045 mL, 0.035 g, 0.30 mmol) and butyllithium (0.20 mL, 1.6 M, 0.32 mmol) were added. The mixture was stirred



for 30 min. Methyl iodide (0.031 mL, 0.071 g, 0.50 mmol) was added. The mixture was allowed to warm to room temperature (2 h) and stirred overnight. The solvent was removed under vacuum. The residue was extracted with benzene (10 mL), the extract was filtered, and the benzene was evaporated. Recrystallization from toluene/hexane yielded a yellow solid, shown by ^1H and ^{31}P NMR spectroscopy to be a pure sample of $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{CH}_3$ ¹ (0.072 g, 0.16 mm, 70%).

^{31}P NMR Spectrum of $[(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)]^-$. $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{H}$ (0.010 g, 0.023 mmol) was placed in a NMR tube. Approximately 1 mL of toluene- d_8 was added. The solution was cooled to -78°C . Butyllithium (0.05 mL, 1.6 M, 0.08 mmol) was added via a syringe. After 30 min, the tube was placed into an NMR spectrometer in which the probe had been cooled to -41°C and the ^{31}P NMR spectrum was recorded. The spectrum showed a broad peak at 14.99 ppm with $J_{\text{PPt}} = 2115$ Hz. This signal disappears when this sample is warmed to room temperature.

Results and Discussion

The reaction of $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{H}$ and butyllithium in THF at -78°C leads to a color change indicative of formation of the desired anion. Addition of Ph_3SnCl , the reagent established by Ellis as the best choice for trapping unstable organometallic anions,⁷ leads to the isolation of $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{SnPh}_3$ in good yield (Scheme 1). Ph_3SnCl must be added to the cooled solution of the anion. None of the desired product is isolated if the anion solution is allowed to warm before addition of Ph_3SnCl .

An analogous reaction with methyl iodide yields $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{Me}$, but also a considerable amount (15–

40%) of $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{I}$ is formed. Following the results of Crocco and Gladysz,⁸ we found that addition of TMEDA to the deprotonation reaction prevents the formation of this byproduct.

Reaction of the anion and I_2 produces $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{I}$, but this product is contaminated with $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)$ (butyl). This byproduct presumably arises from the reaction of the platinum anion with butyl iodide, formed from excess butyllithium and iodine. To circumvent this problem the hydride can be deprotonated with LDA followed by addition of the I_2 to yield exclusively $(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)\text{I}$. Thus, by proper manipulation of the reaction conditions, the anion reacts successfully with a variety of different types of electrophiles.

Although attempts to isolate $[(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)]^-$ have failed, its ^{31}P NMR spectrum could be observed at -41°C . The chemical shift of the anion is not very different from the starting hydride, but the phosphorus–platinum coupling constant drops from 3606 to 2115 Hz. In the ^1H NMR spectrum run on this sample at -41°C , the uniquely shifted hydride resonance of the starting material at $\delta -15$ has completely disappeared and no new hydride resonances are observed. The $[(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)]^-$ anion, formally a d^{10} complex, appears to be the *first monometallic anionic derivative of platinum(0) to be reported*. Nast has reported the preparation of $\text{K}_2\text{Pt}(\text{C}\equiv\text{CR})_2$, an insoluble powder that is likely to be oligomeric.⁸ It is possible that with $[(\text{Me}_2\text{NCS}_2)\text{Pt}(\text{PEt}_3)]^-$ some type of association also takes place, but it remains soluble in THF at low temperatures.

Acknowledgment is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this research. This work was also partially supported by a grant from the University of South Carolina Venture Fund. The NSF (Grant CHE-8411172 and CHE-8904942) and NIH (Grant RR-02425) have supplied funds to support NMR equipment. We thank Professor John Ellis for helpful discussions.

OM9306988

(7) Ellis, J. E.; Parnell, C. P.; Hagen, G. P. *J. Am. Chem. Soc.* 1978, 100, 3605.

(8) Nast, R.; Heinz, W. D. *Chem. Ber.* 1962, 1478.