

**Difference in Reactivity of Cyclopentadienyltitanium
Fluorides and Chlorides Using AlR_3 ($\text{R} = \text{Me}, \text{Et}$):
Syntheses and Structures of $\text{Ti(III)}-\text{F(Cl)}-\text{Al}$
Compounds $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}_2(\mu\text{-Cl})_6\text{Al}_2\text{Me}_4$,
 $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}_2(\mu\text{-F})_8\text{Al}_4\text{Me}_8$, and
 $[(\eta^5\text{-C}_5\text{H}_4\text{Me})_2\text{Ti}(\mu\text{-F})_2\text{AlEt}_2]_2^\dagger$**

Peihua Yu, Peter Müller, Musa A. Said, Herbert W. Roesky,* Isabel Usón,
Guangcai Bai, and Mathias Noltemeyer

*Institut für Anorganische Chemie der Universität Göttingen, Tammannstrasse 4,
D-37077 Göttingen, Germany*

Received November 9, 1998

The reaction of Cp^*TiX_3 [$\text{X} = \text{Cl}$ (**1a**), F (**1b**), $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$] with an excess of AlMe_3 afforded the methylated products $\text{Cp}^*\text{TiCl}_2\text{Me}$ (**2a**) and $(\text{Cp}^*\text{TiF}_2\text{Me})(\text{AlMe}_2\text{F})$ (**2b**), respectively, whereas under thermal conditions the reduced titanium species $\text{Cp}^*_2\text{Ti}_2(\mu\text{-Cl})_6\text{Al}_2\text{Me}_4$ (**3**) and $\text{Cp}^*_2\text{Ti}_2(\mu\text{-F})_8\text{Al}_4\text{Me}_8$ (**4**) were obtained in moderate yields. The reduction of Ti(IV) to Ti(III) is observed when $\text{Cp}'_2\text{TiF}_2$ ($\text{Cp}' = \eta^5\text{-C}_5\text{H}_5$, $\eta^5\text{-C}_5\text{H}_4\text{Me}$, Cp^*) and Cp^*TiF_3 are treated with AlEt_3 , leading to $[\text{Cp}'_2\text{Ti}(\mu\text{-F})_2\text{AlEt}_2]_2$ [$\text{Cp}' = \eta^5\text{-C}_5\text{H}_5$ (**5**), $\eta^5\text{-C}_5\text{H}_4\text{Me}$ (**6**), Cp^* (**7**)] and $\text{Cp}^*_2\text{Ti}_2(\mu\text{-F})_8\text{Al}_4\text{Et}_8$ (**8**), respectively. The crystal structures of complexes **3**, **4**, and **6** have been determined by X-ray single-crystal structure analyses. The core of structure **3** is described as a chair conformation. The environments of the Al and Ti atoms in **3** are distorted tetrahedral and pseudo square pyramidal, respectively. The single-crystal structure analysis of **4** shows an octahedral metal core for Al_4Ti_2 with the Ti atoms occupying trans positions. The core of complex **6** is a nonplanar eight-membered ring consisting of two titanium atoms, two aluminum atoms, and four bridging fluorine atoms.

Introduction

Metallocene complexes of titanium and zirconium (CpTiCl_3 , Cp^*TiCl_3 , and CpZrCl_3 ($\text{Cp} = \eta^5\text{-C}_5\text{H}_5$, $\text{Cp}^* = \eta^5\text{-C}_5\text{Me}_5$)), with methylaluminoxane (MAO) as cocatalyst are known as very active systems for the polymerization of olefins.¹ The investigation of the titanium system by ESR identified the presence of trivalent titanium. The reduction of titanium atoms was interpreted by the formation of a hydride species which is involved in the alkylation of titanium(IV) chloride.^{1a} Obviously MAO plays a crucial role in the formation of the active catalytic site, of which the oxidation state may exert a significant influence on the catalyst performance.² Attempts to isolate well-defined reduced titanium species from CpTiCl_3 and $\text{Cp}^*\text{TiCl}_3/\text{MAO}$ systems were unsuccessful so far.^{1a,2d} In this connection, however, it is known that cyclopentadienyltitanium chlorides can be easily reduced from Ti(IV) to Ti(III) species

using AlEt_3 .^{2a,3} For example the reaction of Cp_2TiCl_2 and $\text{Cp}'\text{TiCl}_3$ ($\text{Cp}' = \text{Cp}, \text{Cp}^*$) with AlEt_3 or Et_2AlCl resulted in the formation of compounds $\text{Cp}_2\text{Ti}(\mu\text{-Cl})_2\text{AlEtX}^{3a,b}$ and $\text{Cp}'\text{Ti}[(\mu\text{-Cl})_2\text{AlEt}_2][(\mu\text{-Cl})_2\text{AlEtX}]^{3c}$ ($\text{X} = \text{Et}$ or Cl). In fact, only a few reductive reactions of cyclopentadienyltitanium chlorides with AlMe_3 have been investigated, and very seldom well-defined products were isolated.⁴ In addition, the products of the reaction of cyclopentadienyltitanium fluorides with MAO have recently been found to exhibit a high catalytic activity for the polymerization of olefins, especially for that of styrene in a syndiotactic manner. The activity of the fluorides is improved by a factor of 30 when compared to the corresponding chlorides.^{2d,5} However, the reactions of organotitanium fluorides with alkylaluminum have attracted less attention to date. There has been a long debate about the interaction of the alkylaluminum with the transition metal center and its ligands with regard to the mechanism of this homogeneous catalysis for the polymerization of olefins.^{1d,2} Therefore, obtaining well-defined $\text{Ti(Zr)}-\text{Al}$ complexes by treating organometallics of group 4 halides with alkylaluminum has

[†] Dedicated to Professor C. N. R. Rao on the occasion of his 65th birthday.

(1) (a) Bueschges, U.; Chien, J. C. W. *J. Polym. Sci., Polym. Chem. Ed.* **1989**, *27*, 1525. (b) Chien, J. C. W.; Wang, W. P. *J. Polym. Sci., Polym. Chem. Ed.* **1987**, *26*, 3089. (c) Zambelli, A.; Olivia, L.; Pellicchia, C. *Macromolecules* **1989**, *22*, 21. (d) Brintzinger, H.-H.; Fischer, D.; Mülhaupt, R.; Rieger, B.; Waymouth, R. M. *Angew. Chem.* **1995**, *107*, 1255; *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1143.


(2) (a) Sinn, H.; Kaminsky, W. *Adv. Organomet. Chem.* **1980**, *18*, 99. (b) Kaminsky, W.; Külper, K.; Brintzinger, H.-H.; Wild, F. R. W. *P. Angew. Chem.* **1985**, *97*, 507; *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 507. (c) Möhring, P. C.; Coville, N. J. *J. Organomet. Chem.* **1994**, *479*, 1. (d) Kaminsky, W. *J. Chem. Soc., Dalton Trans.* **1998**, 1413.

(3) (a) Natta, G.; Mazzanti, G. *Tetrahedron* **1960**, *8*, 86. (b) Hay, J. N.; Obaid, R. M. S. *Eur. Polym. J.* **1978**, *14*, 965. (c) Mach, K.; Varga, V.; Antropiusová, H.; Poláček, J. *J. Organomet. Chem.* **1987**, *333*, 205.

(4) (a) Ott, K. C.; deBoer, E. J. M.; Grubbs, R. H. *Organometallics* **1984**, *3*, 223. (b) Alvanipour, A.; Atwood, J. L.; Bott, S. G.; Junk, P. C.; Kynast, U. H.; Prinz, H. *J. Chem. Soc., Dalton Trans.* **1998**, 1223.

(5) (a) Kaminsky, W.; Lenk, S.; Scholz, V.; Roesky, H. W.; Herzog, A. *Macromolecules* **1997**, *30*, 7647. (b) Murphy, E. F.; Murugavel, R.; Roesky, H. W. *Chem. Rev.* **1997**, *97*, 3425.

Scheme 1




become one of our research interests. Previously, we have shown that reaction of Cp^*MF_3 ($M = \text{Zr}, \text{Hf}$) with AlMe_3 in a 1:1 molar ratio leads to a selective exchange of fluorine atoms for methyl groups, and the byproduct AlMe_2F acts as a ligand bridging two transition metals.^{6a} Furthermore, activation of C–H bonds is observed when an excess of AlMe_3 is reacted with Cp^*MF_3 along with the formation of the fluorine-free cluster $[(\text{Cp}^*\text{M})_3\text{Al}_6\text{Me}_8(\mu_3\text{-CH})_5(\mu_2\text{-CH}_2)_2]$.⁶ The oxidation state of the metal ($M = \text{Zr}, \text{Hf}$), however, remains unchanged. The analogous reaction of the corresponding titanium halides Cp^*TiX_3 ($X = \text{Cl}, \text{F}$) with AlMe_3 , however, has not been fully studied. Herein, we describe the results of the reduction of Cp^*TiX_3 by AlMe_3 and the investigation on the reduction of Cp^*TiF_3 , $\text{Cp}^*_2\text{TiF}_2$, and $(\eta^5\text{-C}_5\text{H}_4\text{Me})_2\text{-TiF}_2$ using AlEt_3 to complete our earlier preliminary results concerning the reaction of Cp_2TiF_2 with AlEt_3 .⁷

Results and Discussion

Preparation of 3 and 4. It is known that Cp^*TiCl_3 (**1a**) can be easily methylated by using 1 equiv of AlMe_3 to yield $\text{Cp}^*\text{TiCl}_2\text{Me}$ (**2a**) in *n*-hexane.⁸ Surprisingly, reacting **1a** with 2 equiv or an excess of AlMe_3 did not yield the expected product $\text{Cp}^*\text{TiClMe}_2$ or Cp^*TiMe_3 ; instead compound **2a** in a 60% yield and a dark green crystalline product identified as $\text{Cp}^*_2\text{Ti}_2(\mu\text{-Cl})_6\text{Al}_2\text{Me}_4$ (**3**) in low yield (10%) are isolated. This suggests that further methylation of **2a** by using an excess of AlMe_3 is not possible (Scheme 1). Heating the reaction mixture resulted in improving the yield of **3** to 25%. Compound **3** was isolated from the reaction mixture at room temperature and compound **2a** at -20°C . Compound **3** is air and moisture sensitive. It decomposes without melting at 200°C . The most intense peaks in the mass spectrum of **3** are observed at m/e 508 ($M^+ - 2\text{AlMe}_2\text{Cl}$, 40%) and 253 (Cp^*TiCl_2 , 100%). The ^1H NMR spectrum of **3** was recorded at room temperature and exhibits a sharp signal (1.75 ppm) for Cp^* -ring protons and a broad signal at relatively higher field (-0.88 ppm) for Al–Me groups.

Attempts to methylate the corresponding fluoride Cp^*TiF_3 (**1b**) using AlMe_3 were unsuccessful. In fact, when **1b** was reacted with various molar ratios of AlMe_3

Scheme 2



at room temperature, a yellow oil-like product was formed after removal of the volatiles. The ^1H and ^{19}F NMR spectra show a mixture of products including $\text{Cp}^*\text{TiF}_2\text{Me}$ and AlMe_2F , which aggregate to form an adduct identified as $(\text{Cp}^*\text{TiF}_2\text{Me})(\text{AlMe}_2\text{F})$ (**2b**). When **1b** was heated under reflux with 2 equiv of AlMe_3 in toluene for 24 h, compound $\text{Cp}^*_2\text{Ti}_2(\mu\text{-F})_8\text{Al}_4\text{Me}_8$ (**4**) resulted in a 30% yield as green crystals (Scheme 2).

Compound **4** is very air and moisture sensitive. It decomposes at 230°C without melting. The most intense peak in the mass spectrum of **4** is at m/e 202 (Cp^*TiF , 100%) and the molecular ion fragment is at m/e 731 ($M^+ - \text{Me}$, 10%). The ^1H NMR spectrum of **4** at room temperature exhibits broad and unresolved signals probably due to the paramagnetic interaction between Ti(III) and ^{19}F atoms.⁹ The protons of the pentamethylcyclopentadienyl ring resonate at 2.10 ppm, and the protons of the Al–Me at relatively higher field (-0.44 to -0.62 ppm). The ^{19}F NMR spectrum of **4** is silent. The reduction of Ti(IV) to Ti(III) in compounds **3** and **4** is evidenced by the greenish color of the products. This observation is in agreement with the solid-state structure showing titanium atoms having oxidation state 3. In this connection, Atwood has recently provided an example of such behavior. Reacting CpTiCl_3 with AlMe_3 in the presence of 18-crown-6 resulted in the formation of $[\text{CpTiCl}\cdot 18\text{-crown-6}]^+[\text{AlCl}_2\text{Me}_2]^-$.^{4b} In this study the reaction of **1a** and **1b** with AlMe_3 leads to the formation of **3** and **4**, respectively, where AlMe_2X ($X = \text{Cl}, \text{F}$) is coordinating to the Ti(III) atoms through the halide atoms forming a four-membered ring in **3** and an eight-membered ring in **4** (Schemes 1 and 2). Surprisingly, it was not possible to obtain compound **4** at room temperature similar to **3**. This is probably due to the higher reduction potential of **1b** compared to **1a**.

As it has been shown so far, only monomethylation of **1a** and **1b** has been observed, yielding **2a** and **2b**, respectively. This obviously indicates that **2a** and **2b** are possible intermediates for the formation of **3** and **4**, respectively. However, the reduction mechanism (**2a** and **2b** to **3** and **4**) is still unclear. A plausible explanation might be given similar to the system $(\text{Cp}_2\text{TiCl}_2/\text{AlMe}_3$ or $\text{Cp}^*\text{TiCl}_3/\text{MAO})$ involving bridging moieties such as methene or hydrides.^{1a,4a}


Preparation of 6, 7, and 8. The reduction of Ti(IV) to Ti(III) using AlEt_3 via the β -hydrogen transfer

(6) (a) Herzog, A.; Roesky, H. W.; Zak, Z.; Noltemeyer, M. *Angew. Chem.* **1994**, *106*, 1035; *Angew. Chem., Int. Ed. Engl.* **1994**, *33*, 967. (b) Herzog, A.; Roesky, H. W.; Jäger, F.; Steiner, A.; Noltemeyer, M. *Organometallics* **1996**, *15*, 909.

(7) Yu, P.; Montero, M. L.; Barnes, C. E.; Roesky, H. W.; Usón, I. *Inorg. Chem.* **1998**, *37*, 2595.

(8) Martin, A.; Mena, M.; Pellinghelli, M. A.; Royo, P.; Serrano, R.; Tiripicchio, A. *J. Chem. Soc., Dalton Trans.* **1993**, 2117.

(9) Yu, P.; Murphy, E. F.; Roesky, H. W.; Lubini, P.; Schmidt, H.-G.; Noltemeyer, M. *Organometallics* **1997**, *16*, 313.




mechanism is demonstrable.² This explains the facile reduction of CpTiCl₃, Cp*TiCl₃, and Cp₂TiCl₂ using AlEt₃ in equimolar ratio leading to the formation of the complexes Cp'Ti(μ-Cl)₄Al₂Et₄^{3c} (Cp' = Cp, Cp*) and Cp₂-Ti(μ-Cl)₂AlEt₂, respectively.^{3a} However, the reduction of cyclopentadienyltitanium fluorides using AlEt₃ is still in the beginning stages. We have recently reported our preliminary results on the reduction of Cp₂TiF₂ with AlEt₃, which resulted in the formation of [Cp₂Ti(μ-F)₂AlEt₂]₂ (**5**).⁷ The F atoms in **5** exhibit completely different coordination modes when compared to the corresponding chlorides.^{3a,5b} The reaction of (η⁵-C₅H₄-Me)₂TiF₂, Cp*₂TiF₂, and Cp*TiF₃ with AlEt₃ in a 1:1 molar ratio at room temperature leads to the formation of blue crystals for [(η⁵-C₅H₄Me)₂Ti(μ-F)₂AlEt₂]₂ (**6**) and [Cp*₂Ti(μ-F)₂AlEt₂]₂ (**7**) (Scheme 3) and green crystals for Cp*₂Ti₂(μ-F)₈Al₄Et₈ (**8**) (Scheme 2) in moderate yields at low temperature (–20 °C). Compounds **5–8** are air and moisture sensitive. They have been characterized by MS, IR, and elemental analyses. The signals in the ¹H and ¹⁹F NMR spectra of **5–8** are broad and unresolved. Compounds **5**, **6**, and **7** decompose at 135, 120, and 105 °C, respectively. The decreasing of the decomposition point might be possibly related to the substituents on the Cp-ring. A similar observation has been made for compound **8** (190 °C) and compound **4** (230 °C), where **8** has ethyl and **4** has methyl substituents. The mass spectra of **6** and **7** indicate on the basis of their molecular ion peaks that they are monomeric in the gaseous state [EI-MS: *m/e* 300 (M⁺ – Et, 8%), 225 ((C₅H₄Me)₂TiF, 100%) for **6**, and 412 (M⁺ – Et, 10%), 337 (Cp*₂TiF, 60%), and 317 (Cp*₂Ti, 100%) for **7**]. The similar behavior of **6** and **7** in the gaseous state might suggest that compound **7** has a structure similar to **6**. The mass spectrum of **8** exhibits fragment ions [EI-MS: *m/e* 828 (M⁺ – Et, 8%), 202 (Cp*TiF, 100%)] which are similar to those observed in compound **4**.

Furthermore, when (η⁵-C₅H₄Me)₂TiF₂, Cp*₂TiF₂, and Cp*TiF₃ are allowed to react with more than 2 equiv of AlEt₃, new deep red crystals are isolated. The identification of these species is in progress. We have shown that the reaction of Cp₂TiF₂ with an excess of AlEt₃ gave a stable hydride compound [Cp(C₅H₄)Ti(μ-H)AlEt₂]₂ (**9**)⁷ in good yield (Scheme 3). This indicates that the

Table 1. Crystallographic Data for **3**, **4**, and **6**

	3	4	6
chemical formula	C ₂₄ H ₄₂ Al ₂ Cl ₆ Ti ₂	C ₂₈ H ₅₄ Al ₄ F ₈ Ti ₂	C ₃₂ H ₄₈ Al ₂ F ₄ Ti ₂
fw	693.04	746.43	658.46
cryst syst	monoclinic	monoclinic	monoclinic
space group	P2 ₁ /n (No. 14)	C2/c (No. 12)	C2/c (No. 12)
T, °C	–123	–140	–140
λ, Å	0.710 73	0.710 73	0.710 73
a, Å	9.852(2)	17.143(3)	24.340(4)
b, Å	15.099(3)	15.078(3)	7.916(1)
c, Å	11.767(2)	15.257(3)	20.128(4)
β, deg	112.09(1)	97.61(3)	121.518(2)
V, Å ³	1621.9(6)	3909.1(1)	3306(1)
Z	2	4	4
ρ _{calcd} , g cm ^{–3}	1.419	1.268	1.323
μ, mm ^{–1}	1.054	0.553	0.579
R1 ^a	0.0520	0.0355	0.0477
wR2 ^b	0.1408	0.0979	0.1178

^a R1 = Σ|F_o – F_c|/Σ|F_o| [I > 2σ(I)]. ^b wR2 = [Σw(F_o² – F_c²)²/ΣwF_o⁴]^{1/2} (all data).

**Figure 1.** Molecular structure of (η⁵-C₅Me₅)₂Ti₂(μ-Cl)₆Al₂-Me₄ (**3**).

reduction of cyclopentadienyltitanium fluorides from Ti(IV) to Ti(III) using AlEt₃ is more facile when compared to AlMe₃. In addition it is reasonable to propose that the formation of the titanium(III) hydride species occurs through a β-hydrogen transfer process. Although the reduction of Cp*TiF₃ using AlMe₃ is still ambiguous, it might occur in a similar fashion through the α-hydrogen transfer process,^{4a,10} or alternatively it might involve free radicals since the reduction is promoted under thermal conditions.

In summary, we have shown that the monoalkylation of Cp*TiX₃ (X = Cl and F) by AlMe₃ is the first step in the reaction followed by the reduction of Ti(IV) to Ti(III). The formation of the reduced chloride species **3** is more facile compared to that of the fluoride **4** due to the lower reduction potential of **1a** than that of **1b**. The resulting aggregates of Ti(III)–X–methylaluminum complexes are well-characterized. Finally the reduction of titanium fluorides is facile by involving the β-hydrogen transfer process. Some of the intermediates and the final Ti(III)–X–ethylaluminum products were isolated and characterized.

Structural Aspects. The solid-state structures of **3**, **4**, and **6** were determined by single-crystal X-ray structure analysis (Table 1). Compound **3** crystallizes as dark green crystals from *n*-hexane. The molecular structure of **3** is shown in Figure 1. Selected bond distances and angles are listed in Table 2. Compound **3** is a tetra-

Table 2. Selected Bond Distances (Å) and Angles (deg) for Compound 3



Ti(1)–Cl(1)	2.463(1)	Ti(1)–Cl(1A)	2.482(1)
Ti(1)–Cl(2)	2.554(1)	Ti(1)–Cl(3)	2.544(1)
Ti(1A)–Cl(1)	2.482(1)	Al(1)–Cl(2)	2.264(2)
Al(1)–Cl(3)	2.264(2)	Al(1)–C(6)	1.941(5)
Al(1)–C(7)	1.945(5)		
Cl(1)–Ti(1)–Cl(1A)	79.8(4)	Cl(1)–Ti(1)–Cl(3)	83.2(5)
Cl(1)–Ti(1)–Cl(2)	131.9(5)	Cl(1A)–Ti(1)–Cl(2)	80.8(4)
Cl(1A)–Ti(1)–Cl(3)	132.0(5)	Cl(3)–Ti(1)–Cl(2)	78.0(4)
Ti(1)–Cl(1)–Ti(1A)	100.2(4)	Cl(2)–Al(1)–Cl(3)	90.2(7)
Al(1)–Cl(3)–Ti(1)	96.0(6)	Al(1)–Cl(2)–Ti(1)	95.7(6)
C(6)–Al(1)–Cl(2)	111.2(2)	C(6)–Al(1)–C(7)	120.1(2)
C(6)–Al(1)–Cl(3)	111.9(2)	C(7)–Al(1)–Cl(2)	109.3(2)
C(7)–Al(1)–Cl(3)	110.0(2)		

Table 3. Selected Bond Distances (Å) and Angles (deg) for Compound 4

Ti(1)–F(1)	2.066(6)	Ti(1)–F(2)	2.052(7)
Ti(1)–F(3)	2.058(7)	Ti(1)–F(4)	2.074(6)
Al(1)–F(4)	1.759(5)	Al(2)–F(1)	1.756(6)
Al(3)–F(2)	1.748(7)	Al(2A)–F(3)	1.771(7)
F(2)–Ti(1)–F(3)	80.2(2)	F(2)–Ti(1)–F(1)	79.9(2)
F(1)–Ti(1)–F(3)	128.9(3)	F(2)–Ti(1)–F(4)	129.0(3)
F(4)–Ti(1)–F(3)	78.0(2)	F(1)–Ti(1)–F(4)	79.1(2)
F(4)–Al(1)–F(4A)	96.6(4)	F(1)–Al(2)–F(3A)	95.4(3)
F(2)–Al(3)–F(2A)	94.7(5)	F(4)–Al(1)–F(4A)	96.6(4)
Al(2)–F(1)–Ti(1)	146.7(3)	Al(3)–F(2)–Ti(1)	147.7(4)
Al(2A)–F(3)–Ti(1)	145.8(4)	Al(1)–F(4)–Ti(1)	145.9(3)

nuclear complex in which two $[\text{Cp}^*\text{TiCl}(\mu\text{-Cl})_2\text{AlMe}_2]$ moieties are bridged by the Cl atoms in an asymmetric arrangement. Compound **3** is a centrosymmetric molecule with the center of inversion in the center of the $\text{Ti}(\mu\text{-Cl})_2\text{Ti}$ plane and with the 2-fold axis going through the bridging Cl(1) and Cl(1A). The four-membered $\text{Al}(1)(\mu\text{-Cl})_2\text{Ti}(1)$, $\text{Ti}(1)(\mu\text{-Cl})_2\text{Ti}(1\text{A})$, and $\text{Al}(1\text{A})(\mu\text{-Cl})_2\text{Ti}(1\text{A})$ rings are planar with small deviations from planarity (rms deviation 0.02, 0.01, and 0.02 Å). The core of complex **3** is described as a chair conformation consisting of three four-membered rings. The Al atoms have a distorted tetrahedral coordination sphere, while the environment of the Ti atoms is pseudo square pyramidal. The average bond distance of $\text{Ti}-\text{Cl}-\text{Ti}$ ($\text{Ti}-\text{Cl}$, 2.47 Å) is slightly shorter than that of $\text{Ti}-\text{Cl}-\text{Al}$ ($\text{Ti}-\text{Cl}$, 2.54 Å). Both of them are longer than those in compound $[\text{CpTiCl-18-crown-6}][\text{AlCl}_2\text{Me}_2]$ ($\text{Ti}-\text{Cl}$, 2.34 Å).^{4b} The Al–Cl (2.26 Å) and Al–C (1.94 Å) average bond distances are comparable to those in compound $\text{Cp}_2\text{Ti}(\mu\text{-Cl})_2\text{AlEt}_2$ (Al–Cl, 2.5 Å, Al–C 1.94 Å). The average $\text{Ti}-\text{Cl}-\text{Ti}$ (100°) angle is slightly larger than that of Al–Cl–Ti (95°). While the Cl–Al–Cl (90°) angle is larger than the Cl–Ti–Cl (79°) one.

Green single crystals of compound **4** are obtained from toluene. The X-ray single-crystal structure analysis shows **4** to crystallize in the monoclinic space group $C2/c$ with half a molecule in the asymmetric unit. The other half is generated by the 2-fold axis through Al(1) and Al(3). The selected bond distances and angles are listed in Table 3. The core of structure **4** consists of a Al_4Ti_2 octahedron with the two Ti atoms occupying trans positions. The edges of the octahedron are μ -bridged by eight F atoms, which are somewhat displaced from their ideal geometrical positions; thus, a $\text{Al}_4\text{F}_8\text{Ti}_2$ cage is formed (Figure 2). This cage has a noncrystallographic 4-fold axis through both the Ti atoms and no mirror plane through the four Al atoms. A Cp^* ligand completes the coordination sphere of Ti, which is a square pyramid

**Figure 2.** Molecular structure of $(\eta^5\text{-C}_5\text{Me}_5)_2\text{Ti}_2(\mu\text{-F})_8\text{Al}_4\text{Me}_8$ (**4**).**Figure 3.** Molecular structure of $[(\eta^5\text{-C}_5\text{H}_4\text{Me})_2\text{Ti}(\mu\text{-F})_2\text{AlEt}_2]_2$ (**6**).

(Cp^*TiF_4). The tetrahedral coordination sphere of Al is completed by two methyl groups; 50% of the cage in the crystal of **4** is disordered about the $\text{Ti}(1)-\text{Ti}(1\text{A})$ pseudo axis by 45° . It is interesting to note that the C atoms of the Al–methyl groups are not directly involved in the disorder; only the Al atoms which are connected to them change and the methyl H atoms are disordered. In compound **4** the average $\text{Ti}-\text{F}$ bond distance is 2.06 Å (average Al–F 1.76 Å) and is comparable to those determined in compound **6** ($\text{Ti}-\text{F}$ 2.11 Å and Al–F 1.74 Å, average) and $[\text{Cp}_2\text{Ti}(\mu\text{-F})_2]_3\text{Al}^{11}$ ($\text{Ti}-\text{F}$ 2.09 Å, Al–F 1.81 Å, average), whereas the average $\text{F}-\text{Al}-\text{F}$ (95.5°) angle is smaller than that in **6** (103°). It is interesting to note that there are two types of $\text{F}-\text{Ti}-\text{F}$ angles in **4** (average 80.1° and 128.9°). Both are larger than that in **6** (average 77°).

Complex **6** crystallizes as blue crystals from toluene. The single-crystal X-ray structure analysis shows **6** consists of two molecules of each of the starting materials with the loss of one ethyl group from each aluminum (Figure 3). Selected bond distances and angles are listed in Table 4. The core of complex **6** is a nonplanar eight-membered ring consisting of two titanium atoms (as

(10) Tebbe, F. N.; Parshall, G. W.; Reddy, G. S. *J. Am. Chem. Soc.* **1978**, *100*, 3611.

(11) Liu, F. Q.; Gornitzka, H.; Stalke, D.; Roesky, H. W. *Angew. Chem.* **1993**, *105*, 447; *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 442.

Table 4. Selected Bond Distances (Å) and Angles (deg) for Compound 6

Ti(1)-F(1)	2.119(2)	Ti(1)-F(2)	2.113(2)
Al(1)-F(1)	1.749(2)	Al(1)-F(2A)	1.741(2)
Al(1A)-F(2)	1.741(2)	Al(1)-C(11)	1.974(4)
Al(1)-C(13)	1.965(4)		
F(2)-Ti(1)-F(1)	76.68(7)	F(2A)-Al(1)-F(1)	102.9(1)
Al(1A)-F(2)-Ti(1)	162.9(1)	Al(1)-F(1)-Ti(1)	154.0(1)
F(1)-Al(1)-C(11)	108.2(1)	F(1)-Al(1)-C(13)	108.0(1)
F(2A)-Al(1)-C(11)	106.0(1)	F(2A)-Al(1)-C(13)	110.0(1)
C(13)-Al(1)-C(11)	120.4(2)		

Cp₂Ti units), two aluminum atoms (as AlEt₂ units), and four bridging fluorine atoms, respectively. Both the Ti and Al atoms have distorted tetrahedral coordination spheres. In compound **5** four of the metal atoms and two of the fluorine atoms form a plane, whereas in compound **6** the corresponding four metal atoms and two of the fluorine atoms (F(2) and F(2A)) are far away from planarity. Instead, the Al₂F₄Ti₂ core of **6** shows a boatlike conformation (Figure 3). The Ti-Ti distance (5.865 Å) indicates that no metal-metal interaction is found in compound **6**. The Ti-F (2.11 Å) and Al-F (1.74 Å) average bond distances are comparable with those in compound **5** (Ti-F 2.09 Å, Al-F 1.73 Å, average respectively).⁷ In **6** the average angles of F-Al-F (102.8°), F-Ti-F (76.7°), and Al-F-Ti (153.9° and 162.9°) are comparable with those in **5** (F-Al-F 100.1°, F-Ti-F 78.5°, and Al-F-Ti 160.1°, 169.1°).

Experimental Section

General Data. All experimental manipulations were carried out under an atmosphere of dinitrogen rigorously excluding air and moisture using Schlenk techniques. Solvents were purified according to conventional procedures and were freshly distilled prior to use. Compounds **1a**,¹² **1b**,¹³ (η^5 -C₅H₄Me)₂TiF₂,¹⁴ and Cp*₂TiF₂¹⁴ were prepared as previously described. AlMe₃ and AlEt₃ were purchased from Aldrich. IR spectra were recorded on a Bio-Rad FTS-7 spectrometer. Mass spectra were obtained on Finnigan MAT 8230 and Varian MAT CH5 spectrometers. Elemental analyses were performed at the analytical laboratory of the Institute of Inorganic Chemistry, University of Göttingen. NMR spectra were recorded on a Bruker AM 250 spectrometer.

Synthesis of Cp*₂Ti₂(μ-Cl)₆Al₂Me₄ (3). To a suspension of Cp*TiCl₃ (0.58 g, 2 mmol) in *n*-hexane (50 mL) was added AlMe₃ (2.5 mL, 5 mmol, 2.0 M in toluene) by a syringe at room temperature. A clear greenish-yellow solution was obtained upon stirring the reaction mixture overnight at ambient temperature. After reducing the volume to about 15 mL, dark green crystals of compound **3** could be obtained in 10% (0.07 g) yield at room temperature along with an orange-yellow crystalline compound **2a** obtained in a 60% (0.32 g) yield by cooling at -10 °C. When the green solution was heated under reflux for 12 h, a brown oil-like product deposited from a brown-green solution on the bottom of the flask upon cooling the flask to room temperature. The dark green solution was transferred by a syringe and then concentrated to about 15 mL. The dark green crystals of **3** could be isolated in a 25% yield (0.17 g) at ambient temperature along with orange crystals of **2a**, which were obtained in a 40% (0.21 g) yield by cooling at -10 °C. Mp: >200 °C (dec). EI-MS *m/e* (%): 508 (M⁺ - 2AlMe₂Cl, 40), 253 (Cp*TiCl₂, 100). ¹H NMR (C₆D₆) δ

(ppm): 1.75 (30 H, Cp*), -0.88 (12 H, Al-Me). IR (Nujol) (cm⁻¹): 1278 m, 1162 m, 1054 s, 902 s, 850 s, 741 s, 665 s, 533 s. Anal. Calcd for C₂₄H₄₂Al₂Cl₆Ti₂: C, 41.5; H, 6.1; Cl, 30.7. Found: C, 42.1; H, 6.3; Cl, 30.2.

Synthesis of Cp*₂Ti₂(μ-F)₈Al₂Me₈ (4). To a solution of **1b** (0.48 g, 2 mmol) in toluene (40 mL) was added AlMe₃ (2 mL, 4 mmol, 2 M solution in toluene) dropwise using a syringe. The reaction mixture was allowed to stir at room temperature to give an orange-yellow solution, which turned to brown-yellow upon heating under reflux overnight. After filtration and partial removal of the solvent in vacuo, compound **4** was obtained in the form of green crystals in a yield of 30% (0.15 g) based on **1b**. Mp: >230 °C (dec). EI-MS *m/e* (%): 713 (M⁺ - Me, 10), 655 (M⁺ - Me - AlMe₂F, 15), 221 (Cp*TiF₂, 90), 202 (Cp*TiF, 100). IR (Nujol) (cm⁻¹): 2002 w, 1945 w, 1592 m, 1488 m, 1429 s, 1220 s, 1146 s, 1078 s, 988 s, 940 s, 755 s, 716 s, 695 s, 553 s, 530 s. Anal. Calcd for C₂₈H₅₄Al₂F₈Ti₂: C, 32.2; H, 7.2; F, 20.4. Found: C, 32.8; H, 7.3; F, 20.1.

Synthesis of [η^5 -C₅H₄Me)₂Ti(μ-F)₂AlEt₂]₂ (6). To a suspension of (η^5 -C₅H₄Me)₂TiF₂ (0.49 g, 2 mmol) in toluene (30 mL) was added an equimolar amount of AlEt₃ (2 mL, 1.0 M solution in *n*-hexane) via a syringe at room temperature, causing a slow gas evolution. The suspension dissolved and turned into a green-blue solution. The solution was further stirred for 1 h. Then the volume of the solution was reduced slowly under vacuum (15 mL) and kept at -20 °C for 1 week. Blue crystals of **6** were obtained in 37% yield (0.11 g) by filtration and washed with cold *n*-hexane (5 mL). Mp: >120 °C (dec). EI-MS *m/e* (%): 300 (M⁺ - Et, 8), 225 ((C₅H₄Me)₂TiF, 100). IR (Nujol) (cm⁻¹): 1036 m, 984 m, 950 m, 859 m, 799 s, 722 s, 649 s. Anal. Calcd for C₃₂H₄₈Al₂F₄Ti₂: C, 58.4; H, 7.3; F, 11.5. Found: C, 57.9; H, 7.3; F, 11.0.

Synthesis of [Cp*₂Ti(μ-F)₂AlEt₂]₂ (7). To a suspension of Cp*₂TiF₂ (0.71 g, 2 mmol) in *n*-hexane (30 mL) was added an equimolar amount of AlEt₃ (2 mL, 1.0 M solution in *n*-hexane) via a syringe at room temperature. The mixture was further stirred overnight. Reducing the solvent to about 10 mL and cooling the solution at -20 °C for 1 week produced air-sensitive, blue crystals of **7** in 35% yield (0.3 g). Mp: >105 °C (dec). EI-MS *m/e* (%): 412 (M⁺ - Et, 10), 337 (Cp*₂TiF, 60), and 317 (Cp*₂Ti, 100). IR (Nujol) (cm⁻¹): 1261 m, 1101 s, 1059 s, 1024 s, 799 m, 722 s, 640 s. Anal. Calcd for C₄₈H₈₀Al₂F₄Ti₂: C, 65.3; H, 9.1; F, 8.6. Found: C, 65.7; H, 9.4; F, 8.2.

Synthesis of Cp*₂Ti₂(μ-F)₈Al₂Et₈ (8). To a suspension of **1b** (0.48 g, 2 mmol) in *n*-hexane (30 mL) was added AlEt₃ (2 mL, 1.0 M solution in *n*-hexane) dropwise via a syringe at room temperature. The color changed from orange to dark red at once. The mixture was further stirred for 2 h to give a brown-yellow solution. After removal of the volatiles, the residue was dissolved in *n*-hexane (10 mL) and cooled to -20 °C to give green crystals of compound **8** in a 27% (0.15 g) yield. Mp: >190 °C (dec). EI-MS *m/e* (%): 828 (M⁺ - Et, 8), 799 (M⁺ - 2Et, 25), 221 (Cp*TiF₂, 90), 202 (Cp*TiF, 100). IR (Nujol) (cm⁻¹): 1108 m, 1098 m, 1024 m, 721 s, 660 s, 636 s, 612 s. Anal. Calcd for C₃₆H₇₀Al₂F₈Ti₂: C, 50.3; H, 8.2; F, 17.7. Found: C, 50.8; H, 8.3; F, 17.9.

X-ray Crystallography. Single crystals of compounds **3**, **4**, and **6** suitable for X-ray diffraction studies were selected and mounted on a glass fiber in a rapidly cooled perfluoropolyether.¹⁵ The diffraction data of **3** were collected on a Stoe-Siemens-AED2 four-circle diffractometer at 150(2) K, using Mo Kα radiation (λ = 0.710 73 Å), performing 2θ/ω scans. The data for **4** and **6** were collected on a Stoe-Siemens-Huber four-circle diffractometer coupled to a Siemens CCD area detector at 133-(2) K, with graphite-monochromated Mo Kα radiation (λ = 0.710 73 Å), performing φ and ω scans. The structures were solved by direct methods using the program SHELXS-97¹⁶ and refined against F² on all data by full-matrix least-squares with

(12) Hidalgo, G.; Mena, M.; Palacios, F.; Royo, P.; Serrano, R. *J. Organomet. Chem.* **1988**, *340*, 37.

(13) Herzog, A.; Liu, F.-Q.; Roesky, H. W.; Demsar, A.; Keller, K.; Noltemeyer, M.; Pauer, F. *Organometallics* **1994**, *13*, 1251.

(14) Murphy, E. F.; Yu, P.; Dietrich, S.; Roesky, H. W.; Parisini, E.; Noltemeyer, M. *J. Chem. Soc., Dalton Trans.* **1996**, 1983.

(15) Kottke, J.; Stalke, D. *J. Appl. Crystallogr.* **1993**, *26*, 615.

(16) Sheldrick, G. M. *Acta Crystallogr. Sect. A* **1990**, *46*, 467.

SHELXL-97.¹⁷ All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in the model at geometrically calculated positions and refined using a riding model. The disorder in **4** and the Cp* ring was modeled with the help of similarity restraints for 1–2 and 1–3 distances and displacement parameters as well as rigid bond restraints for anisotropic displacement parameters. The occupancies for the disordered parts were refined and eventually set at the convergence value.

Acknowledgment. This work was supported by the Göttinger Akademie der Wissenschaften, the Witto

(17) Sheldrick, G. M. *SHELX-97*; University of Göttingen, 1997.

GmbH, and the Deutsche Forschungsgemeinschaft. M.A.S. thanks the Alexander von Humboldt Foundation, Bonn, for a research fellowship.

Supporting Information Available: Listings of X-ray crystallographic data, atomic coordinates and equivalent isotropic displacement parameters for all non-hydrogen atoms, hydrogen positional and thermal parameters, anisotropic displacement parameters, and bond distances and angles for the structures of **3**, **4**, and **6**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM9809163