

Chemistry of $\text{In}(\text{C}_5\text{H}_5)_3$ and Some Heteroleptic Organoindium(III) Derivatives. Crystal and Molecular Structures of $\text{In}(\text{C}_5\text{H}_5)_3$, $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{PPh}_3$, and $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$

O. T. Beachley, Jr.,* David J. MacRae, Andrey Yu. Kovalevsky, Yuegang Zhang, and Xue Li

Department of Chemistry, University at Buffalo, The State University of New York, Buffalo, New York 14260

Received March 27, 2002

Pure $\text{In}(\text{C}_5\text{H}_5)_3$ has been prepared in high yields from InCl_3 and $\text{Li}(\text{C}_5\text{H}_5)$ in THF solution with the absence of external heating. Neat $\text{In}(\text{C}_5\text{H}_5)_3$ readily decomposes at $150\text{ }^\circ\text{C}$ to form $\text{In}(\text{C}_5\text{H}_5)_6$ and cyclopentadiene as the primary products whereas a benzene suspension decomposes under refluxing conditions. $\text{In}(\text{C}_5\text{H}_5)_3$ forms an isolable 1:1 adduct with PPh_3 but NMe_3 , THF, and Et_2O do not form stable adducts. Ligand redistribution reactions with InR_3 ($\text{R} = \text{Me}$ and CH_2CMe_3) in THF solution provided $\text{R}_2\text{In}(\text{C}_5\text{H}_5)$ and $\text{RIn}(\text{C}_5\text{H}_5)_2$. These heteroleptic organoindium(III) compounds have been isolated as analytically pure crystalline solids, but in THF solution they form equilibrium mixtures of the adducts of InR_3 , $\text{R}_2\text{In}(\text{C}_5\text{H}_5)$, $\text{RIn}(\text{C}_5\text{H}_5)_2$, and $\text{In}(\text{C}_5\text{H}_5)_3$, as appropriate. Crystals of $[\text{In}(\text{C}_5\text{H}_5)_3]_n$, $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{PPh}_3$, and $[(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)]_n$ were characterized by single-crystal X-ray structural studies.


Introduction

Our recent research has been designed to gain a fundamental understanding of the detailed nature of heteroleptic organometallic compounds of the heavier group 13 elements. The gallium compounds R_2GaCp and RGaCp_2 ($\text{R} = \text{Me}$,^{1,2} Et ,³ CH_2CMe_3 ,⁴ $\text{Cp} = \text{C}_5\text{H}_5$,^{1,2,4} $\text{C}_5\text{H}_4\text{Me}^{2,3}$) were prepared in nearly quantitative yields by ligand redistribution reaction between GaR_3 and GaCp_3 (eqs 1 and 2). Many of these heteroleptic com-



pounds were isolated as analytically pure solids that were chemically stable at room temperature,^{1–3} whereas in other cases pure single compounds could not be obtained.⁴ The analytically pure derivatives were crystalline solids. In contrast, liquids were typically unstable at room temperature because their ligands redistributed to form more symmetrical derivatives. X-ray structural studies of the crystalline solids revealed linear polymers with four-coordinate gallium atoms that were bridged by cyclopentadienide groups through their 1- and 3-positions. Even though many compounds were isolated as analytically pure crystalline solids, dissolution always produced equilibrium mixtures of multiple species (eqs 3 and 4). Cryoscopic molecular weight studies of benzene

solutions were consistent with the presence of mono-



mers. To prepare heteroleptic organoindium compounds and gain an understanding of their chemistry, we needed pure $\text{In}(\text{C}_5\text{H}_5)_3$, a necessary starting material for the preparation of these types of compounds by ligand redistribution reactions. In this paper, we describe a reliable procedure for the synthesis of $\text{In}(\text{C}_5\text{H}_5)_3$ as well as a description of its physical and chemical properties. This compound was used, in turn, for the preparation of the heteroleptic derivatives $\text{R}_2\text{In}(\text{C}_5\text{H}_5)$ and $\text{RIn}(\text{C}_5\text{H}_5)_2$ ($\text{R} = \text{Me}$, CH_2CMe_3).

Results and Discussion

The synthesis of analytically pure $\text{In}(\text{C}_5\text{H}_5)_3$ in high yields has been achieved by reacting InCl_3 with $\text{Li}(\text{C}_5\text{H}_5)$ in a 1:3.09 mol ratio in THF solution at room temperature. Since all reactants and products were soluble in THF, reaction times were kept short. Furthermore, no external heating was employed during the preparative reaction or during the isolation and purification of the product. The initial indium-containing product $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{THF}$ was readily separated from LiCl and excess $\text{Li}(\text{C}_5\text{H}_5)$ by extraction with benzene at room temperature. The THF was then removed quantitatively from the unstable adduct by vacuum distillation at room temperature. A final washing of the product with pentane produced $\text{In}(\text{C}_5\text{H}_5)_3$ as a bright yellow powder that did not melt but appeared to decompose at $\sim 160\text{--}164\text{ }^\circ\text{C}$ as shown by a color change from bright yellow to orange

(1) Beachley, O. T., Jr.; Royster, T. L.; Arhar, J. R. *J. Organomet. Chem.* **1992**, *434*, 11.

(2) Beachley, O. T., Jr.; Moss crop, M. T. *Organometallics* **2000**, *19*, 4550.

(3) Beachley, O. T., Jr.; Rosenblum, D. B.; Churchill, M. R.; Lake, C. H.; Krajkowski, L. M. *Organometallics* **1995**, *14*, 4402.

(4) Beachley, O. T., Jr.; Maloney, J. P.; Rogers, R. D. *Organometallics* **1997**, *16*, 3267.




Figure 1. Molecular geometry and labeling of atoms for [In(C₅H₅)₃]_n (50% probability ellipsoids for non-hydrogen atoms).

Table 1. Selected Bond Distances (Å) and Angles (deg) for In(C₅H₅)₃^a

(A) Indium–Carbon Distances			
In(1)–C(1)	2.229(4)	In(1)–C(11)	2.229(4)
In(1)–C(6)	2.389(4)	In(1)–C(8)#1	2.482(4)
In(1)#2–C(8)	2.482(4)		
(B) Carbon–Carbon Distances for Cyclopentadienide Rings			
C(1)–C(2)	1.453(6)	C(8)–C(9)	1.441(5)
C(1)–C(5)	1.461(6)	C(9)–C(10)	1.360(5)
C(2)–C(3)	1.366(6)	C(11)–C(15)	1.458(5)
C(3)–C(4)	1.423(6)	C(11)–C(12)	1.476(5)
C(4)–C(5)	1.358(6)	C(12)–C(13)	1.352(6)
C(6)–C(7)	1.419(6)	C(13)–C(14)	1.438(5)
C(6)–C(10)	1.442(5)	C(14)–C(15)	1.363(6)
C(7)–C(8)	1.408(5)		
(C) Angles around Indium			
C(11)–In(1)–C(1)	129.9(2)	C(11)–In(1)–C(6)	104.2(1)
C(11)–In(1)–C(8)#1	108.8(1)	C(6)–In(1)–C(8)#1	99.4(1)
C(1)–In(1)–C(8)#1	102.0(1)		

^a Symmetry transformations used to generate equivalent atoms: #1 $-x + 3/2, y + 1/2, z, z$; #2 $-x + 3/2, y - 1/2, z$.

to brown that occurred slowly. Even though the pure solid appeared to decompose at ~ 160 – 164 °C, heating of a suspension of In(C₅H₅)₃ in benzene to reflux was sufficient to induce decomposition as observed by a change in its color from bright yellow to brownish orange and a shift of the single resonance in its ¹H NMR spectrum to higher field (see later paragraph). As In(C₅H₅)₃ does not melt and ¹H NMR spectra of solutions containing the compound do not identify the presence of multiple In–C₅H₅ species due to rapid exchange of C₅H₅ substituents, the only tool for evaluating the purity of In(C₅H₅)₃ is its elemental analysis. The isolation of pure In(C₅H₅)₃ requires the use of pure reagents and paying careful attention to the details of the preparative procedure.

The structure of In(C₅H₅)₃ (Figure 1) consists of infinite polymeric chains that are located along the crystallographic *b* axis. Each indium is bonded to four cyclopentadienide groups. Two cyclopentadienide groups are monohapto or terminal whereas the other two form bridges between two indium atoms by using the carbon atoms at the 1- and 3-positions of the rings. Selected interatomic distances and angles are collected in Table 1. The structure of another polymorphic modification

of In(C₅H₅)₃ was reported previously.⁵ The space group for the earlier structure was *P*2₁2₁2₁ whereas the current structural study has identified the *Pbca* space group with a unit cell that is almost twice as large as the original. The indium atom has a distorted tetrahedral configuration. The largest C–In–C angle occurs between the two terminal cyclopentadienide groups and is 129.9(2)° for C(11)–In(1)–C(1) whereas the second largest angle occurs between the two bridging cyclopentadienide groups, C(6)–In(1)–C(8') (1.5 – *x*, *y* + 0.5, *z*) at 99.4(1)°. The terminally coordinated cyclopentadienide rings have largely nonconjugated π -systems with C(1)–C(3), C(4)–C(5), C(12)–C(13), and C(14)–C(15) being essentially double bonds. The ipso-carbon atoms C(1) and C(11) are bonded to In(1) by using almost sp³-hybridized orbitals as shown by the positions of the hydrogen atoms. These hydrogen atoms are displaced from the corresponding plane of the cyclopentadienide ring by 0.23 Å for H(1) and –0.43 Å for H(11). The bridging cyclopentadienide groups have an allyl-like conjugated π -system containing C(6)–C(7)–C(8) atoms and a double bond between C(9) and C(10). Thus, atoms C(6) and C(8) have close to sp²-hybridization. The cyclopentadienide ring with atoms C(1) through C(5) is positioned almost on top of the ring with C(11) through C(15). The angle between their planes is 17.7°. The positioning of these two rings results in a short van der Waals contact between C(4) and C(14) of 3.38 Å. The sum of the van der Waals radii is 3.42 Å.⁶ Thus, there might be some degree of π – π interaction. There are no short distances that might be indicative of interactions between the chains.

The preparation of In(C₅H₅)₃ has been described in two articles. The initial preparation⁷ utilized the reaction between InCl₃ and excess Na(C₅H₅) in diethyl ether but the yield was minimal and the elemental analysis suggested that the sample was impure as the percent carbon was low (calcd 58.11, found 56.49). The major indium-containing product, indium(I) cyclopentadienide, was isolated by sublimation when the crude reaction product was heated at 150 °C. The second synthesis⁸ of In(C₅H₅)₃ had two major modifications to the original procedure. The reagent Na(C₅H₅) was replaced with Li(C₅H₅) and the product was isolated with refluxing benzene. The percent carbon observed for this product was low also (calcd 58.11, found 57.6). It is noteworthy that neither of these papers reported a melting point, a decomposition temperature, or a solubility study of the compound in any solvent. When we attempted to prepare In(C₅H₅)₃ using the method described by Poland and Tuck,⁸ impure In(C₅H₅)₃ was obtained. Details of these experiments and the resulting products are provided with the Supporting Information.

The solubility of In(C₅H₅)₃ in benzene and its properties after heating in benzene were investigated in order to learn possible reasons for the poor elemental analyses for the products isolated by us after using the method of Poland and Tuck.⁸ Pure In(C₅H₅)₃ has minimal solubility in benzene at room temperature. Heating to 60 °C increased the solubility but decomposition also

(5) Einstein, F. W. B.; Gilbert, M. M.; Tuck, D. G. *Inorg. Chem.* **1972**, *11*, 2832.

(6) Zefirov, Y. V.; Zorkii, P. M. *Russ. Chem. Rev.* **1989**, *58*, 713.

(7) Fischer, E. O.; Hofmann, H. P. *Angew. Chem.* **1957**, *69*, 639.

(8) Poland, J. S.; Tuck, D. G. *J. Organomet. Chem.* **1972**, *42*, 307.

occurred. All of the originally soluble $\text{In}(\text{C}_5\text{H}_5)_3$ did not redissolve upon reheating. The chemical shift of the resonance for the C_5H_5 protons moved to higher field with heating and new resonances for C_5H_6 and dicyclopentadiene appeared in the spectrum and increased in intensity as the sample was heated for longer times. Thus, C_5H_6 was formed by thermal decomposition, not hydrolysis.

Pure $\text{In}(\text{C}_5\text{H}_5)_3$ decomposes readily at 150°C to form $\text{In}(\text{C}_5\text{H}_5)^{7,9}$ and C_5H_6 as the isolated products. Heating for only 15 min was sufficient to completely decompose a 2.6-mmol sample as $\sim 81\%$ of the indium was converted to $\text{In}(\text{C}_5\text{H}_5)$. These observations of the thermal decomposition of $\text{In}(\text{C}_5\text{H}_5)_3$ are consistent with our observations of heating the compound with benzene and with the observed instability of $\text{In}(\text{C}_5\text{Me}_5)_3$.⁹ Thus, the formation of $\text{In}(\text{C}_5\text{H}_5)$ from the decomposition of $\text{In}(\text{C}_5\text{H}_5)_3$ does not require $\text{Na}(\text{C}_5\text{H}_5)$.⁸

When $\text{In}(\text{C}_5\text{H}_5)_3$ and $\text{In}(\text{C}_5\text{H}_5)$ were present in a THF solution, the indium(III) species stabilized $\text{In}(\text{C}_5\text{H}_5)$. The solution remained pale yellow for more than 4 months while being held at $\sim 20^\circ\text{C}$. In contrast, a THF solution of only $\text{In}(\text{C}_5\text{H}_5)$ that was initially light yellow after warming from -196 to $\sim 20^\circ\text{C}$ became brown after 20 min and very dark brown, almost black, with a metallic mirror on the NMR tube after 45 min. The ^1H NMR spectrum of the THF solution of $\text{In}(\text{C}_5\text{H}_5)_3$ and $\text{In}(\text{C}_5\text{H}_5)$ exhibited only a single line for the C_5H_5 protons. Exchange of C_5H_5 moieties through the formation of an indium(II) species $\text{THF}\cdot(\text{C}_5\text{H}_5)_2\text{In}-\text{In}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$ with σ -bonded C_5H_5 moieties is consistent with indium chemistry¹¹ and with the hypothesis that decomposition of $\text{In}(\text{C}_5\text{H}_5)$ in THF involves a transformation from a π - to a σ -bonded species prior to cleavage of the indium-carbon bond.⁹ Thus, the change in chemical shift of the single resonance for the cyclopentadienide protons to higher field as samples of $\text{In}(\text{C}_5\text{H}_5)_3$ are heated in benzene could be due to the presence of $\text{In}(\text{C}_5\text{H}_5)$ which, in turn, is exchanging C_5H_5 ligands with $\text{In}(\text{C}_5\text{H}_5)_3$ through the formation of an indium(II) species.

Indium(III) cyclopentadienide is a good Lewis acid to phosphines but not to amines and ethers. The pale yellow adduct $\text{In}(\text{C}_5\text{H}_5)_3\cdot\text{PPh}_3$ exists as simple monomeric species in the crystalline state and in benzene solution. No dissociation of the adduct in the concentration range of 0.07 to 0.03 *m* was observed during cryoscopic molecular weight studies. The molecular geometry of the adduct as determined by an X-ray structural study and the labeling of the atoms is shown in Figure 2. The arrangement of the substituents on indium relative to those on phosphorus is shown in Figure 3. Selected interatomic distances and angles are collected in Table 2. The substituents around indium and phosphorus atoms are arranged in distorted tetrahedral environments. The indium-carbon distances of 2.256(2), 2.270(2), and 2.263(2) Å are very similar to the terminal $\text{In}-\text{C}(\text{C}_5\text{H}_5)$ distances in $\text{In}(\text{C}_5\text{H}_5)_3$ of 2.229(4) Å but shorter than the bridging $\text{In}-\text{C}(\text{C}_5\text{H}_5)$ distances of 2.389(4) and 2.482(4) Å determined in the present




Figure 2. Molecular geometry and labeling of atoms for $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{PPh}_3$ (50% probability ellipsoids for non-hydrogen atoms, hydrogen atoms are omitted for clarity).




Figure 3. Arrangement of atoms about the indium-phosphorus vector in $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{PPh}_3$.

Table 2. Selected Bond Distances (Å) and Angles (deg) for $(\text{C}_5\text{H}_5)_3\text{In}\cdot\text{PPh}_3$

(A) Indium-Phosphorus and Indium-Carbon Distances			
$\text{In}(1)-\text{P}(1)$	2.6992(4)	$\text{In}(1)-\text{C}(1)$	2.256(2)
$\text{In}(1)-\text{C}(6)$	2.270(2)	$\text{In}(1)-\text{C}(11)$	2.263(2)
(B) Phosphorus-Carbon Distances			
$\text{P}(1)-\text{C}(16)$	1.816(2)	$\text{P}(1)-\text{C}(28)$	1.821(2)
$\text{P}(1)-\text{C}(22)$	1.814(2)		
(C) Angles around Indium			
$\text{C}(1)-\text{In}(1)-\text{C}(11)$	113.68(6)	$\text{C}(1)-\text{In}(1)-\text{P}(1)$	106.22(5)
$\text{C}(1)-\text{In}(1)-\text{C}(6)$	114.09(7)	$\text{C}(11)-\text{In}(1)-\text{P}(1)$	102.53(5)
$\text{C}(11)-\text{In}(1)-\text{C}(6)$	115.02(7)	$\text{C}(6)-\text{In}(1)-\text{P}(1)$	103.57(5)
(D) Angles around Phosphorus			
$\text{C}(22)-\text{P}(1)-\text{C}(16)$	105.10(7)	$\text{C}(16)-\text{P}(1)-\text{C}(28)$	105.46(7)
$\text{C}(22)-\text{P}(1)-\text{C}(28)$	105.93(7)	$\text{C}(22)-\text{P}(1)-\text{In}(1)$	114.63(5)
$\text{C}(16)-\text{P}(1)-\text{In}(1)$	111.46(5)	$\text{C}(28)-\text{P}(1)-\text{In}(1)$	113.48(5)

study. The $\text{In}-\text{C}(\text{C}_5\text{H}_5)$ distance in the five-coordinate adduct $\text{Cl}_2(\text{C}_5\text{H}_5)\text{In}\cdot(\text{THF})_2$ ¹² is longer at 2.377(2) Å. The cyclopentadienide ligands are arranged around the indium with a propeller-like orientation in an *-sc*-conformation with respect to the phenyl substituents as seen in Figure 3. The phenyl groups are rotated with respect to each other so that the dihedral angles between their planes are -81.2° , -52.5° , and 53.1° . The torsion angles along the indium-phosphorus bond of $\text{C}(1)-\text{In}(1)-\text{P}(1)-\text{C}(22)$, $\text{C}(6)-\text{In}(1)-\text{P}(1)-\text{C}(16)$, and $\text{C}(11)-\text{In}(1)-\text{P}(1)-\text{C}(28)$ are $-25.29(7)^\circ$, $-26.56(7)^\circ$, and $-27.64(7)^\circ$, respectively. The indium-phosphorus bond

(9) Beachley, O. T., Jr.; Pazik, J. P.; Glassman, T. E.; Churchill, M. R.; Fettinger, J. C.; Blom, R. *Organometallics* **1988**, *7*, 1051.

(10) Beachley, O. T., Jr.; Blom, R.; Churchill, M. R.; Faegri, K., Jr.; Fettinger, J. C.; Pazik, J. P.; Victoriano, L. *Organometallics* **1989**, *8*, 346.

(11) Tuck, D. G. *Chem. Soc. Rev.* **1993**, 269.

(12) Visona, P.; Benetollo, F.; Rossetto, G.; Zanella, P. *Polyhedron* **1996**, *15*, 1743.

distance of 2.699(2) Å is shorter than the corresponding distances in both (Me₃In)₂·diphos of 2.755(4) Å¹³ and H₃-In·[P(C₆H₁₁)₃]₂ of 2.9869(5) Å.¹⁴ The cyclopentadienide groups have the characteristic features of simple σ-bonded monohapto rings.

The possibility of weak intermolecular interactions between some *p*-hydrogen atoms on the phenyl rings in one adduct molecule and a carbon-carbon π-bond of a cyclopentadienide ring in a neighboring molecule exists. The distance between the appropriate atoms for this possible interaction between H(27A)···π C(14')-C(15') (2.5 - *x*, 0.5 + *y*, 0.5 - *z*) is 2.62 Å, and the angle between C(27)-H(27A)···π(C(14')-C(15')) is 162°. The distance between H(32A)···C(7) (1 + *x*, *y*, *z*) is 2.836 Å. The sum of van der Waals radii for carbon and hydrogen is 2.870 Å.⁶ These intermolecular contacts might help the adduct crystallize by making 3-dimensional networks of the molecules.

The heteroleptic organoindium compounds Me₂In(C₅H₅) and (Me₃CCH₂)₂In(C₅H₅) were prepared as colorless solids in nearly quantitative yields by facile stoichiometric ligand redistribution reactions between In(C₅H₅)₃ and InR₃ (R = Me and CH₂CMe₃) in THF solution (eq 1) at room temperature. The THF was readily removed from these compounds by vacuum distillation at ~20 °C. Each product was purified by sublimation at 100–110 °C. The physical properties of Me₂In(C₅H₅) as prepared by the ligand-redistribution reaction were identical with those previously reported.¹⁵ Samples of pure Me₂In(C₅H₅) decompose at ~200 °C to an unknown dark brown material. No In(C₅H₅) was observed. An analytically pure sample of (Me₃CCH₂)₂-In(C₅H₅) also decomposed at 188–190 °C to an unknown dark brown material. No In(C₅H₅) was observed.

An X-ray structural study of (Me₃CCH₂)₂In(C₅H₅) revealed that the compound exists as a polymer with cyclopentadienide units bridging In(CH₂CMe₃)₂ moieties. The structure is fully ordered in the monoclinic cell and the *P*₂₁ space group. However, as the angle β in the unit cell is essentially 90°, twinning exists in the structure. Thus, the absolute configuration of the molecule could not be determined. The unit cell contains two asymmetric units, each of which consists of two (Me₃CCH₂)₂In(C₅H₅) monomers as depicted in Figure 4. Selected interatomic distances and angles are collected in Table 3. Each asymmetric unit contains two monomeric segments of the polymeric chain which have slightly different geometries with respect to the In-C₅H₅-In connections. Every indium atom is in contact with four carbon atoms, two from terminal CH₂CMe₃ groups and two from bridging C₅H₅ moieties, in a highly distorted tetrahedral environment. The C(CH₂Me₃)-In-C(CH₂Me₃) angles are significantly larger than the angle for an ideal trigonal-planar arrangement of atoms with values of 150.9(2)° [In(1)], 152.3(2)° [In(2)], 145.8(2)° [In(3)], and 145.6(2)° [In(4)], while the C(C₅H₅)-In-C(C₅H₅) angle is less than the ideal tetrahedral angle with values of 91.9(2)° [In(1)], 91.5(2)° [In(2)], 100.8(2)° [In(3)], and 102.1(2)° [In(4)]. The bridging indium-




Figure 4. Molecular geometry and labeling of atoms for [(Me₃CCH₂)₂In(C₅H₅)]_n (50% probability ellipsoids for non-hydrogen atoms, hydrogen atoms are omitted for clarity).

Table 3. Selected Bond Distances (Å) and Angles (deg) for (Me₃CCH₂)₂In(C₅H₅)^a

(A) Indium-Carbon Distances			
In(1)-C(1)	2.517(5)	In(1)-C(11)	2.180(5)
In(1)-C(16)	2.148(5)	In(1)-C(9)#1	2.543(5)
In(1)#2-C(9)	2.543(5)	In(2)-C(3)	2.550(5)
In(2)-C(6)	2.539(5)	In(2)-C(21)	2.160(5)
In(2)-C(26)	2.190(4)	In(3)-C(31)	2.534(4)
In(3)-C(41)	2.183(4)	In(3)-C(39)#2	2.530(5)
In(3)-C(46)	2.135(5)	In(3)#2-C(39)	2.530(5)
In(4)-C(33)	2.539(5)	In(4)-C(36)	2.519(4)
In(4)-C(51)	2.162(5)	In(4)-C(56)	2.179(4)
(B) Carbon-Carbon Distances for Cyclopentadienide Ring			
C(1)-C(2)	1.418(7)	C(1)-C(5)	1.417(7)
C(2)-C(3)	1.404(7)	C(3)-C(4)	1.419(7)
C(4)-C(5)	1.380(7)	C(6)-C(10)	1.401(6)
C(6)-C(7)	1.436(7)	C(7)-C(8)	1.388(7)
C(8)-C(9)	1.429(6)	C(9)-C(10)	1.408(6)
C(31)-C(32)	1.403(6)	C(31)-C(35)	1.426(6)
C(32)-C(33)	1.397(6)	C(33)-C(34)	1.443(6)
C(34)-C(35)	1.386(6)	C(36)-C(37)	1.442(6)
C(37)-C(38)	1.380(6)	C(38)-C(39)	1.438(7)
C(39)-C(40)	1.418(7)	C(36)-C(40)	1.411(7)
(C) Angles around Indium			
C(16)-In(1)-C(11)	150.9(2)	C(16)-In(1)-C(1)	103.8(2)
C(11)-In(1)-C(1)	100.3(2)	C(16)-In(1)-C(9)#1	106.1(2)
C(11)-In(1)-C(9)#1	89.0(2)	C(1)-In(1)-C(9)#1	91.9(5)
C(21)-In(2)-C(26)	152.3(2)	C(21)-In(2)-C(6)	102.4(2)
C(26)-In(2)-C(6)	100.1(2)	C(21)-In(2)-C(3)	105.4(2)
C(26)-In(2)-C(3)	89.9(2)	C(6)-In(2)-C(3)	91.5(2)
C(46)-In(3)-C(41)	145.8(2)	C(46)-In(3)-C(39)#2	103.1(2)
C(41)-In(3)-C(39)#2	91.8(2)	C(46)-In(3)-C(31)	107.98(16)
C(41)-In(3)-C(31)	98.9(2)	C(39)#2-In(3)-C(31)	100.8(2)
C(51)-In(4)-C(56)	145.6(2)	C(51)-In(4)-C(36)	109.0(2)
C(56)-In(4)-C(36)	98.7(2)	C(51)-In(4)-C(33)	102.1(2)
C(56)-In(4)-C(33)	91.2(2)	C(36)-In(4)-C(33)	102.1(2)

^a Symmetry transformations used to generate equivalent atoms: #1 - *x* + 1, *y*, *z*; #2 - *x* - 1, *y*, *z*.

carbon(C₅H₅) bond distances (Å) of 2.517(5) for In(1)-C(1), 2.543(5) for In(1)-C(9, #1), 2.550(5) for In(2)-C(3), 2.530(5) for In(3)-C(39, #2), 2.534(4) for In(3)-C(31), 2.539(5) for In(4)-C(33), and 2.519(4) for In(4)-C(36) are slightly longer than those in In(C₅H₅)₃ at 2.389(4) and 2.482(4) Å and in Me₂In(C₅H₅) at 2.457(12) and 2.484(12) Å.¹⁵ The In-C(CH₂CMe₃) bond distances are similar to the In-Me distances of 2.118(14) and 2.129(13) Å in Me₂In(C₅H₅)¹⁵ and the In-C(CH₂Me₃) distance of 2.12(2) Å in [In(CH₂CMe₃)Cl₂]_x.¹⁶ The In-C bond vectors are almost parallel in the asymmetric unit for indium-

(13) Bradley, D. C.; Chudzynska, H.; Faktor, M. M.; Frigo, D. M.; Hursthouse, M. B.; Hussain, B.; Smith, L. M. *Polyhedron* **1988**, *7*, 1289.

(14) Cole, M. L.; Hibbs, D. E.; Jones, C.; Smithies, N. A. *J. Chem. Soc., Dalton Trans.* **2000**, 545.


(15) Beachley, O. T., Jr.; Robirds, E. S.; Atwood, D. A.; Wei, P. *Organometallics* **1999**, *18*, 2561.

(1) and -(2) with the angles between these vectors being 177.5° and 176.6°. These vectors are slightly bent in the asymmetric unit for indium(3) and -(4) where the angles are 169.7° and 169.3°. The planes of the C₅H₅ rings are slightly tilted with respect to the C–In bonds by 12.2° and 11.3° for C1...C5, 13.6° and 11.2° for C6...C10, 6.2° and 13.5° for C31...C35, and 15.9° and 13.5° for C36...C40. The carbon–carbon distances within each C₅H₅ ring are appropriate for a delocalized allyl-type bonding between the carbon atoms at the 1, 2, and 3 positions of the ring whereas the carbon–carbon distance between the atoms at the 4 and 5 positions is shorter and representative of a double bond. Thus, each indium(III) atom can be considered to have two normal σ -bonds to the neopentyl carbon atoms and a three-centered two-electron bond across the C(C₅H₅)–In–C(C₅H₅) system.


The first examples of heteroleptic organoindium compounds that incorporate two cyclopentadienide groups MeIn(C₅H₅)₂ and (Me₃CCH₂)In(C₅H₅)₂ have been prepared as analytically pure yellow solids by ligand redistribution reactions (eq 2) in THF solutions. The preparative reactions utilized excess In(C₅H₅)₃. The equilibrium in which the ligands of RIn(C₅H₅)₂ redistribute to form R₂In(C₅H₅) and In(C₅H₅)₃ in THF solution (eq 4) must be shifted to minimize the amount of R₂In(C₅H₅), a potential impurity that has a sublimation temperature that is similar to or slightly less than that of RIn(C₅H₅)₂. Analytically pure MeIn(C₅H₅)₂ was isolated easily by sublimation at 110–120 °C. Decomposition, rather than melting, occurred at 150–160 °C but only very small amounts of In(C₅H₅) were formed. Analytically pure, yellow (Me₃CCH₂)In(C₅H₅)₂ was also isolated by sublimation at 90–95 °C but it was necessary to remove a small amount of (Me₃CCH₂)₂In(C₅H₅) that sublimed as a colorless solid prior to the sublimation of (Me₃CCH₂)In(C₅H₅)₂. The two compounds (Me₃CCH₂)₂In(C₅H₅) and (Me₃CCH₂)In(C₅H₅)₂ are readily distinguished by their color. Careful control of the temperature during sublimation is necessary for the isolation of analytically pure (Me₃CCH₂)In(C₅H₅)₂. It is of interest that (Me₃CCH₂)In(C₅H₅)₂ sublimed at a lower temperature than MeIn(C₅H₅)₂. This observation suggests that the cyclopentadienide bridge bonds that stabilize the compounds as solids appear to be weaker in (Me₃CCH₂)In(C₅H₅)₂ than in MeIn(C₅H₅)₂. X-ray structural studies of (Me₃CCH₂)In(C₅H₅)₂ and MeIn(C₅H₅)₂ were prevented by severe twinning of the crystals.

The neopentyl derivative (Me₃CCH₂)In(C₅H₅)₂ melts at 126.4–126.9 °C. If heating is immediately quenched after the compound melts, the observed melting point is reproducible. However, the melt does decompose. When a sample of pure (Me₃CCH₂)In(C₅H₅)₂ was heated at 150 °C in a sealed, evacuated tube, pale yellow crystals of In(C₅H₅) collected in the coolest part of the tube whereas colorless needles of (Me₃CCH₂)₂In(C₅H₅) formed on the warmer walls. A condensable gaseous product was identified as C₅H₆ with a trace of CMe₄. These data suggest that (Me₃CCH₂)In(C₅H₅)₂ melted and then redistributed its ligands to form initially a

mixture of (Me₃CCH₂)₂In(C₅H₅) and In(C₅H₅)₃ (eq 5).



The homoleptic derivative In(C₅H₅)₃ then decomposed to In(C₅H₅) and C₅H₆ (eq 6) whereas the (Me₃CCH₂)₂-



In(C₅H₅) sublimed without decomposition to the cooler walls of the apparatus.

Even though all of the heteroleptic organoindium derivatives of the types R₂In(C₅H₅) and RIn(C₅H₅)₂ (R = Me, CH₂CMe₃) can be isolated as pure compounds in the solid state, they dissolve in THF to form adducts that redistribute their ligands to form equilibrium mixtures of InR₃·THF, R₂In(C₅H₅)·THF, RIn(C₅H₅)₂·THF, and In(C₅H₅)₃·THF, as appropriate (eqs 3 and 4). Dissolution of all known organoindium(III) cyclopentadienide derivatives at room temperature requires the use of a Lewis base with significant donor ability. The cyclopentadienide bridges in the polymer must be broken for dissolution to occur. Thus, these compounds are exceedingly soluble in THF, a strong base, but are essentially insoluble in nonbasic solvents and in weakly basic solvents such as C₅H₁₂, C₆H₆, Et₂O, HCCl₃, and CCl₄. In contrast, the cyclopentadienide bridges in the related gallium derivatives are weaker as Me₂Ga(C₅H₄-Me)₂,² Et₂Ga(C₅H₄Me)₂,² Et₂Ga(C₅H₅)₂,³ and EtGa(C₅H₅)₂³ dissolve in benzene and form monomers in solution. The ¹H NMR spectrum of (Me₃CCH₂)₂In(C₅H₅) in THF solution at ~20 °C clearly demonstrates the occurrence of the ligand redistribution equilibria and the presence of (Me₃CCH₂)₂In(C₅H₅)·THF, (Me₃CCH₂)In(C₅H₅)₂·THF, In(CH₂CMe₃)₃·THF, and In(C₅H₅)₃·THF (eqs 3 and 4). Six resonances are observed for the methyl and methylene protons for the neopentyl groups. However, only one resonance for the cyclopentadienide protons can be seen. Cooling of the solution to –60 °C, the lowest temperature studied, did not change the spectrum. The cyclopentadienide protons for the three different species were not resolved. The resonances for the neopentyl substituents associated with each of the three compounds in the solution were identified by comparing the chemical shifts of each line with the resonances of the independent species.

Although Me₂In(C₅H₅) is isolable as an analytically pure solid, it exists in THF solution as a mixture of Me₂In(C₅H₅)·THF, MeIn(C₅H₅)₂·THF, InMe₃·THF, and In(C₅H₅)₃·THF according to NMR spectroscopy. Even though four compounds are present in THF solution, only three resonances, two for In–Me protons and one for C₅H₅ protons, are observed at ~20 °C. As the solution was cooled to –40 °C, one of the original In–Me resonances moved from –0.52 ppm to slightly higher field and split into two lines at –0.53 and –0.56 ppm. The second In–Me resonance at –1.10 ppm shifted to higher field also. The new line at –0.53 ppm is assigned to Me₂In(C₅H₅)·THF whereas the one at –0.56 ppm is for InMe₃·THF on the basis of their relative intensities. Pure InMe₃·THF dissolved in THF has a single resonance at –0.55 ppm. The second In–Me resonance that

(16) Beachley, O. T., Jr.; Spiegel, E. F.; Kopasz, J. P.; Rogers, R. D. *Organometallics* **1989**, *8*, 1915.

shifted from -1.10 to -1.64 ppm over the temperature range of 20 to -90 °C may be assigned to the In–Me group in $\text{MeIn}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$. The resonance corresponding to the C₅H₅ protons shifted from 6.04 to 5.97 ppm over the temperature range of 20 to -90 °C. Conformation of these assignments for the various In–Me resonances was obtained by preparing a THF solution of $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)$ with one small crystal of InMe_3 . The resonances for the In–Me groups occurred at -0.55 and -0.56 ppm at -90 °C whereas the resonance at higher field disappeared because the equilibrium had been shifted. These temperature-dependent NMR spectroscopic data are consistent with the occurrence of ligand redistribution equilibria (eqs 3 and 4) and with the exchange of methyl groups and cyclopentadienide groups between the different species. The resonances of the cyclopentadienide ligands for the different species could not be resolved at any temperature studied.

The monomethyl derivative $\text{MeIn}(\text{C}_5\text{H}_5)_2$ exists as a mixture of $\text{MeIn}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$, $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$, $\text{In}(\text{C}_5\text{H}_5)_3\cdot\text{THF}$, and possibly $\text{InMe}_3\cdot\text{THF}$ in THF solution. Two In–Me lines and one C₅H₅ line are observed at ~ 20 °C. Thus, the methyl groups are exchanging between the different species as are the cyclopentadienide groups. When the temperature was lowered from 20 to -80 °C, the most intense In–Me resonance shifted from -1.09 to -1.49 ppm whereas the smaller line moved from -0.37 to -0.03 ppm. The resonance corresponding to the C₅H₅ protons broadened and shifted from 5.98 to 5.85 ppm between 20 and -40 °C. At -60 °C the C₅H₅ protons exhibited lines at 6.06 and 5.72 ppm. Additional lowering of the temperature to -80 °C caused the cyclopentadienide resonances to shift to 6.02 and 5.71 ppm. The presence of an In–Me resonance associated with $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$ at a chemical shift other than at 0.53 ppm suggests that $\text{InMe}_3\cdot\text{THF}$ is present in solution. Even though no resonance for $\text{InMe}_3\cdot\text{THF}$ was observed, its presence is required by the balanced chemical equation for the ligand redistribution reaction of $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$ (eq 3).

¹H NMR spectroscopy suggests that $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2$ exists in THF solution as an equilibrium mixture of $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$, $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$, $(\text{Me}_3\text{CCH}_2)_3\text{In}\cdot\text{THF}$, and $\text{In}(\text{C}_5\text{H}_5)_3\cdot\text{THF}$ (eqs 3 and 4). The resonances for the methyl protons associated with the neopentyl groups for all three species were observed at 20 °C. However, only two of the three resonances for the methylene protons for the neopentyl groups could be observed. The line for the methylene protons for $(\text{Me}_3\text{CCH}_2)_3\text{In}\cdot\text{THF}$ was not observed, possibly because its intensity is expected to be extremely low and/or the resonance is masked by other resonances. The compound $(\text{Me}_3\text{CCH}_2)_3\text{In}\cdot\text{THF}$ will be formed by the redistribution of $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$, a product of the initial redistribution reaction (eq 3). Only minor changes for the chemical shifts for the resonances of the neopentyl methyl protons were observed as the temperature was lowered. Thus, the resonances at 0.99 , 0.83 , and 1.10 ppm for $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$, $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$, and $(\text{Me}_3\text{CCH}_2)_3\text{In}\cdot\text{THF}$, respectively, at 20 °C were observed at 1.00 , 0.84 , and 1.10 ppm, respectively, at -50 °C. In contrast, the methylene protons on the neopentyl substituents shifted more significantly as the lines for $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$

and $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$ at 0.61 and 0.37 ppm, respectively, moved to 0.54 and 0.25 ppm at -50 °C. The single resonance associated with the C₅H₅ protons appeared at 6.03 ppm at 20 °C but was broader and slightly shifted to 6.04 ppm at -30 °C. At -40 °C the resonance split into two lines at 5.95 and 6.05 ppm. The resonance at 6.05 ppm may be assigned to the C₅H₅ protons for the cyclopentadienide groups for both $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2\cdot\text{THF}$ and $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)\cdot\text{THF}$ whereas the resonance at 5.95 ppm may be assigned to $\text{In}(\text{C}_5\text{H}_5)_3\cdot\text{THF}$. The chemical shift of the resonance at 6.05 ppm remained constant to -60 °C, whereas the resonance at 5.95 assigned to $\text{In}(\text{C}_5\text{H}_5)_3\cdot\text{THF}$ shifted to 5.93 ppm at -60 °C, the lowest temperature studied.

The equilibrium constants for the ligand redistribution reactions for THF solutions of $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)$ (eq 3), $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$ (eq 3), $\text{MeIn}(\text{C}_5\text{H}_5)_2$ (eq 4), and $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2$ (eq 4) have calculated values of 3.9×10^{-3} , 3.2×10^{-3} , 0.15 , and 2.0×10^{-2} , respectively, at the normal operating temperature of the instrument. The equilibrium constants for the methyl derivatives were calculated on the basis of the integrations of the methyl resonances whereas the equilibrium constants for the neopentyl derivatives were based on the integrations for the methyl protons for the *tert*-butyl groups on the neopentyl substituents. More accurate, detailed, and complete thermodynamic data results could not be obtained for these systems as the resonances were not resolved fully.

Experimental Section

All compounds described in this investigation were sensitive to oxygen and moisture and were manipulated either under a purified argon atmosphere in a Vacuum Atmospheres drybox or by using standard vacuum line techniques. The starting materials $\text{Li}(\text{C}_5\text{H}_5)$,⁹ InMe_3 ,¹⁷ and $\text{In}(\text{CH}_2\text{CMe}_3)_3$ ¹⁶ were prepared by literature methods whereas InCl_3 was purchased from Strem Chemicals, Inc. and was used as received. Special care was taken during the preparation of $\text{Li}(\text{C}_5\text{H}_5)$ to ensure its purity.⁹ All solvents were carefully dried by using conventional procedures. Elemental analyses were performed by either E&R Microanalytical Laboratory, Parsippany, NJ, or Oneida Research Services, Whitesboro, NY. Melting points were determined with a Mel-Temp by using flame-sealed capillaries filled with argon and are uncorrected. ¹H NMR (400 MHz) spectra were recorded with a Varian Unity-Nova 400 spectrometer whereas ¹³C NMR (125.7 MHz) spectra were recorded with a Varian Unity-Nova 500 spectrometer. Proton chemical shifts are reported in δ (ppm) units and are referenced to SiMe_4 at δ 0.00 ppm and $\text{C}_6\text{D}_5\text{H}$ at δ 7.15 ppm or the proton impurities in *d*₈-THF at 1.73 and 3.58 ppm. Carbon chemical shifts are referenced to SiMe_4 at δ 0.00 and to *d*₈-THF at δ 67.40 and 25.27 ppm. All samples for NMR spectra were contained in flame-sealed NMR tubes. Deuterated solvents, benzene-*d*₆ and THF-*d*₈, were purchased from Aldrich Chemical Co. and Cambridge Isotopes, Inc., respectively, dried with P_4O_{10} , and vacuum distilled into tubes coated with sodium mirrors. The benzene-*d*₆ was purchased as 99.6% atom percent deuterated. Molecular weights were measured cryoscopically in benzene solution by using an instrument similar to that described by Shriver and Drezdon.¹⁸

Synthesis of $\text{In}(\text{C}_5\text{H}_5)_3$ in THF Solution. A three-necked Solv-Seal flask that contained 2.19 g (30.4 mmol) of $\text{Li}(\text{C}_5\text{H}_5)$

(17) Bradley, D. C.; Chudzynska, H. C.; Harding, I. S. *Inorganic Syntheses*; Cowley, A. H., Ed.; John Wiley and Sons Inc.: New York, 1997; Vol. 31, p 67.

(18) Shriver, D. F.; Drezdon, M. A. *The Manipulation of Air-Sensitive Compounds*; Wiley: New York, 1986; p 38.

and 75 mL of THF was connected to a sidearm dumper charged with 2.17 g (9.81 mmol) of InCl_3 and to a medium porosity frit. A THF solution of InCl_3 was added to the $\text{Li}(\text{C}_5\text{H}_5)/\text{THF}$ solution over a period of 1 h and then the resulting solution was stirred for 3 h at room temperature. Flask-to-flask vacuum distillation at room temperature was used to remove most of the THF. The color changed from colorless to orange/yellow. When a thick, viscous material was present, it was dynamically evacuated for an additional 2 h until a thick paste remained. Then approximately 75 mL of benzene was added by vacuum distillation to the resulting mixture of $\text{In}(\text{C}_5\text{H}_5)_3 \cdot \text{THF}$, LiCl , and excess $\text{Li}(\text{C}_5\text{H}_5)$. The resulting solution of $\text{In}(\text{C}_5\text{H}_5)_3 \cdot \text{THF}$ was separated from the insoluble material by filtration. The benzene and the remaining THF were removed by flask-to-flask vacuum distillation and then by dynamic evacuation for ~ 4 h. Finally, approximately 75 mL of pentane was added to the product and the resulting suspension was stirred for 2 h. The pentane was removed rapidly by vacuum distillation to produce a dry powder that was dynamically evacuated for an additional 4 h. The final product $\text{In}(\text{C}_5\text{H}_5)_3$ was a bright yellow powder that weighed 2.36 g (7.61 mmol, 77.6% yield based on InCl_3). **$\text{In}(\text{C}_5\text{H}_5)_3$** : Mp: color change from bright yellow to orange at ~ 160 – 164 °C dec. ^1H NMR ($\text{THF}-d_6$): δ 5.90 (s, C_5H_5). ^{13}C NMR ($\text{THF}-d_6$): δ 111.09 (s, C_5H_5). Anal. Calcd for $\text{C}_{15}\text{H}_{15}\text{In}$: C, 58.10; H, 4.88; In, 37.03. Found: C, 58.06; H, 4.87; In, 36.96. Solubility: soluble in THF; no appreciable solubility in Et_2O , C_6H_6 , C_5H_{12} , or CHCl_3 .

Thermal Decomposition of $\text{In}(\text{C}_5\text{H}_5)_3$. An evacuated break-seal tube that contained 0.803 g (2.59 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$ was placed halfway into a 150 °C oven. After ~ 1 min of heating pale yellow crystals began to collect at the cool end of the tube and a brown solid was observed in the bottom of the tube. A total of 15 min of heating was used to completely decompose the sample. The tube was opened and the volatile, condensable material was transferred to an NMR tube and identified as C_5H_6 . The pale, yellow crystalline solid was identified as $\text{In}(\text{C}_5\text{H}_5)$ (0.379 g, 2.11 mmol, 81.3% yield based on $\text{In}(\text{C}_5\text{H}_5)_3$). The unidentified brown residue that was insoluble in common organic solvents weighed 0.392 g. **Pale yellow crystals:** $\text{In}(\text{C}_5\text{H}_5)$. Mp 168.6–169.7 °C dec (lit.⁹ mp 169.3–171 °C dec). ^1H NMR ($\text{THF}-d_6$): δ 5.99 (s, C_5H_5). **Volatile liquid (C_5H_6 and $\text{C}_{10}\text{H}_{12}$):** ^1H NMR (C_6D_6): δ 6.49 (m, 27.99 H, C_5H_6), 6.29 (m, 29.37 H, C_5H_6), 5.93 (m, 1.01 H, C_5H_6), 5.48 (m, 0.97 H, C_5H_6), 3.57 (m, 0.73 H, $\text{C}_{10}\text{H}_{12}$), 3.12 (m, 0.82 H, $\text{C}_{10}\text{H}_{12}$), 2.70 (m, 34.27 H, $\text{C}_{10}\text{H}_{12}$), 2.59 (m, 1.02 H, $\text{C}_{10}\text{H}_{12}$), 2.11 (m, 1.06 H, $\text{C}_{10}\text{H}_{12}$), 1.58 (m, 1.76 H, $\text{C}_{10}\text{H}_{12}$), 1.18 (s, 0.57 H, $\text{C}_{10}\text{H}_{12}$), 1.16 (s, 0.43 H, $\text{C}_{10}\text{H}_{12}$).

Reaction of $\text{In}(\text{C}_5\text{H}_5)_3$ with $\text{In}(\text{C}_5\text{H}_5)$. A sidearm dumper was charged with 0.15 g (0.48 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$ and connected to a reaction vessel charged with 0.086 g (0.48 mmol) of $\text{In}(\text{C}_5\text{H}_5)$ and a magnetic stir bar. Solvent (1 mL of $\text{THF}-d_6$) was added by vacuum distillation, the reactants were mixed, and the resulting solution was poured into the NMR tube. The NMR tube was flame-sealed and maintained at -196 °C until the ^1H NMR spectrum was recorded. Solvent ($\text{THF}-d_6$) was vacuum distilled into a second NMR tube that contained 0.126 g (0.700 mmol) of $\text{In}(\text{C}_5\text{H}_5)$. This NMR tube was flame-sealed and maintained at -196 °C until the ^1H NMR spectrum was recorded. **$\text{In}(\text{C}_5\text{H}_5)_3$ and $\text{In}(\text{C}_5\text{H}_5)$** : ^1H NMR ($\text{THF}-d_6$): initial spectrum, 20 min after warming, pale yellow solution, δ 5.940 (s, C_5H_5), 4 months after solution preparation, pale yellow solution, δ 5.934 (s, C_5H_5). ^{13}C NMR ($\text{THF}-d_6$): 4 months after solution preparation, δ 108.07 (s, C_5H_5). **$\text{In}(\text{C}_5\text{H}_5)$** : ^1H NMR ($\text{THF}-d_6$): initial spectrum, 10 min after warming, brown solution, δ 5.985 (s, C_5H_5), C_5H_6 negligible; 20 h after warming, black precipitate, δ 6.500 (m, C_5H_6 , 0.21 H), 6.407 (m, C_5H_6 , 0.23 H), 5.986 (s, C_5H_5 , 5.00 H), 2.935 (s, C_5H_6 , 0.35 H); 72 h after warming, black precipitate, δ 6.503 (m, C_5H_6 , 0.20 H), 6.403 (m, C_5H_6 , 0.23 H), 5.986 (s, C_5H_5 , 5.00 H), 2.934 (s, C_5H_6 , 0.37 H); 4 months after solution preparation, black precipitate, δ 6.503 (m, C_5H_6 , 0.20 H), 6.404 (m, C_5H_6 , 0.23 H), 5.985 (s,

C_5H_5 , 5.00 H), 2.932 (s, C_5H_6 , 0.41 H). ^{13}C NMR ($\text{THF}-d_6$) 4 months after solution preparation, δ 106.82 (s, C_5H_5).

Studies of Lewis Acidity of $\text{In}(\text{C}_5\text{H}_5)_3$. (a) NMe_3 . A flask was charged with 0.052 g (0.017 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$, 0.020 g (0.085 mmol) of NMe_3 , and 5 mL of C_6H_6 . The resulting cloudy yellow suspension was maintained at room temperature for 2 h. The material, volatile at room temperature, was removed by trap-to-trap vacuum distillation to leave a cream-colored product. Finally, the flask was dynamically evacuated for an additional 24 h and the product turned yellow. The final product weighed 0.050 g. Thus, the $\text{In}(\text{C}_5\text{H}_5)_3$ retained only a very small amount of NMe_3 . Mp: 121 °C (sample began to turn tan), 136 °C (sample turned brown), 141 °C (apparent melting). Crystals of $\text{In}(\text{C}_5\text{H}_5)$ were observed in the cooler part of the melting point capillary at the conclusion of the melting point experiment. ^1H NMR ($\text{THF}-d_6$): δ 5.85 (s, C_5H_5 , 15.0 H), 1.12 (s, NCH_3 , 1.8 H). **(b) PPh_3** . A flask was charged with 0.805 g (2.59 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$ and 0.525 g (2.00 mmol) of PPh_3 and approximately 10 mL of C_6H_6 . The resulting bright-yellow suspension in C_6H_6 was stirred for 1 h and filtered through a medium porosity frit. The benzene was removed by vacuum distillation and replaced with 10 mL of C_5H_{12} . Washing of the product with cold pentane provided 1.02 g (1.79 mmol, 89.5% yield based on PPh_3) of $\text{In}(\text{C}_5\text{H}_5)_3 \cdot \text{PPh}_3$ as pale yellow crystals. **$\text{In}(\text{C}_5\text{H}_5)_3 \cdot \text{PPh}_3$** : Mp 159 °C (sample began to turn orange), 169 °C (sample suddenly melted and became orange red), 180 °C (sample turned brown). Crystals of $\text{In}(\text{C}_5\text{H}_5)$ were observed in the cooler part of the melting point capillary at the conclusion of the melting point experiment. Anal. Calcd for $\text{C}_{33}\text{H}_{30}\text{InP}$: C, 69.25; H, 5.28. Found: C, 68.83; H, 5.24. Solubility: soluble in THF and C_6H_6 ; no appreciable solubility in pentane. ^1H NMR ($\text{THF}-d_6$): δ 5.95 (s, C_5H_5 , 15H), 7.30 (br m, PPh , 15H). ^1H NMR (C_6D_6): δ 6.20 (s, C_5H_5 , 15H), 7.00 (m, PPh), 7.43 (m, PPh). $^{31}\text{P}\{^1\text{H}\}$ NMR ($\text{THF}-d_6$): δ -3.08 (s). $^{31}\text{P}\{^1\text{H}\}$ NMR (C_6D_6) δ 5.95 (s). Cryoscopic molecular weight, benzene solution, formula weight 572 (observed molality, observed molecular weight, association): 0.0688, 582, 1.02; 0.0406, 588, 1.03; 0.0274, 584, 1.02.

Synthesis of $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)$ by a Ligand Redistribution Reaction. A THF solution of 0.655 g (4.10 mmol) of InMe_3 was allowed to react with 0.636 g (2.05 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$ at room temperature for 1 h. After the THF was removed by vacuum distillation, the flask was dynamically evacuated for an additional 3 h. The resulting colorless solid was sublimed at 100–110 °C to produce 1.19 g (5.67 mmol, 92.2% yield) of $\text{Me}_2\text{In}(\text{C}_5\text{H}_5)$. **$\text{Me}_2\text{In}(\text{C}_5\text{H}_5)$** : Colorless solid. Mp 190–196 °C dec (color change to dark brown) (lit.¹⁵ mp 195–200 °C dec). Solubility: soluble in THF; no appreciable solubility in Et_2O , C_6H_6 , or C_5H_{12} . ^1H NMR ($\text{THF}-d_6$): δ 6.04 (s, C_5H_5 , 5.00H), -0.52 (s, $\text{Me}_2\text{In}(\text{C}_5\text{H}_5) \cdot \text{THF}$ and $\text{InMe}_3 \cdot \text{THF}$, 5.78H), -1.10 (s, $\text{MeIn}(\text{C}_5\text{H}_5)_2 \cdot \text{THF}$, 0.22 H) (lit. ^1H NMR (d_6 -THF): δ 6.05 (s, C_5H_5 , 2.19H), 0.11 (s, $\text{InMe}_3 \cdot \text{THF}$, 0.48H), -0.52 (s, $\text{Me}_2\text{In}(\text{C}_5\text{H}_5) \cdot \text{THF}$, 6.0H), -1.11 (s, $\text{MeIn}(\text{C}_5\text{H}_5)_2 \cdot \text{THF}$, 0.11H).

Synthesis of $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$ by a Ligand Redistribution Reaction. A THF solution of 1.42 g (4.33 mmol) of freshly sublimed $\text{In}(\text{CH}_2\text{CMe}_3)_3$ was allowed to react with 0.672 g (2.17 mmol) of $\text{In}(\text{C}_5\text{H}_5)_3$ dissolved in THF. After the THF was removed by vacuum distillation, the product was sublimed at 100–110 °C to produce 1.923 g (5.97 mmol, 91.7% yield) of $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$. **$(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$** : Colorless solid. Mp 188–190 °C dec (color change to brown). Anal. Calcd for $\text{C}_{15}\text{H}_{27}\text{In}$: C, 55.92; H, 8.45. Found: C, 55.91; H, 8.33. Solubility: soluble in THF; no appreciable solubility in Et_2O , C_6H_6 , or C_5H_{12} . ^1H NMR ($\text{THF}-d_6$): δ 6.07 (s, C_5H_5 , 4.78H), 0.99 (s, $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$, 15.81H), 0.61 (s, $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$, 3.76H), 0.82 (s, $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2$, 0.47H), 0.49 (s, $(\text{Me}_3\text{CCH}_2)\text{In}(\text{C}_5\text{H}_5)_2$, 0.01H), 1.11 (s, $(\text{Me}_3\text{CCH}_2)_3\text{In}$, 1.83H), 0.81 (s, $(\text{Me}_3\text{CCH}_2)_3\text{In}$, 0.35H).

Table 4. Data for X-ray Crystallographic Studies of [In(C₅H₅)₃]_n, (C₅H₅)₃In·PPh₃, and [(Me₃CCH₂)₂In(C₅H₅)]_n

	[In(C ₅ H ₅) ₃] _n	(C ₅ H ₅) ₃ In·PPh ₃	[(Me ₃ CCH ₂) ₂ In(C ₅ H ₅)] _n
molec. formula	C ₁₅ H ₁₅ In	C ₃₃ H ₃₀ InP	C ₁₅ H ₂₇ In
M _r	310.09	572.36	322.19
cryst shape	yellow plate	yellow brick	colorless brick
cryst syst	orthorhombic	monoclinic	monoclinic
space group	<i>Pbca</i>	<i>P2₁/n</i>	<i>P2₁</i>
<i>a</i> , Å	9.8029(3)	11.0573(2)	9.7386(1)
<i>b</i> , Å	9.5808(3)	15.3171(3)	14.9887(2)
<i>c</i> , Å	24.6957(7)	16.8052(3)	20.7791(3)
α, deg	90	90	90
β, deg	90	108.559(1)	90.0068(4)
γ, deg	90	90	90
<i>V</i> , Å ³	2319.4(1)	2698.21(9)	3033.10(7)
<i>D</i> _{calcd} , g/cm ³	1.776	1.409	1.411
<i>Z</i>	8	4	4
μ(Mo Kα), mm ⁻¹	2.005	0.954	1.535
<i>T</i> (K)	90(1)	90.0(1)	90.0(1)
max. 2θ, deg	50.0	60.0	58.5
absorp. corr. meth.	SADABS ¹⁹	SADABS ¹⁹	SADABS ¹⁹
reflms measd	21659	48046	32340
unique reflms (<i>R</i> _{int})	2039(0.067)	7876 (0.059)	14405 (0.047)
reflms <i>I</i> > 4σ(<i>I</i>)	1526	6571	13854
refined parameters	190	406	629
<i>R</i> [<i>I</i> > 2σ(<i>I</i>)]	0.028	0.028	0.026
<i>wR</i> ₂	0.065	0.065	0.061
goodness-of-fit	1.028	1.204	1.013
extinction coeff.	None	None	0.0068(1)

Synthesis of MeIn(C₅H₅)₂ by a Ligand Redistribution Reaction with Excess In(C₅H₅)₃. The same procedure as described for the preparation of Me₂In(C₅H₅) was used to prepare MeIn(C₅H₅)₂. Reagents: 0.164 g (1.02 mmol) of freshly sublimed InMe₃, 0.953 g (3.07 mmol) of In(C₅H₅)₃, and 10 mL of THF. The resulting yellow solid was sublimed at 110–120 °C to produce 0.649 g (2.497 mmol, 81.6% yield based on InMe₃) of MeIn(C₅H₅)₂. **MeIn(C₅H₅)₂:** Yellow solid. Mp 150.7–160.2 °C dec (gradual change from yellow to viscous brown liquid). Anal. Calcd for C₁₁H₁₃In: C, 50.81; H, 5.04; In, 44.15. Found: C, 50.57; H, 5.32; In, 44.03. Solubility: soluble in THF; no appreciable solubility in Et₂O, C₆H₆, or C₅H₁₂. Attempts to grow crystallographic quality crystals by slow sublimation at 130–140 °C produced severely twinned crystals that were not useful. ¹H NMR (THF-*d*₈): δ 5.98 (s, C₅H₅, 10.7H), -0.37 (s, Me₂In(C₅H₅), 0.23H), -1.09 (s, MeIn(C₅H₅)₂, 2.06H).

Synthesis of (Me₃CCH₂)In(C₅H₅)₂ by a Ligand Redistribution Reaction with Excess In(C₅H₅)₃. The same procedure as described for the preparation of Me₂In(C₅H₅) was used to prepare (Me₃CCH₂)In(C₅H₅)₂. Reagents: 0.282 g (0.859 mmol) of freshly sublimed In(CH₂CMe₃)₃, 0.799 g (2.58 mmol) of In(C₅H₅)₃, and 10 mL of THF. The resulting yellow solid was heated at 90–95 °C to isolate the product by sublimation. A small amount of a colorless solid identified as (Me₃CCH₂)₂In(C₅H₅) that sublimed initially was removed. Then the sample was reheated and 0.766 g (2.42 mmol, 94% yield based on In(CH₂CMe₃)₃) of (Me₃CCH₂)In(C₅H₅)₂ as a yellow solid was isolated from the coldfinger of the sublimator. **(Me₃CCH₂)In(C₅H₅)₂:** Yellow solid. Mp 126.4–126.9 °C dec (gradual change from yellow to viscous brown liquid). Anal. Calcd for C₁₅H₂₁In: C, 56.99; H, 6.70. Found: C, 56.86; H, 6.69. Solubility: soluble in THF; no appreciable solubility in Et₂O, C₆H₆, or C₅H₁₂. ¹H NMR (THF-*d*₈): δ 6.03 (s, C₅H₅, 10.14H), 0.99 (s, (Me₃CCH₂)₂In(C₅H₅), 0.91H), 0.61 (s, (Me₃CCH₂)₂In(C₅H₅), 0.19H), 0.83 (s, (Me₃CCH₂)In(C₅H₅)₂, 7.87H), 0.37 (s, (Me₃CCH₂)In(C₅H₅)₂, 1.77H), 1.10 (s, (Me₃CCH₂)₃In, 0.11H).

Thermal Decomposition of (Me₃CCH₂)In(C₅H₅)₂. A break-seal tube that contained 0.972 g (3.07 mmol) of (Me₃CCH₂)In(C₅H₅)₂ was evacuated and flame sealed. The tube was placed halfway into a 150 °C oven. Immediately, the (Me₃CCH₂)In(C₅H₅)₂ began to melt and pale yellow crystals collected at the coolest end of the tube while colorless needles formed on the walls of the tube immediately above the oven. After the tube was heated for 15 min, a trace of a brown

residue was observed at the bottom of the tube. The tube was opened and the condensable gases transferred to an NMR tube. The yellow crystalline solid was collected in the drybox and identified as In(C₅H₅) (0.211 g, 1.17 mmol, 76% yield based on eqs 5 and 6) by its physical properties and ¹H NMR spectrum. The colorless needles were identified as (Me₃CCH₂)₂In(C₅H₅) (0.452 g, 1.40 mmol, 91% based on eqs 5 and 6). **Pale yellow crystals:** In(C₅H₅). Mp 169.8–170.4 °C dec. ¹H NMR (THF-*d*₈): δ 5.99 (s, C₅H₅). **Colorless needles:** (Me₃CCH₂)₂In(C₅H₅). Mp 188.2–191.4 °C dec. ¹H NMR (THF-*d*₈): δ 6.08 (s, C₅H₅, 4.89H), 0.99 (s, (Me₃CCH₂)₂In(C₅H₅), 15.75H), 0.61 (s, (Me₃CCH₂)₂In(C₅H₅), 3.74H), 0.82 (s, (Me₃CCH₂)In(C₅H₅)₂, 0.51H), 0.49 (s, (Me₃CCH₂)In(C₅H₅)₂, 0.01H), 1.11 (s, (Me₃CCH₂)₃In, 1.79H), 0.82 (s, (Me₃CCH₂)₃In, 0.31H). **Volatile liquid:** C₅H₆ and CMe₄. ¹H NMR (C₆D₆): δ 6.48 (m, C₅H₆, 15.13 H), 6.29 (m, C₅H₆, 16.25 H), 3.58 (m, C₅H₆, 1.22 H), 2.68 (m, C₁₀H₁₂, 27.78 H), 1.41 (m, C₁₀H₁₂, 1.28 H), 1.24 (m, C₁₀H₁₂, 1.20 H), 0.91 (s, CMe₄, 12 H), 0.86 (m, C₁₀H₁₂, 1.35 H), 0.39 (s, 0.38 H).

Collection of X-ray Diffraction Data and Structural Solutions for [In(C₅H₅)₃]_n, In(C₅H₅)₃·PPh₃, and [(Me₃CCH₂)₂In(C₅H₅)]_n. **General Features.** Well-defined crystals of the title compounds were covered with Infineum V8512 oil (Infineum USA L. P., 1900 East Linden Avenue, Linden, NJ 07036) and mounted on a Bruker SMART1000 CCD diffractometer equipped with the rotating anode (Mo Kα radiation, λ = 0.71073 Å). X-ray diffraction data were collected at 90 K. Details for all compounds are provided in Table 4. Data collection for each compound involved four sets of frames (600 frames in each set) and covered half-reciprocal space using the ω-scans technique (0.3° frame width) with different φ angles. Reflection intensities were integrated by using the SAINT-PLUS program.¹⁹ The solution and refinement of the structures were performed by use of the SHELXTL program package.²⁰ The structures were refined by full-matrix least squares against *F*². Non-hydrogen atoms were refined in anisotropic approximation. The hydrogen atoms were located

(19) SMART and SAINTPLUS, Area detector control and integration software, Ver. 6.0 1; Bruker Analytical X-ray Systems: Madison, WI, 1999.

(20) SHELXTL, Ver. 5.10; An integration system for solving, refining, and displaying crystal structures from diffraction data; Bruker Analytical X-ray Systems: Madison, WI, 1997.

from difference electron density Fourier syntheses and refined in isotropic approximation for $[\text{In}(\text{C}_5\text{H}_5)_3]_n$, with $U_{\text{iso}} = 1.2 U_{\text{eq}}$ of the preceding carbon atom for $\text{In}(\text{C}_5\text{H}_5)_3 \cdot \text{PPh}_3$, or as idealized CH_3 groups with $U_{\text{iso}} = 1.5 U_{\text{eq}}$ and by using the riding model with $U_{\text{iso}} = 1.2 U_{\text{eq}}$ of the preceding carbon atom for $[(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)]_n$. Data were corrected for absorption by using the Bruker AXS SADABS program that is a part of the SAINTPLUS package.¹⁹

The structure of $(\text{Me}_3\text{CCH}_2)_2\text{In}(\text{C}_5\text{H}_5)$ provides an example of twinning by *pseudo*-merohedry,^{21,22} which occurs when the metric symmetry is higher than the symmetry of the structure. The structure appeared to be orthorhombic with all angles in the unit cell of exactly 90° . Although the $|E^2 - 1|$ value of 0.867 was close to that for the centrosymmetric space group, the systematic absences only for three mutually orthogonal 2₁ screw axes were found giving only one acceptable space group $P2_12_12_1$ and indicating the presence of both racemic twinning and general twinning in the structure. The structure could then be solved in the monoclinic space group $P2_1$, which

(21) Giacovazzo, C. *Fundamentals of Crystallography*; Oxford University Press: Oxford, UK, 1992.

(22) Herbst-Irmer, R.; Sheldrick, G. M. *Acta Crystallogr.* **1998**, *B54*, 443.

requires the transformation of the unit cell. The R_{int} value for the orthorhombic space group was 0.083, significant but just slightly higher than that for the monoclinic space group (0.047). The structure was solved by the Patterson method, which gave the positions of the indium atoms. Structural refinement was completed by using the twinning law (1 0 0; 0 -1 0; 0 0 -1) for the generation of the unequal components, which were doubled to satisfy the racemic twinning. The three scale factors for twin components were refined to values 0.20385, 0.35935, and 0.16834.

Supporting Information Available: Experimental details for the attempted synthesis of $\text{In}(\text{C}_5\text{H}_5)_3$ by the literature⁸ method; variable-temperature NMR spectral data for heteroleptic organoindium cyclopentadienide compounds; solubility and stability of $\text{In}(\text{C}_5\text{H}_5)_3$ in benzene; and complete tables of positional parameters, interatomic distances and angles, anisotropic thermal parameters, and positions for hydrogen atoms and packing diagrams for the three compounds studied. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM0202322