

Polymorphism in the Crystal Structures of the Group 13 Trimethyls

Roland Boese,[†] Anthony J. Downs,[‡] Timothy M. Greene,[§] Alexander W. Hall,[‡]
Carole A. Morrison,^{||} and Simon Parsons^{*,||}

Institut für Anorganische Chemie, Universität Essen, Universitätsstrasse 3-5, 45117 Essen, Germany; Inorganic Chemistry Laboratory, University of Oxford, South Parks Road, Oxford, England OX1 3QR; School of Chemistry, University of Exeter, Chemistry Building, Stocker Road, Exeter, England EX4 4QD; and School of Chemistry, The University of Edinburgh, King's Buildings, West Mains Road, Edinburgh, Scotland EH9 3JJ

Received January 14, 2003

Crystal structures have been determined for trimethylboron, BMe_3 , and for a new polymorph of trimethylgallium, GaMe_3 ; in addition, the crystal structure of trimethylthallium, TlMe_3 , has been redetermined. The BMe_3 crystal structure represents a new structural type for the group 13 trimethyl derivatives in the solid state. In contrast to its heavier analogues, it consists of layers containing only very weakly interacting BMe_3 molecules. GaMe_3 forms a ladder-like pseudo-polymer via long gallium-to-methyl intermolecular interactions with $\text{Ga}\cdots\text{C}$ distances in the range 3.096(3)–3.226(4) Å. This is compared with a recently reported crystal structure of a polymorph, which, like InMe_3 and TlMe_3 , is characterized by the formation of pseudo-tetramers. The effects of crystallization and secondary interactions have been analyzed by comparison with related crystallographic, gas-phase electron diffraction, and spectroscopic studies of these and other trimethyl derivatives of the group 13 elements. The energetic differences between polymorphs of BMe_3 , GaMe_3 , and InMe_3 have been explored by plane wave DFT calculations. The energy differences between the BMe_3 -like layered structure and the InMe_3 -like pseudo-tetrameric structure are calculated to be -1.7 , $+3.6$, and $+10.4$ kJ mol⁻¹ for BMe_3 , GaMe_3 , and InMe_3 , respectively.

Introduction

There are three different structures that are observed for the group 13 trimethyl derivatives in the crystalline state. One is the dimeric form observed for trimethylaluminum,¹ in which methyl groups participate in strong metal–metal bridges, with C–Al interaction distances commensurate with those of terminal Al–C bonds. Methyl bridging is also observed in a second structural type featured by the derivatives with Ga, In, and Tl, but the interaction distances are substantially longer than the primary metal–carbon bonds. Prior to this study, crystal structures had been determined for GaMe_3 ,² InMe_3 ,^{3,4} and TlMe_3 ,⁵ here we describe crystal structure determinations of a new polymorph of GaMe_3 and a redetermination of the structure of TlMe_3 . The third structural type is represented by BMe_3 . As we show below, this crystal structure consists of layers in which the molecules interact by weak van der Waals,

or possibly electrostatic, interactions. The longstanding debate over the nature and geometry of the bridging methyl groups in Al_2Me_6 has recently been resolved in a neutron powder diffraction study at 4.5 K,⁶ and in this paper we limit our attention to the second and third structural types.

We have also investigated theoretically the energetic differences between polymorphs of BMe_3 , GaMe_3 , and InMe_3 in which the molecules form layers (as in the observed structure of BMe_3) or weak methyl bridges (as in InMe_3). Polymorphism has been described as the supramolecular equivalent of molecular isomerism.⁷ Traditional ab initio modeling procedures (notably GAUSSIAN) simulate isolated molecules, but while this style of calculation is suitable for studying gaseous isomers, it is not so readily applicable to solids. Plane wave density functional theory (DFT) can simulate the periodic wave function characteristics of a repeating unit such as a crystallographic unit cell. The lattice parameters and atomic positions can all be varied to minimize the crystal lattice energy, atomic forces, and unit cell stress, and we therefore use these methods to draw energetic comparisons between trimethyl polymorphs.

Our results complete the series of crystal structures for the group 13 trimethyls. The trimesityls (mesityl =

* To whom correspondence should be addressed. Tel: +44 131 650 5804. Fax: +44 131 650 4743. E-mail: S.Parsons@ed.ac.uk.

[†] Universität Essen.

[‡] University of Oxford.

[§] University of Exeter.

^{||} The University of Edinburgh.

(1) Vranka, R. G.; Amma, E. L. *J. Am. Chem. Soc.* **1967**, *89*, 3121.
Byram, S. K.; Fawcett, J. K.; Nyberg, S. C.; O'Brien, R. J. *J. Chem. Soc. D* **1970**, 16. Huffman, J. C.; Streib, W. E. *J. Chem. Soc. D* **1971**, 911.

(2) Mitzel, N. W.; Lustig, C.; Berger, R. J. F.; Runeberg, N. *Angew. Chem., Int. Ed.* **2002**, *41*, 2519.

(3) Amma, E. L.; Rundle, R. E. *J. Am. Chem. Soc.* **1958**, *80*, 4141.

(4) Blake, A. J.; Craddock, S. *J. Chem. Soc., Dalton Trans.* **1990**, 2393.

(5) Sheldrick, G. M.; Sheldrick, W. S. *J. Chem. Soc. A* **1970**, 28.

(6) McGrady, G. S.; Turner, J. F. C.; Ibberson, R. M.; Prager, M. *Organometallics* **2000**, *19*, 4398.

(7) Bernstein, J. *Polymorphism in Molecular Crystals*, Clarendon Press: Oxford, U.K., 2002.


2,4,6-trimethylphenyl) are the only other group 13 organometallics for which a complete series of crystal structures has been determined.

Experimental Section

Synthesis of Compounds. (i) Trimethylboron was prepared by direct metathesis between trimethylaluminum (obtained from Aldrich and purified by fractional condensation in vacuo) and tri-*n*-butyl borate (also obtained from Aldrich) as neat liquids.⁸ To moderate the exothermic reaction, the mixture was held in a Pyrex glass vessel fitted with a greaseless (Young's) valve at temperatures <233 K. Trimethylboron was the only volatile product. It was purified by fractional condensation in vacuo with traps held at 162, 144, and 77 K. The fraction collected at 144 K was identified as essentially pure trimethylboron on the evidence of the IR spectrum of the vapor.⁹

(ii) Trimethylgallium was prepared similarly by ligand redistribution between trimethylaluminum and gallium(III) chloride (obtained from Aldrich).¹⁰ Warming the mixture to room temperature caused a vigorous reaction to occur. After the mixture was stirred for 90 min to ensure completion of the reaction, the volatile products were vaporized in vacuo and fractionated via traps held at 222, 178, and 77 K. Trimethylgallium was collected at 178 K and was authenticated by the IR spectrum of the vapor.¹¹

(iii) Trimethylthallium was synthesized rather differently, namely in accordance with eq 1 by the addition of an excess of methylolithium in ether solution (1.4 M) to thallium(I) iodide in the presence of iodomethane (all reagents being used as supplied by Aldrich).¹² After the mixture had been



stirred for 1 h at room temperature, the ether solution was siphoned off into a clean, preconditioned Schlenk tube. The solvent was evaporated under reduced pressure and the trimethylthallium isolated as a crystalline solid by fractionation in vacuo, being retained by a trap held at 242 K. The purity of the product was checked by reference to the IR spectrum of its vapor at ambient temperatures.¹³

Crystal Growth. Samples of BMe₃, GaMe₃, and TlMe₃ were loaded into Pyrex capillaries and mounted on a Bruker Smart Apex CCD diffractometer equipped with an Oxford Cryosystems low-temperature device.¹⁴ The boron and gallium compounds (the former being a gas and the latter a liquid under ambient conditions) were respectively frozen at 100 and 213 K, and crystals were grown in situ by means of Boese's zone refinement method using an OHCD infrared laser-assisted crystallization device.¹⁵ The sample of TlMe₃ was treated similarly, but since this tended to sublime rather than melt under laser irradiation, a suitable crystal was obtained by careful sublimation inside the capillary at 273 K. Data collections were carried out using graphite-monochromated Mo

Table 1. Crystallographic Data Collection and Refinement Parameters

	BMe ₃	GaMe ₃	TlMe ₃
formula	C ₃ H ₉ B	C ₃ H ₉ Ga	C ₃ H ₉ Tl
<i>M_r</i>	55.91	114.82	249.47
cryst syst	monoclinic	monoclinic	tetragonal
space group	<i>C2/c</i>	<i>C2/c</i>	<i>P4₂/n</i>
<i>a</i> /Å	6.3473(9)	18.409(3)	13.3049(7)
<i>b</i> /Å	10.9284(16)	6.2652(9)	13.3049(7)
<i>c</i> /Å	14.224(2)	18.268(3)	6.2891(5)
β /deg	91.099(3)	91.361(2)	90
<i>V</i> /Å ³	986.5(2)	2106.4(5)	1113.30(12)
<i>Z</i>	95	120	150
<i>D_c</i> /Mg m ⁻³	8	16	8
μ /mm ⁻¹	0.753	1.448	2.977
range of transmission	0.038	5.044	28.844
cryst dimens/mm	0.396–1	0.071–0.272	0.233–1
cryst habit	0.5 × 0.5 × 1	0.26 × 0.26 × 1	0.4 × 0.2 × 0.2
θ_{max} /deg	colorless	colorless	colorless
no. of rflns:	cylinder	cylinder	block
total/unique	25.00	26.49	29.00
no. of data with <i>F</i> > 4σ(<i>F</i>)	12 095/1804	8030/2168	8902/1413
<i>R</i> _{int}	1487	1789	1143
no. of restraints	0.0328	0.0346	0.067
no. of params	300	0	0
<i>R</i> (<i>F</i> > 4σ(<i>F</i>))	102	85	41
<i>R_w</i> (<i>F</i> ² , all data)	0.0456	0.0294	0.0351
max shift/su	0.1344	0.0753	0.0890
final diff map extremes/e Å ⁻³ .	0.001	0.002	0.002
	+0.11, -0.13	+0.81, -0.50	+1.75, -1.69

K α radiation ($\lambda = 0.71073 \text{ \AA}$); collection parameters are listed in Table 1 or in the Supporting Information.

Crystal Structure Determination of BMe₃. Diffraction data were collected for a sample at 95 K. Complete indexing of the diffraction pattern of the sample of BMe₃ described above required two orientation matrices.¹⁶ The relationship between the component orientations could be described with the matrix

$$\begin{pmatrix} -1.00 & -0.03 & 0.00 \\ -0.09 & 1.00 & 0.04 \\ 0.01 & 0.06 & -1.00 \end{pmatrix},$$

which approximates to a 180° rotation about [010]. Since this is a symmetry operation of a monoclinic lattice, the result implies that the sample consisted of two slightly misaligned crystals. Such "twinning" conditions may give rise to refinement difficulties because reflections from different domains partially overlap, but such problems were avoided by simultaneous integration of both components, which ensures that fully and partially overlapping reflections are treated correctly.¹⁷ An absorption correction was applied using the recently written program TWINABS,¹⁸ which is based on the multiscan procedure of Blessing¹⁹ and designed to treat twinned data. Data from both components were used in refinement, the final residuals being only some 0.2% higher than if pure single-component data were used.

The structure was solved by direct methods and refined by full-matrix least squares against $|F|^2$ using all data (SHELXTL),²⁰ with anisotropic displacement parameters modeled for the B and C atoms. It was clear from electron density difference maps that the methyl groups were disordered by a

(8) Köster, R. *Justus Liebigs Ann. Chem.* **1958**, 618, 31.
 (9) Woodward, L. A.; Hall, J. R.; Dixon, R. N.; Sheppard, N. *Spectrochim. Acta* **1959**, 15, 249.
 (10) Gaines, D. F.; Borlin, J.; Fody, E. P. *Inorg. Synth.* **1974**, 15, 203.
 (11) McKean, D. C.; McQuillan, G. P.; Duncan, J. L.; Sheppard, N.; Munro, B.; Fawcett, V.; Edwards, H. G. M. *Spectrochim. Acta* **1987**, 43A, 1405.
 (12) Gilman, H.; Jones, R. G. *J. Am. Chem. Soc.* **1946**, 68, 517.
 (13) Schwerdtfeger, P.; Bowmaker, G. A.; Boyd, P. D. W.; Ware, D. C.; Brothers, P. J.; Nielson, A. J. *Organometallics* **1990**, 9, 504.
 Johnson, E. M. D.Phil. Thesis, University of Oxford, 1971.
 (14) Cosier, J.; Glazer, A. M. *J. Appl. Crystallogr.* **1986**, 19, 105.
 (15) Boese, R.; Nussbaumer, M. In *Correlations, Transformations, and Interactions in Organic Crystal Chemistry*; Jones, D. W., Katrusiak, A., Eds.; IUCr Crystallographic Symposia 7; Oxford University Press: Oxford, U.K., 1994; p 20.

(16) Sparks, R. A. GEMINI: Program for Indexing X-ray Diffraction Patterns Derived from Multiple or Twinned Crystals; Bruker-AXS, Madison, WI, 2000.
 (17) SAINT, version 6.33; Bruker-Nonius, Madison, WI, 2002.
 (18) Sheldrick, G. M. TWINABS: Program for Performing Absorption Corrections to X-ray Diffraction Patterns Collected from Non-Merohedrally Twinned and Multiple Crystals; University of Göttingen, Göttingen, Germany, 2002.
 (19) Blessing, R. H. *Acta Crystallogr.* **1995**, A51, 33.

Table 2. Observed Bond and Contact Distances (Å) and Angles (deg) in MMe_3 ($M = B, Ga, Tl$)^a

BMe_3 (obsd, $C2/c$)		$GaMe_3$ (obsd, $C2/c$)		$GaMe_3$ (obsd, $P4_2/n$)		$TlMe_3$ (obsd, $P4_2/n$)	
B1–C1	1.5548(11)	Ga1–C1	1.956(3)	Ga1–C1	1.952(3)	Tl1–C1	2.196(8)
B1–C2	1.5565(11)	Ga1–C2	1.958(3)	Ga1–C2	1.962(2)	Tl1–C2	2.206(8)
B1–C3	1.5541(12)	Ga1–C3	1.968(3)	Ga1–C3	1.958(2)	Tl1–C3	2.216(7)
		Ga1...C5 ^b	3.096(3)	Ga1...C2 ^f	3.149(3)	Tl1...C2 ^h	3.243(8)
		Ga1...C6 ^c	3.512(4)	Ga1...C3 ^g	3.647(3)	Tl1...C3 ⁱ	3.364(7)
		Ga2–C4	1.964(3)				
		Ga2–C5	1.970(3)				
		Ga2–C6	1.956(3)				
		Ga2...C2 ^d	3.226(3)				
		Ga2...C3 ^e	3.204(3)				
C1–B1–C2	120.06(7)	C1–Ga1–C2	122.02(16)	C1–Ga1–C2	119.72(14)	C1–Tl1–C2	120.7(3)
C1–B1–C3	120.09(7)	C1–Ga1–C3	119.46(16)	C1–Ga1–C3	121.29(14)	C1–Tl1–C3	124.1(3)
C2–B1–C3	119.86(7)	C2–Ga1–C3	118.27(16)	C2–Ga1–C3	118.85(13)	C2–Tl1–C3	115.1(3)
		Ga1...C5 ^b –Ga2 ^b	171.09(18)	Ga1...C2 ^f –Ga1 ^f	167.22(12)	Tl1...C2 ^h –Tl1 ^h	167.8(4)
		Ga1...C6 ^c –Ga2 ^c	120.02(17)	Ga1...C3 ^g –Ga1 ^g	164.67(13)	Tl1...C3 ⁱ –Tl1 ⁱ	167.5(4)
		C4–Ga2–C5	120.45(17)				
		C4–Ga2–C6	120.16(18)				
		C5–Ga2–C6	119.39(17)				
		Ga2...C2 ^d –Ga1 ^d	162.14(18)				
		Ga2...C3 ^e –Ga1 ^e	168.71(17)				

^a Dimensions for the tetragonal phase of $GaMe_3$ were calculated from the data in ref 2 (data taken from CCDC 163477). All standard uncertainties were calculated with a full variance–covariance matrix, with the exception of those given in italics. ^b $-x, y + 1, 1/2 - z$. ^c $1/2 - x, 1/2 + y, 1/2 - z$. ^d $x, y - 1, z$. ^e $x, -y, 1/2 + z$. ^f $y, 3/2 - x, 1/2 - z$. ^g $y - 1/2, 1 - x, z - 1/2$. ^h $3/2 - y, x, 1/2 - z$. ⁱ $1 - y, x - 1/2, z - 1/2$.

180° rotation about their B–C axes. The relative occupancies were initially refined independently, but later these were modeled with one variable for all three methyl groups after they had converged to common values. The occupancy of the major component (labeled H1A–H1C etc. in the Supporting Information) was 0.577(6).

The orientation of the methyl groups was such that one C–H bond lay in the molecular BC_3 plane, and some distortion from ideal, local C_{3v} symmetry was anticipated (see below). The H atom positions were refined subject to the restraint that all in-plane BCH angles were similar. Similarity restraints were also applied to the out-of-plane BCH angles and all 1,2-C–H and 1,3-H...H distances. "Opposite" H atoms attached to the same carbon atom but in different disorder components were constrained to have equal isotropic displacement parameters.

Crystal Structure Determinations of $GaMe_3$ and $TlMe_3$. Diffraction data were collected for crystals held at 120 K ($GaMe_3$) and 150 K ($TlMe_3$), and absorption corrections were applied using the program SADABS.²¹ The structures were solved by direct methods and refined by full-matrix least squares against $|F|^2$ using all data (SHELXTL), with anisotropic displacement parameters modeled for the metal and C atoms. The methyl groups were treated as freely rotating rigid groups. Four out of the six independent methyl groups in $GaMe_3$ were found to be disordered in a fashion similar to that described above for BMe_3 . The relative occupancies were fixed at 0.5:0.5. No attempt was made to model a deviation of the methyl groups from local C_{3v} symmetry; the improvement to refinement statistics on introduction of a more flexible model was marginal for BMe_3 and would have been negligible for this compound, where H atom scattering contributes relatively much less to the diffraction pattern.

Refinement and geometric data for all compounds are collected in Tables 1 and 2, respectively. Structural analyses used the program PLATON,²² and figures were drawn using SHELXTL or CAMERON.²³ The file CCDC 209601–209603 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via

www.ccdc.cam.ac.uk/conts/retrieving.html (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, U.K.; fax +44 1223 336033; e-mail deposit@ccdc.cam.ac.uk).

Theoretical Methods. Total energy plane-wave DFT calculations were performed using the CASTEP 4.2 simulation code²⁴ for the compounds BMe_3 , $GaMe_3$, and $InMe_3$ in ordered layered monoclinic ($C2/c$) and tetragonal ($P4_2/n$) polymorphic forms. Periodic boundary conditions allow the valence electronic wave function to be expanded in terms of a discrete plane-wave basis set (set at 540 eV for BMe_3 and 500 eV for $GaMe_3$ and $InMe_3$), while the core wave function is described by standard ultrasoft pseudopotentials available with the software package. The symmetry-reduced set of k points used to sample the reciprocal space were generated using Monkhorst–Pack grids²⁵ (dimensions $2 \times 1 \times 1$ and $1 \times 1 \times 2$ for the monoclinic and tetragonal lattices, respectively, both generating one k point in the symmetry-reduced first Brillouin zones). The generalized gradient approximation (GGA) functional PW91²⁶ was used to model electronic correlation and exchange. Simultaneous optimization of lattice vectors and atomic positions was performed until the convergence criteria were met (maximum energy change per atom 2×10^{-5} eV, maximum RMS displacement 1.0×10^{-3} Å, maximum RMS force 0.05 eV Å⁻¹, and maximum RMS stress 0.1 GPa). The starting geometries used for the optimizations of the monoclinic lattice polymorphs were taken from the experimental structure determination of BMe_3 reported in this paper, with the boron atoms simply replaced by gallium or indium to generate input coordinates for the other two structures. The tetragonal lattice atomic coordinates and cell vectors for the BMe_3 and $InMe_3$ structures originated from Blake's $InMe_3$ X-ray structure determination;⁴ calculations on $GaMe_3$ used Mitzel's data² on the tetragonal polymorph as the starting

(22) Spek, A. L. PLATON: A Multipurpose Crystallographic Tool; Utrecht University, Utrecht, The Netherlands, 2002. PC version: Farrugia, L. J. *J. Appl. Crystallogr.* **1999**, *32*, 837.

(23) Watkin, D. J.; Pearce, L.; Prout, C. K. CAMERON—A Molecular Graphics Package; Chemical Crystallography Laboratory, University of Oxford, Oxford, U.K., 1993.

(24) Payne, M. C.; Teter, M. P.; Allan, D. C.; Arias, T. A.; Joannopoulos, J. D. *Rev. Mod. Phys.* **1992**, *64*, 1045. CASTEP 4.2 Academic version, licensed under the UKCP-MSI agreement, 1999.

(25) Monkhorst, H. J.; Pack, J. D. *Phys. Rev. B* **1976**, *13*, 5188.

(26) Perdew, J. P.; Chevary, J. A.; Vosko, S. H.; Jackson, K. A.; Pederson, M. R.; Singh, D. J.; Fiolhais, C. *Phys. Rev. B* **1992**, *46*, 5571.

(20) Sheldrick, G. M. SHELXTL: Suite of Programs for Crystal Structure Analysis, Incorporating Structure Solution (XS), Least-Squares Refinement (XL), and Graphics (XP); University of Göttingen, Göttingen, Germany, 2001.

(21) Sheldrick, G. M. SADABS: Program for Performing Absorption Corrections to Single-Crystal X-ray Diffraction Patterns; University of Göttingen, Göttingen, Germany, 2002.

point (as this structure is the most directly comparable with the others in our series).

The Supporting Information contains tables of crystallographic data for BMe_3 , GaMe_3 , and TlMe_3 (also deposited with the CCDC, as described above) and tables of optimized theoretical coordinates for polymorphs of BMe_3 , GaMe_3 , and InMe_3 .

Results and Discussion

Under ambient conditions, the trimethyl derivatives of the group 13 elements range from a gas (BMe_3), through a liquid (AlMe_3 and GaMe_3), to a low-melting solid (InMe_3 and TlMe_3).²⁷ On the evidence of mass and vibrational spectroscopic and electron diffraction measurements, the vapors of all but the aluminum compound consist of monomeric MMe_3 molecules ($\text{M} = \text{B}, \text{Ga}, \text{In}, \text{Tl}$), each with a trigonal-planar MC_3 skeleton and more or less freely rotating methyl groups.^{9,11,13,28–31} In contrast, the analogous aluminum species, AlMe_3 , is found in appreciable concentrations only at elevated temperatures and/or low pressures;^{32,33} otherwise, the dimer $\text{Me}_2\text{Al}(\mu\text{-Me})_2\text{AlMe}_2$ prevails throughout the condensed and vapor states.⁶ Increasing the atomic number of the group 13 element results in an overall rise in melting and boiling points consistent with the expected strengthening of van der Waals interactions. That the pattern is far from regular, however, is evidenced by the following melting points (in K):²⁷ BMe_3 , 112; AlMe_3 , 288; GaMe_3 , 257; InMe_3 , 362; TlMe_3 , 312. To what extent the implied cohesive energies of the crystal reflect differences of structure and/or variations in the type or degree of the intermolecular interactions—possibly including so-called “agostic” interactions³⁴—is not possible to judge on the evidence available to date. Previous studies involving X-ray crystallography,^{1–5} gas electron diffraction,^{28–30} and vibrational spectroscopy^{4,11,13} argue that perturbation of the MMe_3 units in the crystal is quite modest, with the sole exception of $\text{M} = \text{Al}$, for which the dimer $\text{Me}_2\text{Al}(\mu\text{-Me})_2\text{AlMe}_2$ holds sway (but is itself subject to relatively little change with the transition from the vapor to the crystalline state⁶). One or two of these studies have also alluded to the possibility of polymorphism, for example in the case of InMe_3 .^{3,4} The aims of the present study have been to enlarge on knowledge of the crystal structures of these compounds and of the secondary interactions they reveal and to explore the possibilities of polymorphism, partly by experiment (in the case of GaMe_3) but, more widely, by plane wave DFT analysis.

Trimethylboron. The boron-to-carbon distances in BMe_3 are equal within error, and the BC_3 framework




Figure 1. Crystal structure of BMe_3 viewed along the crystallographic c direction. The molecules are related either by lattice translations or C -centering operations. The minor disorder component has been omitted for clarity. Displacement ellipsoids enclose 50% probability surfaces.

adopts the expected D_{3h} symmetry (Figure 1). The most closely related crystal structure in the literature is that of triethylboron, BEt_3 ; the B–C distances in that compound lie around 1.573(1) Å.³⁵ The corresponding distance in BMe_3 is 1.555(1) Å, but a riding analysis³⁶ suggests that this difference is probably owed to the relatively high librational motion of the methyl groups at the temperature used for data collection (95 K), which is only 17 K below the melting point. In this context, it is perhaps significant that the B–C distance in the gaseous BMe_3 molecule (determined by electron diffraction) is reported to be 1.5783(11) Å.²⁸

The BCC angle in BEt_3 would be expected, on the basis of simple predictions using VSEPR theory, for example, to be close to 109.5°, and the most remarkable feature of the structure of this compound is the rather large BCC angle, reported to be 118.9(2)°. A similar effect has recently been observed in GaEt_3 .² The reason for this deviation has been ascribed to hyperconjugation between the out-of-plane CH_2 bonds and the vacant p orbital on the central group 13 atom. The methyl groups in BMe_3 are disordered by a 180° rotation about the B–C vector, each component containing one CH bond in the BC_3 plane. The ab initio optimized crystal structure of an ordered model of BMe_3 (see below) revealed that the average in-plane BCH angle was 115°, whereas the average out-of-plane BCH angle was 110°. Scattering from the H atoms in BMe_3 contributes some 28% to $F(000)$, giving them a significant influence on data fitting, and it seemed possible that a deviation from ideal tetrahedral geometry about the carbon atoms in BMe_3 might be detectable, despite the disorder. Restrained refinement of a model in which the BCH angles

(27) Starowieyski, K. In *Chemistry of Aluminium, Gallium, Indium and Thallium*; Downs, A. J., Ed.; Blackie Academic and Professional: Glasgow, U.K., 1993; p 339. Macintyre, J. E., Ed. *Dictionary of Organometallic Compounds*, 2nd ed.; Chapman and Hall: London, U.K.; 1995.

(28) Bartell, L. S.; Carroll, B. L. *J. Chem. Phys.* **1965**, *42*, 3076.

(29) Beagley, B.; Schmidling, D. G.; Steer, I. A. *J. Mol. Struct.* **1974**, *21*, 437.

(30) Fjeldberg, T.; Haaland, A.; Seip, R.; Shen, Q.; Weidlein, J. *Acta Chem. Scand., Ser. A* **1982**, *36*, 495.

(31) Hall, J. R.; Woodward, L. A.; Ebsworth, E. A. V. *Spectrochim. Acta* **1964**, *20*, 1249.

(32) Almenningen, A.; Halvorsen, S.; Haaland, A. *Acta Chem. Scand.* **1971**, *25*, 1937.

(33) Atiya, G. A.; Grady, A. S.; Russell, D. K.; Claxton, T. A. *Spectrochim. Acta* **1991**, *47A*, 467. Kvisle, S.; Rytter, E. *Spectrochim. Acta* **1984**, *40A*, 939.

(34) Brookhart, M.; Green, M. L. H.; Wong, L.-L. *Prog. Inorg. Chem.* **1988**, *36*, 1. Scherer, W.; McGrady, G. S. Submitted for publication in *Angew. Chem., Int. Ed.*

(35) Boese, R.; Bläser, D.; Niederprüm, N.; Nüsse, M.; Brett, W. A.; Schleyer, P. v. R.; Bühl, M.; van Eikema Hommes, N. J. R. *Angew. Chem., Int. Ed. Engl.* **1992**, *31*, 314.

(36) Johnson, C. K. In *Crystallographic Computing, Proceedings of the International Summer School*; Munksgaard: Copenhagen, Denmark, 1970; pp 220–226.




Figure 2. Formation of layers in the crystal structure of BMe_3 (view along the crystallographic b direction). Packing within the layers is illustrated in Figure 1.

were allowed to vary revealed a trend rather similar to that observed in the BCC angles in BEt_3 , with the in-plane BCH angles averaging 116° , compared with 109° for the out-of-plane BCH angles, in very good agreement with the theoretical results. The standard uncertainties of these quantities (excluding the effects of restraints) fall in the range $1.1\text{--}1.8^\circ$. This places the difference on the limit of statistical significance. A refinement model in which the methyl groups were constrained to adopt perfect C_{3v} symmetry yielded a conventional R factor of 5.2% (48 parameters), compared with 4.5% for the model described here (102 parameters). For what it is worth, the Hamilton test³⁷ using weighted residuals implies that the improvement is significant.

In the crystal structure of BMe_3 the molecules pack in layers which stack along the c direction (Figure 2). The boron and hydrogen atoms respectively carry small positive and negative charges, and the boron atoms lie between methyl groups in neighboring layers. The shortest intermolecular $\text{B}\cdots\text{H}$ contact is 3.04 \AA , which is well beyond the sum of the van der Waals radii of B and H (2.83 \AA),³⁸ a finding consistent no doubt with the high volatility of trimethylboron. The arrangement of the molecules within the layers (Figure 1) resembles a close-packed array.

Trimethylgallium and Trimethylthallium. Trimethylboron is unique among the group 13 trimethyls in showing no significant association in the solid state. Trimethylaluminum exists as a methyl-bridged dimer both in the solid state and in the gas phase. Although such behavior is not observed for Ga, In, and Tl, it has long been clear that these trimethyls are also associated via secondary metal \cdots methyl contacts in the solid state. The melting points of GaMe_3 (257 K), InMe_3 (362 K), and TlMe_3 (312 K) alternate along the series, and are all significantly higher than that of BMe_3 (112 K), partly as a result of these interactions.




Figure 3. Structure of TlMe_3 projected onto (001), showing methyl bridge formation. Hydrogen atoms have been omitted for clarity. Ellipsoids enclose 50% probability surfaces.

Trimethylindium has the highest melting point of the trimethyl derivatives formed by the three heaviest group 13 metals, and it might be inferred from this that it exhibits the strongest degree of association in the solid state. Its crystal structure, first investigated by Amma and Rundle in 1958³ and then again by Blake and Cradock in 1990,⁴ is characterized by the formation of pseudo-tetramers (Rundle's term) which consist of four InMe_3 molecules connected via long $\text{In}\cdots\text{methyl}$ bridges ($\text{In}\cdots\text{C} = 3.083(12)\text{ \AA}$). These are disposed about a crystallographic 4 site, forming a flattened tetrahedron. Longer $\text{In}\cdots\text{methyl}$ bridges ($\text{In}\cdots\text{C} = 3.558(15)\text{ \AA}$) connect the tetramers. Projections of this structure along [001] (Figure 3 shows the isostructural Tl derivative in this projection) can beguile one into thinking that this structure is a two-dimensional network. In fact, it is three-dimensional, consisting of two mutually exclusive networks which are interlaced by means of a 4-fold screw axis.³

An atoms-in-molecules analysis³⁹ on the related gallium system has shown that the critical point in the metal \cdots methyl bridge region occurs along the $\text{Ga}\cdots\text{C}$ vector, implying that this is not an agostic interaction.² In the following sections we discuss intermolecular interactions in terms of metal-to-carbon distances partly for this reason, but also because H atom positions have not been determined very precisely and because they are consistent with the contemporary literature in this area.

The crystal structure of TlMe_3 was investigated using photographic methods by Sheldrick and Sheldrick in 1970.⁵ Our data set establishes the structural parameters to greater precision than was possible in that study, although the conclusions are unchanged. TlMe_3 adopts the InMe_3 structure (Figure 3), but there is a much smaller difference between the lengths of the short and long secondary metal to methyl contacts: the $\text{Tl}\cdots\text{C}$ distances are $3.243(8)$ and $3.364(7)\text{ \AA}$ within and

(37) Hamilton, W. C. *Acta Crystallogr.* **1965**, *18*, 502.

(38) Bondi, A. J. *Phys. Chem.* **1964**, *68*, 441.

(39) Bader, R. F. W. *Atoms in Molecules: A Quantum Theory*; Clarendon Press: Oxford, U.K., 1994.




Figure 4. Pseudo-polymer formed in the crystal structure of the monoclinic polymorph of GaMe_3 . The heavy dotted lines are short contacts of 3.096(3) Å; the light dotted lines are longer contacts of 3.204(3) and 3.226(3) Å. The ellipsoids enclose 50% probability surfaces; H atoms have been omitted for clarity.

between the tetramers, respectively. The primary Tl–C bond distances, averaging 2.206(8) Å, are identical with those in the gaseous molecule (2.206(3) Å), as gauged by electron diffraction.³⁰

Very recently Mitzel et al. reported a crystal structure of GaMe_3 which is isostructural with that of InMe_3 .² In this tetragonal phase both the $\text{Ga}\cdots\text{methyl}$ contacts which form the tetramers and those between the tetramers are slightly longer than in InMe_3 (3.134 and 3.647 Å, respectively). This phase was obtained by cooling a sample of GaMe_3 through its melting point, but crystal growth by laser-assisted zone refinement yielded a new polymorph of GaMe_3 , which is *C*-centered monoclinic. The structure contains two crystallographically independent molecules, both of which adopt the expected trigonal-planar geometry in the primary coordination sphere.

The length of the *b* axis of the unit cell of the monoclinic phase is similar to that of the *c* axis of the tetragonal phase, and when projected along these directions the structures bear a close resemblance to each other. However, the tetramers which characterize the tetragonal phase of GaMe_3 are replaced in the monoclinic phase by a polymer. The two crystallographically independent molecules alternate along chains with $\text{methyl}\cdots\text{Ga}$ contacts of 3.226(3) and 3.204(3) Å formed above and below the planes of the molecules containing Ga2. Shorter contacts of 3.096(3) Å made to Ga1 serve to link the chains together into a ladderlike array (Figure 4). Long contacts measuring 3.512(4) Å are formed between the ladders, completing the trigonal-bipyramidal coordination about Ga1. In the tetragonal forms of MMe_3 (*M* = Ga, In, Tl) the angles subtended at bridging carbon atoms fall in the range 160–170°. This trend is also followed in the structure of monoclinic GaMe_3 , except in the case of the long interconnecting methyl bridge (C6), where the angle is 120.02(17)°.

Figure 5 shows a projection of the monoclinic structure along the *c* direction; it should be compared with Figure 6, which shows the structure of TlMe_3 (which can be taken to be representative of all the tetragonal MMe_3 phases) projected perpendicular to the (110) plane. These rather similar packing arrangements are related by a shift in the relative positions of layers containing the short contacts. The pattern of contacts in the two phases is represented schematically in Figure 7, which is intended to illustrate the transition from a structure consisting of two independent three-dimen-




Figure 5. Structure of monoclinic GaMe_3 projected onto (001). The polymers shown in Figure 5 pass into the page, and different polymers have been shown in different colors. Domains of short (between 3.0 and 3.3 Å) and long (3.512 Å) $\text{Ga}\cdots\text{C}$ contacts are indicated by the letters S and L, respectively.




Figure 6. Structure of TlMe_3 projected onto (110). The two different three-dimensional networks are shown in blue and red. Domains of shorter (3.243 Å) and longer (3.364 Å) $\text{Tl}\cdots\text{C}$ contacts are indicated by the letters S and L, respectively. Isostructural tetragonal polymorphs are known for GaMe_3 (in which the short and long contacts are 3.149(3) and 3.647(3) Å, respectively) and InMe_3 (contact distances 3.083(12) and 3.558(15) Å).

sional networks in the tetragonal phases of MMe_3 (*M* = Ga, In, Tl) to a single three-dimensional network in the monoclinic phase of GaMe_3 .

Ab Initio Calculations of Polymorphs of Group 13 Trimethyls. There is some evidence in the literature that InMe_3 may exist in at least two different polymor-




Figure 7. Schematic representations of Figure 5 (top, monoclinic GaMe₃) and Figure 6 (bottom, TlMe₃) showing the formation of two independent networks following displacement of layers. In the top part of the figure, the ellipses represent polymers passing into the plane of the paper. In the bottom part of the figure, they represent columns of tetramers.

phic forms. Blake and Cradock⁴ refer to an alternative form of InMe₃, which they describe as being less volatile, and more stable, than the tetragonal form. However, a careful variable-temperature X-ray diffraction study between 273 and 113 K did not reveal any new phases. In their study of the same compound, Amma and Rundle³ noted that, in addition to the tetragonal phase, a less common, *less* stable, pseudo-hexagonal (more likely triclinic) form was found to exist. This conclusion was based on crystal morphology, and no diffraction data have ever been collected on this form of InMe₃. However, the reduced cell dimensions of BMe₃ are pseudo-hexagonal, with $a = b = 6.319 \text{ \AA}$, $c = 14.224 \text{ \AA}$, $\alpha = \beta = 90.55^\circ$, and $\gamma = 119.70^\circ$, and it is possible that the less stable phase is closely related to the structure of BMe₃ described here. The ab plane of this pseudo-hexagonal lattice is evident in Figure 1, perhaps most clearly by treating each B atom as a lattice point, though this would not be the conventional choice of origin.

Polymorphism is a very common phenomenon, and it seems perfectly reasonable that the group 13 trimethyls should be as susceptible to it as any other class of compound. Indeed, as described above, we have observed it for GaMe₃. However, our new polymorph falls into the same structural category, being characterized by long methyl bridges, as the known tetragonal structures; whereas the latter are pseudo-tetrameric, the new polymorph is pseudo-polymeric, and the energy difference between the two forms is presumably small. The observations by previous workers concerning trimethylindium suggest that it may be possible to observe transitions *between* structural types by varying the conditions of temperature and/or pressure. With this in mind, we have investigated the energetic differences between layered and pseudo-tetrameric polymorphs of BMe₃, GaMe₃, and InMe₃ using plane wave density functional theory (DFT). Previous work in our research groups has shown plane wave DFT calculations to be a

very successful and useful tool to investigate (a) polymorphic transitions which occur in small organic systems under the application of high pressure,^{40,41} (b) the properties of hydrogen bonds,⁴² and (c) crystal disorder in PbCp₂.⁴³ Our calculations on GaMe₃ used Mitzel's coordinates for the tetragonal polymorph, as this is most directly comparable with InMe₃.

The results of the calculations are shown in Table 3. Bond and contact distances tend to be overestimated at this level of theory, an effect which carries through to the unit cell dimensions, which are also overestimated. Of course, the C–H bond lengths are all ca. 0.1 Å longer than those obtained experimentally, but most of this disagreement arises from the systematic shortening of these parameters when derived from X-ray data. Nevertheless, both the present results and those of our previous work in this area show that structural trends (for example, in a set of bond lengths within a structure) are reliably reproduced. So too are relative energies. In all three cases, the lower energy structure corresponds to the experimentally observed polymorph. The calculated tetragonal polymorph of BMe₃ has a very long B···C "contact" of 3.715 Å, exceeding even the related distances in either GaMe₃ or InMe₃. Modeling suggests that shortening this contact to a more reasonable distance between 3.0 and 3.3 Å begins to incur repulsive H···H interactions between methyl groups of less than twice the van der Waals radius of H (2.4 Å). This does not occur in the heavy-atom derivatives because of their longer metal-to-carbon bonds. Conversely, the short B–C bond enables the electron deficiency of the boron to be relieved by hyperconjugation between the empty 2p orbital on the boron and the out-of-plane C–H bonds of the methyl groups, a circumstance supported both experimentally and by the results of these calculations (see above). Presumably the reverse of this argument explains the preference for the methyl-bridged structures by the heavier group 13 trimethyls. As the experimental structure of GaEt₃ shows, hyperconjugation is also possible in these systems, although it is notable that the shortest calculated interplanar M···C distances in the *C2/c* polymorphs become shorter along the series B > Ga > In. There is also a tendency along this series for the methyl groups to rotate about the C–metal bond away from the conformation described above for BMe₃.

Mitzel et al. performed calculations at the MP2/TZVP level on an isolated pair of GaMe₃ molecules in which a Ga···C bridging interaction measuring 3.206 Å was found to have an energy of 11.4 kJ mol⁻¹. Of this, only 3.4 kJ mol⁻¹ was ascribable to electrostatic forces; the remainder arose from a dispersion interaction (7.5 kJ mol⁻¹), an ionic correlation contribution (4.2 kJ mol⁻¹), from a reduction in intramolecular correlation on approach of two molecules), and a negative repulsive term (–3.8 kJ mol⁻¹).² The total interaction energy is rather similar to that of a weak hydrogen bond. The energy differences between polymorphs are of the same order of magnitude as the terms given above. The two

(40) Allan, D. R.; Clark, S. J.; Ibberson, R. M.; Parsons, S.; Pulham, C. R.; Sawyer, L. *Chem. Commun.* **1999**, 751.

(41) Allan, D. R.; Clark, S. J.; Dawson, A.; McGregor, P. A.; Parsons, S. *Acta Crystallogr.* **2002**, B58, 1018.

(42) (a) Morrison, C. A.; Siddick, M. M. *Chem. Eur. J.*, in press. (b) Wilson, C. C.; Morrison, C. A. *Chem. Phys. Lett.* **2002**, 362, 85.

(43) Morrison, C. A.; Wright, D. S.; Layfield, R. A. *J. Am. Chem. Soc.* **2002**, 124, 6775.

Table 3. Calculated Solid State Structural Parameters for Polymorphs of MMe₃ (M = B, Ga, In)^a

model	BMe ₃			GaMe ₃			InMe ₃		
	obsd C2/c	calcd C2/c	calcd P4 ₂ /n	obsd P4 ₂ /n ²	calcd C2/c	calcd P4 ₂ /n	obsd P4 ₂ /n ⁴	calcd C2/c	calcd P4 ₂ /n
a/Å	6.3473(9)	6.6648	13.0536	12.9532(3)	7.3159	13.4392	13.2166(11)	7.4839	14.0377
b/Å	10.9284(16)	11.2908	13.0536	12.9532(3)	12.1062	13.4392	13.2166(11)	12.5123	14.0377
c/Å	14.224(2)	14.6057	6.371433	6.2588(1)	14.7211	6.4588	6.4039(9)	14.6060	7.1097
β/deg	91.099(3)	90.002	90.0	90	88.536	90	90	88.973	90
M1–C1/Å	1.5548(11)	1.554	1.556	1.952(3)	1.971	1.973	2.136(13)	2.192	2.180
M1–C2/Å	1.5565(11)	1.558	1.556	1.962(2)	1.970	1.985	2.179(12)	2.185	2.211
M1–C3/Å	1.5541(12)	1.555	1.554	1.958(3)	1.972	1.977	2.121(14)	2.190	2.189
M1···C2'/Å ^b			3.715	3.149(3)		3.266	3.083(12)		3.181
M1···C3'/Å ^b			4.009	3.647(3)		3.848	3.558(15)		4.091
C1–M1–C2/deg	120.06(7)	119.5	119.6	119.72(14)	119.2	120.4	119.7(5)	119.0	118.2
C2–M1–C3/deg	120.09(7)	120.0	121.5	121.29(14)	120.3	119.1	116.8(5)	119.8	118.3
C1–M1–C3/deg	119.86(7)	120.5	120.5	118.85(13)	120.5	120.3	123.5(5)	121.1	122.9
M1···C2'–M1'/deg ^b			170.0	167.23(12)		168.9	168.1(6)		167.8
M1···C3'–M1'/deg ^b			164.2	164.67(13)		162.8	166.2(6)		164.0
M1···C1'/deg ^c	3.7908(13)	3.826	3.758					3.525	
E _{tot} /eV		–5547.369 2869	–5547.224 9200		–21 406.543 6047	–21 406.841 8865		–17 435.088 4853	–17 435.951 7724
E _{rel} /kJ mol ^{–1}		0	+1.7		+3.6	0.0		+10.4	0.0

^a The structures in C2/c are analogous to the observed layered structure of BMe₃ (this should not be confused with the monoclinic polymorph of GaMe₃, which is also described in this paper). Structures in P4₂/n are based on the pseudo-tetrameric structure of InMe₃. All structures have Z = 8 and α = γ = 90°. E_{tot} refers to one unit cell. E_{rel} is the energy per molecule relative to the lower energy polymorph, which is given the value E_{rel} = 0. ^b Tetragonal structures only; these are the distances and angles involving the methyl bridges. ^c Monoclinic structures only; this is the shortest interplanar M···C distance. Primes refer to symmetry-equivalent atoms; refer to Table 2 for full details.

optimized structures of BMe₃ both consist of very weakly interacting molecules, and not surprisingly, the energy difference between them is small. The corresponding energy difference between the polymorphs of InMe₃ is greater than for GaMe₃, consistent with the shorter bridges formed in InMe₃. Enthalpy differences between real polymorphs usually fall in the range 0–10 kJ mol^{–1} and probably do not exceed 25 kJ mol^{–1}.⁷ Therefore, the results of these calculations show it to be quite possible that the unstable phase of InMe₃ observed by Amma and Rundle had the BMe₃ structure.

Distortions Caused by Methyl Bridging. When an atom forms a strong contact, any other bonds that it forms are generally weakened as a result. In our structure of GaMe₃ the primary Ga–C distances span the range 1.956(3)–1.970(3) Å; the average distance in the gaseous molecule, deduced by electron diffraction,²⁹ is 1.967(2) Å. The carbon atom involved in the shortest intermolecular contact (C5···Ga1 = 3.096(3) Å) also makes the longest primary C–Ga bond, but there is no discernible relationship between the other bond and contact lengths in the structure. The calculations on tetragonal InMe₃ show that the In–C bond length involving the carbon atom making the stronger secondary contact is 0.03 Å longer than the bond involving the carbon atom, which is not involved in bridging. The relative lengthening of the In–C bond involving the carbon that makes the longer contact is much less, viz. 0.009 Å.⁴⁴ The corresponding figures for tetragonal GaMe₃ are 0.012 and 0.004 Å, in excellent agreement with the experimental values. The relative lengthening of the bond involving the more strongly bridging methyl group which has been discussed by several authors^{2–5} thus appears to be a genuine effect, even though the differences observed crystallographically tend to teeter on the brink of statistical insignificance.

Bond angles are somewhat “softer” interactions than bond lengths, and all structures exhibit C–M–C angles which differ from 120°. Where a significant deviation occurs, the largest bond angle is invariably the one not involving the carbon atom making the stronger intermolecular contact. There is no deviation from planarity in the molecular MC₃ unit in any of the crystal structures. In contrast, the very strong bridging contacts established in monomeric aluminum derivatives containing bulky groups, such as Al(CH₂Ph)₃⁴⁵ and Al-*t*-Bu₃,⁴⁶ have been observed to lift the Al atom out of the plane formed by the three directly bound carbon atoms by up to 0.4 Å.

Acknowledgment. We thank the EPSRC for research funding and for an Advanced Fellowship to T.M.G. and the Royal Society for a University Research Fellowship to C.A.M.

Supporting Information Available: Tables giving crystallographic data for BMe₃, GaMe₃, and TlMe₃ and optimized theoretical coordinates for polymorphs of BMe₃, GaMe₃, and InMe₃. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM0300272

(44) The experimentally observed differences for these In–C bond lengths are 0.041(18) and –0.015(19) Å, respectively; neither of these is statistically significant.

(45) Rahman, A. F. M. M.; Siddiqui, K. F.; Oliver, J. P. *Organometallics* **1982**, *1*, 881.

(46) Downs, A. J.; Greene, T. M.; Parsons, S.; Taylor, R. A. Unpublished results.