

# Chelate-Controlled Synthesis of *rac*- and *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>ZrCl<sub>2</sub>

Matthew D. LoCoco and Richard F. Jordan\*

Department of Chemistry, The University of Chicago, 5735 South Ellis Avenue,  
Chicago, Illinois 60637

Received September 22, 2003

The reaction of the chelated bis-amide complex Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}Cl<sub>2</sub>(THF)<sub>2</sub> (**2**) with Li<sub>2</sub>[Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>] yields *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} (*meso*-**3**) in >98% yield. In contrast, the reaction of Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>}Cl<sub>2</sub>(THF)<sub>2</sub> (**4**) or the related mono-THF adduct Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>}Cl<sub>2</sub>(THF) (**5**) with Li<sub>2</sub>[Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>] yields *rac*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>} (*rac*-**6**) in quantitative NMR yield and 89% isolated yield. X-ray crystallographic analyses show that the Zr{RN(CH<sub>2</sub>)<sub>3</sub>NR} chelate ring in *rac*-**6** has a pronounced twist conformation, while that in *meso*-**3** has a flatter, envelope conformation. It is proposed that the conformations of the Zr{RN(CH<sub>2</sub>)<sub>3</sub>NR} chelate rings in the stereodetermining transition states for addition of the second Cp<sup>-</sup> ring in these reactions are similar to those in the metallocene products and control the diastereoselectivity. *meso*-**3** and *rac*-**6** are converted to *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>ZrCl<sub>2</sub> (*meso*-**1**) and *rac*-**1**, respectively, by reaction with HCl in Et<sub>2</sub>O.

## Introduction

The application of structurally complex *ansa*-zirconocenes, particularly bis-indenyl systems, as stereoselective catalysts has been studied extensively.<sup>1</sup> This field would be advanced significantly by the development of efficient syntheses of *rac*-metallocenes derived from simple, easily accessible, bis-Cp<sup>-</sup> ligands. In this regard, the synthesis of *rac*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>ZrCl<sub>2</sub> (*rac*-**1**) has attracted considerable interest.<sup>2</sup> This metallocene has been prepared by salt elimination and amine elimination reactions, which produce *rac*/*meso* mixtures,<sup>3,4</sup> and by photochemical isomerization of *rac*-**1**/*meso*-**1** mixtures.<sup>5</sup> To date, the most efficient route to **1**, developed by Brintzinger and co-workers, comprises reaction of Sn(NMe<sub>2</sub>)<sub>4</sub> with Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub> to yield a silastannaindacene, followed by transmeta-

lation with ZrCl<sub>4</sub> to produce *rac*-**1** (>98% de, 60% overall isolated yield).<sup>6</sup> Here we describe highly stereoselective syntheses of *rac*-**1** and *meso*-**1**.

We recently reported a highly selective, "chelate controlled" synthesis of *rac*-*ansa*-bis(indenyl) zirconocenes, which is based on the reaction of lithium *ansa*-bis(indenyl) salts with the chelated bis-amide compound Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}Cl<sub>2</sub>(THF)<sub>2</sub> (**2**) and is shown in Scheme 1.<sup>7</sup> This procedure works well for *rac*-(SBI)Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} and 2-Me-substituted and 2-Me-4-aryl-substituted derivatives thereof, as well as for *rac*-(EBI)-Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}.<sup>7</sup>

The key to Scheme 1 is the use of the [PhN(CH<sub>2</sub>)<sub>3</sub>NPh]<sup>2-</sup> ligand to control diastereoselectivity. The Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} chelate rings in **2** and several (C<sub>5</sub>R<sub>5</sub>)<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} metallocenes adopt twist conformations which place the N-Ph groups on opposite sides of the N-Zr-N plane, as shown in Scheme 1.<sup>7</sup> This arrangement complements the *rac*-metallocene structure but strongly disfavors the *meso* isomer. The mechanism of [X(indenyl)<sub>2</sub>]<sup>2-</sup> substitution of the chloride and THF ligands of **2** is unknown and may be complicated by ion-pairing, solvation, and other factors.<sup>3</sup> However, the simple model in Scheme 1 provides useful insights. We postulate that, while rotation around the Zr-(indenyl centroid) in the mono-indenyl intermediate **A** can occur, the bridge (X) must be close to the "back" position in the transition state for the second indenyl

\* Corresponding author.

(1) (a) Brintzinger, H. H.; Fischer, D.; Mülhaupt, R.; Rieger, B.; Waymouth, R. M. *Angew. Chem., Int. Ed. Engl.* **1995**, *34*, 1143. (b) Resconi, L.; Cavallo, L.; Fait, A.; Piemontesi, F. *Chem. Rev.* **2000**, *100*, 1253. (c) Hoveyda, A. H.; Morken, J. P. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1262. (d) Halterman, R. L. In *Synthesis of Chiral Titanocene and Zirconocene Dichlorides*; Togni, A., Halterman, R. L., Eds.; Metallocenes: Synthesis, Reactivity, Applications: Wiley-VCH: Weinheim, 1998; Vol. 1, Chapter 8.

(2) For polymerization studies using **1** see: (a) Suzuki, N.; Yamaguchi, Y.; Fries, A.; Mise, T. *Macromolecules* **2000**, *33*, 4602. (b) Mise, T.; Miya, S.; Yamazaki, H. *Chem. Lett.* **1989**, 1853. (c) Röhl, W.; Brintzinger, H. H.; Rieger, B.; Zolk, R. *Angew. Chem., Int. Ed. Engl.* **1990**, *29*, 279. For other examples of metallocenes based on *ansa*-bis(3-R-C<sub>5</sub>H<sub>3</sub>) ligands see: (d) Sutton, S. C.; Nantz, M. H.; Parkin, S. R. *Organometallics* **1993**, *12*, 2248. (e) Collins, S.; Hong, Y.; Taylor, N. J. *Organometallics* **1990**, *9*, 2695. (f) Chen, Z.; Halterman, R. L. *Organometallics* **1994**, *13*, 3932. (g) Schnutenhaus, H.; Brintzinger, H. H. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 777. (h) Kuntz B. J.; Ramachandran, R.; Taylor N. J.; Guan, J.; Collins, S. *J. Organomet. Chem.* **1995**, *497*, 133.


(3) Wiesenfeldt, H.; Reinmuth, A.; Barsties, E.; Evertz, K.; Brintzinger, H. H. *J. Organomet. Chem.* **1989**, *369*, 359.

(4) Diamond, G. M.; Jordan, R. F.; Petersen, J. F. *Organometallics* **1996**, *15*, 4045.

(5) Schmidt, K.; Reinmuth, A.; Rief, U.; Diebold, J.; Brintzinger, H. H. *Organometallics* **1997**, *16*, 1724.

(6) (a) Hüttenhofer, M.; Weeber, A.; Brintzinger, H. H. *J. Organomet. Chem.* **2002**, *663*, 58. (b) Hüttenhofer, M.; Schaper, F.; Brintzinger, H. H. *Angew. Chem., Int. Ed.* **1998**, *37*, 2268.

(7) (a) Zhang, X.; Zhu, Q.; Guzei, I. A.; Jordan, R. F. *J. Am. Chem. Soc.* **2000**, *122*, 8093. (b) LoCoco, M. D.; Jordan, R. F. *Abstracts of Papers*, 225th ACS National Meeting, New Orleans, LA, March 23–27, 2003, INOR-723. (c) SBI = Me<sub>2</sub>Si(indenyl)<sub>2</sub>, EBI = 1,2-bis(indenyl)ethane. (d) X-ray crystallographic analyses show that the Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} chelate rings in Cp<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}, *rac*-(SBI)-Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}, and *rac*-(Me<sub>2</sub>Si(2-Me-4,5-benz-1-indenyl)<sub>2</sub>)Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} adopt twist conformations.<sup>7a,b</sup>


Scheme 1<sup>a</sup>

addition, as required in the *ansa*-metallocene product. As the second indenyl adds, N–Ph/indenyl steric interactions block formation of the *meso* product by path (ii), and the *rac* product forms by path (i). Significant N–Ph/indenyl steric interactions would arise on the crowded side of the *meso* isomer (and transition state leading thereto) unless the Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} chelate ring adopted a different conformation, which is unfavorable.<sup>7,8</sup>

Application of this strategy to the synthesis of **1** led to a surprising result. The reaction of Li<sub>2</sub>[Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>] with **2** in Et<sub>2</sub>O yields pure *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} (*meso*-**3**) quantitatively (Scheme 2)! The <sup>1</sup>H NMR spectrum of *meso*-**3** contains two sets of phenyl resonances and two Si–Me resonances, which establishes the *meso* stereochemistry. Reaction of *meso*-**3** with HCl (1 M Et<sub>2</sub>O solution) in benzene yields *meso*-**1** in >95% NMR yield. The use of 4 equiv of HCl in Scheme 2 facilitates the identification of *meso*-**1** by precipitation of the HNPh(CH<sub>2</sub>)<sub>3</sub>NPhH coproduct as [H<sub>2</sub>NPh(CH<sub>2</sub>)<sub>3</sub>NPhH<sub>2</sub>]Cl<sub>2</sub>.

To probe the stereocontrol mechanism in Scheme 2, the molecular structure of *meso*-**3** was determined by X-ray diffraction (Figure 1). Surprisingly, the Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} ring adopts an envelope conformation in which C(10), C(9), and C(8) are close to the N(1)–Zr(1)–N(2) plane (respective deviations: –0.056, 0.083, 0.148 Å), while C(7) lies 0.63 Å below and C(1) lies 0.56 Å above this plane. This unusual conformation is enforced by the bulky <sup>t</sup>Bu groups, which make close contact with and thus flatten the C(10)–C(8) segment of the chelate.<sup>9</sup> Puckering of the C(8)–C(7)–N(1) seg-

## Scheme 2



ment is required to maintain a normal N(1)–Zr(1)–N(2) angle of 95.3(1)°. The C(10) phenyl group on the crowded side of the molecule lies in the N–Zr–N plane. The steric crowding associated with the <sup>t</sup>Bu groups is manifested in several other ways. The <sup>t</sup>Bu groups are bent 9.5° out of their respective Cp planes, and the Zr–C(<sup>t</sup>Bu) distances (Zr–C(28) 2.693(3) Å; Zr–C(17) 2.782(3) Å) are longer than the other Zr–Cp distances (range 2.49–2.65 Å). Additionally, the two Cp rings are canted such that the angle between the C(27)–C(28) and C(18)–C(17) bond vectors is 24.9°. The Cp rings are rotated around the Zr–centroid axes such that the SiMe<sub>2</sub> bridge is displaced by 16.3° from the back position of the metallocene, and the <sup>t</sup>Bu groups are displaced toward the front of the metallocene by a corresponding amount. This “lateral distortion”<sup>10</sup> is smaller than those in *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr(1,1'-binaphth-2'-ol-2-olato)Cl (27.1°)<sup>5</sup> and *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>ZrMe<sub>2</sub> (24.6°)<sup>11</sup> and appears to be limited by steric crowding between the C(10) phenyl group and the Cp and C(34) methyl groups.<sup>12</sup>

These structural data provide clues to the origin of the surprising selectivity for the *meso* product in Scheme 2. As for Scheme 1, we postulate that steric interactions in the mono-Cp intermediate and the transition state for addition of the second Cp<sup>–</sup> ring are similar to those in the metallocene product. That is, as illustrated in Scheme 3, we postulate that steric interactions force the Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} chelate ring of mono-Cp intermediate **B** into an envelope conformation. Steric interactions between the <sup>t</sup>Bu group of the incoming Cp<sup>–</sup> ligand and the puckered segment of the chelate ring block formation of *rac*-**3** via path (i), so that *meso*-**3** forms via path (ii).

To favor the formation of a *rac*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{RN(CH<sub>2</sub>)<sub>3</sub>NR} metallocene, we sought to modify the controlling [RN(CH<sub>2</sub>)<sub>3</sub>NR]<sup>2–</sup> ligand to increase the steric


(9) Close contacts (Å): H(24C)–H(9B) 2.30, H(9A)–H(31A) 2.43, H(32C)–H(8A) 2.20, H(23A)–H(7A) 2.13, C(10)–H(24C) 2.75.

(10) (a) Burger, P.; Diebold, J.; Gutman, S.; Hund, H. U.; Brintzinger, H. H. *Organometallics* **1992**, *11*, 1319. (b) Burger, P.; Hortman, K.; Brintzinger, H. H. *Makromol. Chem., Macromol. Symp.* **1993**, *66*, 127.

(11) Chirik, P. J.; Henling, L. M.; Bercaw, J. E. *Organometallics* **2001**, *20*, 534.

(12) Close H–H contacts (Å): H(15)–H(29) 2.26, H(15)–H(16) 2.47, H(15)–H(34A) 2.40.

(8) In contrast, metallocene syntheses using Ti binaphtholate or Zr biphenolate compounds that adopt twist conformations gave low yields<sup>a</sup> or *rac/meso* mixtures that were isomerized to *rac* products using TEMPO.<sup>b</sup> (a) Erikson, M. S.; Fronczek, F. R.; McLaughlin, M. L. *J. Organomet. Chem.* **1991**, *415*, 75. (b) Damrau, H.-R.; Royo, E.; Obert, S.; Schaper, F.; Weeber, A.; Brintzinger, H.-H. *Organometallics* **2001**, *20*, 5258.




**Figure 1.** Molecular structure and corresponding space-filling diagram of *meso*- $\text{Me}_2\text{Si}(3\text{-}^t\text{Bu-C}_5\text{H}_3)_2\text{Zr}\{\text{PhN}(\text{CH}_2)_3\text{NPh}\}$  (*meso*-**3**). H atoms are omitted in the ORTEP view. Bond distances (Å): Zr–N(1) 2.076(3), Zr–N(2) 2.176(3), Zr–{C(16)–C(20) centroid} 2.328(3), Zr–{C(25)–C(29) centroid} 2.308(3). Bond angles (deg): centroid–Zr–centroid 124.2(9).


crowding between the Cp-<sup>t</sup>Bu and N–R groups, to provide a stronger preference for the chelate twist conformation in the metallocene product (and presumably the transition state for the second Cp<sup>−</sup> addition). This can be achieved by replacing the N–Ph groups with N–SiMe<sub>3</sub> groups, which feature greater three-dimensional steric bulk. The reaction of  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}\text{Cl}_2(\text{THF})_2$  (**4**)<sup>13</sup> or the related mono-THF adduct  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}\text{Cl}_2(\text{THF})$  (**5**) with  $\text{Li}_2\{\text{Me}_2\text{Si}(3\text{-}^t\text{Bu-C}_5\text{H}_3)_2\}$  in THF yields pure *rac*- $\text{Me}_2\text{Si}(3\text{-}^t\text{Bu-C}_5\text{H}_3)_2\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  (*rac*-**6**) in quantitative NMR yield and 89% isolated yield (Scheme 4). The <sup>1</sup>H NMR spectrum of *rac*-**6** contains singlets for the <sup>t</sup>Bu, SiMe<sub>2</sub>, and SiMe<sub>3</sub> groups, which confirms the *rac* stereochemistry. *rac*-**6** is cleanly converted to *rac*-**1** by reaction with 6 equiv of HCl in Et<sub>2</sub>O. The 3-fold excess of HCl is

(13) (a) Freidrich, S.; Gade, L. H.; Scowen, I. J.; McPartlin, M. *Organometallics* **1995**, *14*, 5344. (b) Freidrich, S.; Gade, L. H.; Trosch, D. J. M.; Scowen, I. J.; McPartlin, M. *Inorg. Chem.* **1999**, *38*, 5295.

## Scheme 3




## Scheme 4



necessary because the  $\text{Me}_3\text{SiNH}(\text{CH}_2)_3\text{NHSiMe}_3$  coproduct reacts with HCl to yield  $[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_3]\text{Cl}_2$  and  $\text{ClSiMe}_3$ .

The molecular structure of *rac*-**6** is shown in Figure 2. *rac*-**6** has approximate *C*<sub>2</sub> symmetry with the *C*<sub>2</sub> axis lying along the Zr–C(12) vector. In contrast to *meso*-**3**, the  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  chelate ring has the familiar twist conformation,<sup>7</sup> which places the N–SiMe<sub>3</sub> groups in the sterically open quadrants of the *rac*-metallocene and minimizes SiMe<sub>3</sub>/<sup>t</sup>Bu steric interactions.<sup>14,15</sup> We propose that the twist conformation

(14) (a) Deviations (Å) of C and Si atoms from the N(1)–Zr–N(2) plane: Si(2) 0.88, C(11) −0.87, C(12) 0.10, C(13) 0.71, Si(1) −0.61. (b) The conformation of the  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  chelate ring in *rac*-**6** is similar to that in **4**; see ref 13b.



**Figure 2.** Molecular structure and corresponding space-filling diagram of *rac*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>} (*rac*-6). H atoms are omitted from the ORTEP view. Bond distances (Å): Zr–N(1) 2.114(2), Zr–N(2) 2.103(2), N(1)–Si(1) 1.725(2), N(2)–Si(2) 1.737(2), Zr–{C(1)–C(5) centroid} 2.328(3), Zr–{C(6)–C(10) centroid} 2.340(3). Bond angles (deg): N(1)–Zr–N(2) 92.63(8), centroid–Zr–centroid 122.5(9).

dictates the formation of *rac*-6 in a manner analogous to Scheme 1.

### Conclusions

This work expands the scope of the “chelate-controlled” approach to metallocene synthesis to include simple *ansa*-bis-Cp systems. The present results suggest that suitable modification of the controlling bis-amide ligand may enable the synthesis of a wide range of *rac*-metallocenes by this method. These studies also provide new insights into the factors that control diastereoselectivity in *ansa*-metallocene formation.

### Experimental Section

**General Procedures.** All reactions were performed under nitrogen or vacuum using standard Schlenk techniques or in

(15) In contrast to *meso*-3, *rac*-6 does not display a significant lateral distortion. However, the Zr–N bond distances, N–Zr–N bond angle, displacement of the <sup>t</sup>Bu groups from the Cp planes, and the trend in Zr–C(Cp) distances in *rac*-6 are similar to the data for *meso*-3. The angle between the C(2)–C(3) and C(10)–C(9) bond vectors is 9.9°.

a nitrogen-filled drybox. Nitrogen was purified by passage through columns containing activated molecular sieves and Q-5 oxygen scavenger. Pentane, hexanes, toluene, and benzene were distilled from sodium/benzophenone or purified by passage through columns of activated alumina and BASF R3-11 oxygen removal catalyst. Benzene-*d*<sub>6</sub> was distilled from sodium/benzophenone. The compounds Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>,<sup>3</sup> Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}Cl<sub>2</sub>(THF)<sub>2</sub>,<sup>7a,b</sup> Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>}<sub>2</sub>,<sup>13a</sup> and Zr{Me<sub>3</sub>SiN(CH<sub>2</sub>)<sub>3</sub>NSiMe<sub>3</sub>}Cl<sub>2</sub>(THF)<sub>2</sub><sup>13a</sup> were prepared by literature procedures.

Elemental analyses were performed by Midwest Microlabs. ESI-mass spectra were obtained with an Agilent 1100 instrument using direct injection via a syringe pump (ca. 10<sup>−6</sup> M solutions). Good agreement between observed and calculated isotope patterns was observed in all cases. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker AC-500 or AC-400 spectrometer in flame-sealed or Teflon-valved tubes at ambient probe temperatures. <sup>1</sup>H and <sup>13</sup>C chemical shifts are reported relative to SiMe<sub>4</sub> and were determined by reference to the residual <sup>1</sup>H and <sup>13</sup>C solvent resonances. Coupling constants are given in Hz.

**Li<sub>2</sub>[Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>].<sup>3</sup>** A Schlenk flask was charged with a solution of Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub> (2.75 g, 9.03 mmol) in Et<sub>2</sub>O (50 mL). The solution was cooled to −78 °C, and a solution of n-BuLi in hexanes (2.5 M, 7.22 mL, 18 mmol) was added by syringe. The mixture was stirred for 4 h at −78 °C, warmed to 0 °C in an ice bath, and then stirred overnight while being allowed to warm gradually to room temperature. The volatiles were removed under vacuum to yield a white solid. Toluene (50 mL) was added by cannula transfer. The slurry was stirred for 30 min and filtered through a medium-porosity glass frit (10–20 μm pore size) to yield a white solid. The solid was dried under vacuum overnight (2.55 g, 89%). <sup>1</sup>H NMR (THF-*d*<sub>6</sub>): δ 5.86 (m, 4H, Cp), 5.74 (t, *J* = 3, 2H, Cp), 1.21 (s, 18H, <sup>t</sup>Bu), 0.26 (s, 6H, SiMe<sub>2</sub>).

**meso-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} (meso-3).** A flask was charged with Li<sub>2</sub>[Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>] (0.705 g, 2.22 mmol) and Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh}Cl<sub>2</sub>(THF)<sub>2</sub> (1.18 g, 2.22 mmol). Diethyl ether (100 mL) was added by vacuum transfer at −196 °C. The mixture was warmed to 0 °C and stirred for 6 h and then warmed to room temperature and stirred overnight. The volatiles were removed under vacuum at 25 °C to yield a red solid. Toluene (100 mL) was added by vacuum transfer at −196 °C, and the mixture was warmed to room temperature and stirred for 30 min. The mixture was filtered through a Celite column, and the volatiles were removed from the filtrate under vacuum at 25 °C. The remaining red solid was dried under vacuum (1.36 g, 99%). This material, which was spectroscopically pure, was recrystallized from pentane/hexamethyldisiloxane.<sup>16</sup> <sup>1</sup>H NMR (C<sub>6</sub>D<sub>6</sub>): δ 7.28 (t, *J* = 8, 2H, *m*-Ph), 7.22 (t, *J* = 8, 2H, *m*-Ph), 6.96 (t, *J* = 8, 1H, *p*-Ph), 6.79 (m, 3H, *o*-Ph and *p*-Ph), 6.47 (d, *J* = 8.0 2H, *o*-Ph), 6.22 (m, 4H, Cp), 6.00 (t, *J* = 3.0, 2H, Cp), 3.57 (t, *J* = 6, 2H, NCH<sub>2</sub>), 3.26 (t, *J* = 6, 2H, NCH<sub>2</sub>), 1.67 (p, *J* = 6, 2H, CH<sub>2</sub>), 0.92 (s, 18H, <sup>t</sup>Bu), 0.60 (s, 3H, SiMe), 0.55 (s, 3H, SiMe). <sup>13</sup>C{<sup>1</sup>H} NMR (THF-*d*<sub>6</sub>): δ 162.9, 160.0, 148.4, 129.2, 128.5, 122.6, 120.8, 120.4, 118.3, 117.4, 111.8, 109.7, 106.0, 53.6, 52.9, 33.7, 31.5, 29.7, −2.6, −5.4; the signal at δ 118.3 is broadened by restricted rotation around an N–Ph bond at room temperature but is sharp at 60 °C. Anal. Calcd for C<sub>35</sub>H<sub>46</sub>N<sub>2</sub>SiZr: C, 68.45; H, 7.54; N, 4.56. Found: C, 68.30; H, 7.52; N, 4.61.

**meso-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>ZrCl<sub>2</sub> (meso-1).** An NMR tube was charged with *meso*-Me<sub>2</sub>Si(3-<sup>t</sup>Bu-C<sub>5</sub>H<sub>3</sub>)<sub>2</sub>Zr{PhN(CH<sub>2</sub>)<sub>3</sub>NPh} (0.023 g, 0.037 mmol), and C<sub>6</sub>D<sub>6</sub> (1.0 mL) was added by syringe. A solution of HCl in Et<sub>2</sub>O (1.0 M, 77.5 μL, 0.077 mmol) was added by syringe. The volatiles were removed under vacuum

(16) For other examples of the use of hexamethyldisiloxane as a recrystallization solvent see: (a) Kloppenburg, L.; Petersen, J. L. *Polyhedron* **1995**, *14*, 69. (b) Strauch, J. W.; Petersen, J. L. *Organometallics* **2001**, *20*, 2623.

immediately, and fresh  $C_6D_6$  was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ . The tube was warmed to room temperature, and a  $^1\text{H}$  NMR spectrum was obtained, which showed that complete conversion to  $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (*meso/rac* 30:1)<sup>3</sup> and  $\text{HNPh}(\text{CH}_2)_3\text{NPh}$  had occurred.  $^1\text{H}$  NMR ( $C_6D_6$ ) for *meso*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$ :  $\delta$  6.84 (t,  $J = 2$ , 2H, Cp), 5.87 (t,  $J = 2$ , 2H, Cp), 5.54 (t,  $J = 2$ , 2H, Cp), 1.45 (s, 18H,  $^i\text{Bu}$ ), 0.29 (s, 3H, SiMe), 0.08 (s, 3H, SiMe).  $^1\text{H}$  NMR ( $C_6D_6$ ) for  $\text{HNPh}(\text{CH}_2)_3\text{NPh}$ :  $\delta$  7.17 (t,  $J = 7$ , 4H, Ph), 6.76 (t,  $J = 7$ , 2H, Ph), 6.46 (d,  $J = 7$ , 4H, Ph), 3.50 (br s, 2H, NH), 2.73 (t,  $J = 7$ , 4H,  $\text{NCH}_2$ ), 1.32 (p,  $J = 7$ ,  $\text{CH}_2$ ).

**Generation of *meso*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (*meso-1*) from *meso-3* and 4 Equiv of HCl.** An NMR tube was charged with *meso*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{Zr}[\text{Ph}(\text{CH}_2)_3\text{NPh}]$  (0.028 g, 0.045 mmol), and  $C_6D_6$  (1.0 mL) was added by syringe. A solution of HCl in  $\text{Et}_2\text{O}$  (1.0 M, 181.0  $\mu\text{L}$ , 0.181 mmol) was added by syringe. The volatiles were removed under vacuum immediately, and fresh  $C_6D_6$  was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ . The tube was warmed to room temperature to yield a slurry of a white precipitate in a yellow supernatant. The supernatant was decanted from the precipitate to yield a clear yellow solution.  $^1\text{H}$  NMR analysis of the solution established that complete conversion to  $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (*meso/rac* 20:1) had occurred. The white solid was isolated and identified as  $[\text{H}_2\text{NPh}(\text{CH}_2)_3\text{NH}_2\text{Ph}]_2\text{Cl}_2$  by NMR and ESI-MS.

**$[\text{H}_2\text{NPh}(\text{CH}_2)_3\text{NH}_2\text{Ph}]_2\text{Cl}_2$ .** A flask was charged with a solution of  $\text{HNPh}(\text{CH}_2)_3\text{NPh}$  (1.12 g, 4.94 mmol) in benzene (25 mL). A solution of HCl in  $\text{Et}_2\text{O}$  (1.0 M, 9.89 mL, 9.89 mmol) was added over a 10 min period by syringe. The mixture was stirred for 1 h at room temperature and filtered to afford a white solid. The solid was washed with hexanes (2  $\times$  20 mL) and benzene (20 mL). The solid was then taken up as a slurry in hexanes (20 mL), and the volatiles were removed under vacuum to yield a white solid.  $^1\text{H}$  NMR ( $D_2O$ ):  $\delta$  7.46 (br m, 6H, Ph), 7.32 (br d,  $J = 7$ , 4H, Ph), 3.43 (t,  $J = 8$ , 4H,  $\text{NCH}_2$ ), 2.03 (p,  $J = 8$ , 2H,  $\text{CH}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $D_2O$ ):  $\delta$  134.0, 130.3, 129.7, 122.0, 47.6, 21.4. ESI-MS:  $[\text{HNPh}(\text{CH}_2)_3\text{NH}_2\text{Ph}]^+$  calcd  $m/z$  227.1, found 227.1.

**$\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}\text{Cl}_2(\text{THF})$  (5).** A flask was charged with  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}_2$  (3.04 g, 5.79 mmol) and  $\text{ZrCl}_4$  (1.35 g, 5.79 mmol). Toluene (100 mL) was added, and the mixture was heated to  $60\text{ }^\circ\text{C}$  for 3 days. The volatiles were removed under vacuum, and the resulting white solid was washed with pentane (75 mL) and dried under vacuum (4.20 g, 95%). The solid was dissolved in THF (40 mL). The bis-THF adduct  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}_2\text{Cl}_2(\text{THF})_2$  (4) can be isolated from this solution by crystallization at  $-35\text{ }^\circ\text{C}$  as described by Gade<sup>13a,b</sup> or by removal of the volatiles under vacuum and vacuum-drying the sample for a few minutes. Drying under vacuum overnight yielded  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}_2\text{Cl}_2(\text{THF})$  as a white solid (5.01 g, 95%).  $^1\text{H}$  NMR ( $C_6D_6$ ):  $\delta$  3.72 (m, 4H,  $\text{OCH}_2$ ), 3.23 (t,  $J = 5$ , 4H,  $\text{NCH}_2\text{CH}_2$ ), 1.74 (p,  $J = 5$ , 2H,  $\text{NCH}_2\text{CH}_2$ ), 1.25 (m, 4H,  $\text{OCH}_2\text{CH}_2$ ), 0.31 (s, 18H, SiMe<sub>3</sub>).

***rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  (*rac-6*).** A flask was charged with  $\text{Li}_2[\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2]$  (0.853 g, 2.69 mmol) and  $\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}_2\text{Cl}_2(\text{THF})$  (1.21 g, 2.69 mmol), and THF (100 mL) was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ . The mixture was warmed to  $0\text{ }^\circ\text{C}$  in an ice bath and then stirred for 20 h while the bath was allowed to warm to room temperature. The volatiles were removed under vacuum at  $25\text{ }^\circ\text{C}$  to yield a yellow solid. Benzene (100 mL) was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ , and the mixture was warmed to room temperature and stirred for 3 h. The mixture was filtered through a medium-porosity glass frit, and the volatiles were removed from the filtrate under vacuum at  $25\text{ }^\circ\text{C}$ . The resulting yellow solid was dried under vacuum (1.46 g, 89%). This material, which was spectroscopically pure, was recrystallized from pentane.  $^1\text{H}$  NMR ( $C_6D_6$ ):  $\delta$  6.85 (t,  $J = 2$ , 2H, Cp), 5.91 (t,  $J = 2$ , 2H, Cp), 5.84 (t,  $J = 2$ , 2H, Cp), 3.32 (dt,  $J = 16$ , 4; 2H,  $\text{NCH}_2$ ), 3.04 (m, 2H,  $\text{NCH}_2$ ), 1.23 (s, 18H,

**Table 1. Summary of Crystal Data for *meso-3* and *rac-6***

	<i>meso-3</i>	<i>rac-6</i>
formula	$\text{C}_{35}\text{H}_{46}\text{N}_2\text{SiZr}$	$\text{C}_{29}\text{H}_{54}\text{N}_2\text{Si}_3\text{Zr}$
cryst syst	monoclinic	monoclinic
space group	$P2_1/n$	$P2_1/n$
<i>a</i> (Å)	10.324(2)	14.712(7)
<i>b</i> (Å)	18.201(3)	11.872(6)
<i>c</i> (Å)	16.753(3)	19.306(9)
$\beta$ (deg)	100.203(2)	104.373(8)
<i>V</i> (Å <sup>3</sup> )	3094.8(8)	3266(3)
<i>Z</i>	4	4
$\mu$ (mm <sup>-1</sup> )	0.420	0.466
cryst dimens (mm)	$0.2 \times 0.2 \times 0.2$	$0.40 \times 0.25 \times 0.25$
cryst color, habit	red, fragment	yellow, fragment
<i>T</i> (K)	100	135
diffractometer	Bruker Smart Apex	Bruker Smart Apex
radiation, $\lambda$ (Å)	Mo K $\alpha$ , 0.71073	Mo K $\alpha$ , 0.71073
$\theta$ range (deg)	2.24–25.03	2.03–25.02
data collected: <i>h</i> ; <i>k</i> ; <i>l</i>	$\pm 12$ ; $-21$ ; $19$ ; $-18$ ; $19$	$\pm 17$ ; $\pm 14$ ; $\pm 22$
no. of reflns	19 612	23 087
no. of indep reflns	5476 (0.0322)	5755 (0.0252)
$(R_{\text{int}})^a$		
abs corr	SADABS	SADABS
max., min. transmn	1.00, 0.854	1.00, 0.852
no. of data/restraints/params	5467/0/360	5755/0/330
<i>R</i> indices ( $I > 2\sigma(I)$ ) <sup>b</sup>	$R_1 = 0.0456$ $wR_2 = 0.0899$	$R_1 = 0.0331$ $wR_2 = 0.0817$
<i>R</i> indices (all data) <sup>b</sup>	$R_1 = 0.0465$ $wR_2 = 0.0904$	$R_1 = 0.0347$ $wR_2 = 0.0826$

<sup>a</sup>  $R_{\text{int}} = \sum |F_o^2 - \langle F_o^2 \rangle| / \sum F_o^2$ . <sup>b</sup>  $R_1 = \sum ||F_o| - |F_c|| / \sum |F_o|$ ;  $wR_2 = \sum [w(F_o^2 - F_c^2)^2] / \sum [w(F_o^2)^2]^{1/2}$ , where  $w = 1 / [\sigma^2(F_o^2) + (aP)^2 + bP]^{-1}$ .

$^i\text{Bu}$ , 1.00 (br m, 2H,  $\text{CH}_2$ ), 0.47 (s, 6H, SiMe<sub>2</sub>), 0.27 (s, 18H, SiMe<sub>3</sub>).  $^{13}\text{C}\{^1\text{H}\}$  NMR:  $\delta$  148.8, 116.1, 111.3, 107.9, 106.5, 46.2, 33.7, 31.3, 28.5, 3.6,  $-4.3$ . Anal. Calcd for  $\text{C}_{29}\text{H}_{54}\text{N}_2\text{Si}_3\text{Zr}$ : C, 57.45; H, 8.98; N, 4.57. Found: C, 57.19; H, 8.97; N, 4.57.

***rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (*rac-1*).** **NMR Scale.** An NMR tube was charged with *rac*- $\text{Me}_2\text{Si}(1\text{-C}_5\text{H}_3\text{-}3\text{-}^i\text{Bu})_2\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  (0.0274 g, 0.0451 mmol) and  $C_6D_6$  (1 mL). A solution of HCl in  $\text{Et}_2\text{O}$  (1.0 M, 271  $\mu\text{L}$ , 0.271 mmol) was added by syringe. The volatiles were removed under vacuum to yield a yellow solid. THF-*d*<sub>8</sub> (1 mL) was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ , and the tube was warmed to room temperature to yield a slurry of a white solid in a yellow supernatant. The supernatant was decanted off and transferred to a new NMR tube. NMR analysis confirmed that quantitative conversion to *rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  had occurred.  $^1\text{H}$  NMR of *rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (THF-*d*<sub>8</sub>):  $\delta$  6.74 (t,  $J = 3$ , 2H), 6.06 (t,  $J = 3$ , 2H, Cp), 5.92 (t,  $J = 3$ , 2H, Cp), 1.32 (s, 18H,  $^i\text{Bu}$ ), 0.68 (s, 6H, SiMe<sub>2</sub>). The white solid was washed with benzene (2 mL) and hexanes (2 mL) and dried under vacuum. NMR and ESI-MS analysis established that the white solid was  $[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_3]\text{Cl}_2$ . Data for  $[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_3]\text{Cl}_2$ :  $^1\text{H}$  NMR ( $D_2O$ ):  $\delta$  3.14 (t,  $J = 8$ , 4H,  $\text{NCH}_2$ ), 2.10 (p,  $J = 8$ , 2H,  $\text{CH}_2$ ).  $^{13}\text{C}\{^1\text{H}\}$  NMR ( $D_2O$ ):  $\delta$  36.59, 25.41. ESI-MS (DMSO):  $[\text{H}_3\text{N}(\text{CH}_2)_3\text{NH}_2]^+$  calcd  $m/z$  75.1, found 75.2.

***rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{ZrCl}_2$  (*rac-1*).** **Preparative Scale.** A flask was charged with *rac*- $\text{Me}_2\text{Si}(3\text{-}^i\text{Bu-C}_5\text{H}_3)_2\text{Zr}\{\text{Me}_3\text{SiN}(\text{CH}_2)_3\text{NSiMe}_3\}$  (1.554 g, 2.563 mmol) and benzene (80 mL). A solution of HCl in  $\text{Et}_2\text{O}$  (1.0 M, 15.38 mL, 15.38 mmol) was added by syringe over a 10 min period. The volatiles were removed under vacuum to yield a yellow solid. Benzene (50 mL) was added by vacuum transfer at  $-196\text{ }^\circ\text{C}$ , and the mixture was warmed to room temperature and stirred for 45 min. The mixture was filtered through a medium-porosity glass frit. The volatiles were removed from the filtrate under vacuum. The remaining yellow solid was dried under vacuum (1.00 g, 84%).  $^1\text{H}$  NMR ( $C_6D_6$ ):  $\delta$  6.72 (t,  $J = 3$ , 2H), 5.69 (t,  $J = 3$ , 2H, Cp), 5.65 (t,  $J = 3$ , 2H, Cp), 1.40 (s, 18H,  $^i\text{Bu}$ ), 0.18

(s, 6H, SiMe<sub>2</sub>). <sup>13</sup>C{<sup>1</sup>H} NMR: δ 152.9, 125.8, 116.2, 110.2, 105.4, 33.6, 30.3, -5.8.

**Reaction of Me<sub>3</sub>SiNH(CH<sub>2</sub>)<sub>3</sub>NHSiMe<sub>3</sub> with 4 Equiv of HCl.** An NMR tube was charged with a solution of Me<sub>3</sub>SiNH-(CH<sub>2</sub>)<sub>3</sub>NHSiMe<sub>3</sub> (0.049 g, 0.23 mmol) in C<sub>6</sub>D<sub>6</sub> (2 mL). A solution of HCl in Et<sub>2</sub>O (1.0 M, 0.9 mL, 0.9 mmol) was added by syringe, and a white precipitate formed. The volatiles were vacuum transferred to a new NMR tube. The white solid was identified as [H<sub>3</sub>N(CH<sub>2</sub>)<sub>3</sub>NH<sub>3</sub>]Cl<sub>2</sub> by NMR and ESI-MS. The presence of ClSiMe<sub>3</sub> in the volatiles was established by <sup>1</sup>H NMR and confirmed by spiking with additional ClSiMe<sub>3</sub>.

**X-ray Structure Determinations.** Crystal data, data collection details, and solution and refinement procedures are summarized in Table 1, and full details are provided in the Supporting Information. Crystals of *meso*-**3** were obtained from a 5:1 pentane/hexamethyldisiloxane solution which was heated

to 80 °C and then allowed to cool to room temperature over a 24 h period. Crystals of *rac*-**6** were obtained from a saturated pentane solution cooled to -35 °C.

**Acknowledgment.** We thank Dr. Ian Steele for assistance with the X-ray crystallographic analyses. This work was supported by the National Science Foundation (CHE-0212210).

**Supporting Information Available:** Additional experimental details and tables of crystallographic data for *meso*-**3** and *rac*-**6**; data are also available for these compounds in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OM034184C