

This article was downloaded by:

On: 29 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

ISOLATION OF THE FIRST ANIONIC FIVE-COORDINATED SILICATES WITH SIX-AND SEVEN-MEMBERED OXYGEN-CONTAINING RINGS

K. C. Kumara Swamy^{ab}; Channareddy Sreelatha^a; Roberta O. Day^a; Joan Holmes^a; Robert R. Holmes^a

^a Department of Chemistry, University of Massachusetts, Amherst, Massachusetts ^b School of Chemistry, University of Hyderabad, Hyderabad, India

To cite this Article Swamy, K. C. Kumara , Sreelatha, Channareddy , Day, Roberta O. , Holmes, Joan and Holmes, Robert R.(1995) 'ISOLATION OF THE FIRST ANIONIC FIVE-COORDINATED SILICATES WITH SIX-AND SEVEN-MEMBERED OXYGEN-CONTAINING RINGS', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 100: 1, 107 — 121

To link to this Article: DOI: 10.1080/10426509508037703

URL: <http://dx.doi.org/10.1080/10426509508037703>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

ISOLATION OF THE FIRST ANIONIC FIVE-COORDINATED SILICATES WITH SIX- AND SEVEN-MEMBERED OXYGEN-CONTAINING RINGS^{1,2}

K. C. KUMARA SWAMY,³ CHANNAREDDY SREELATHA,
ROBERTA O. DAY, JOAN HOLMES, and ROBERT R. HOLMES*

*Contribution from the Department of Chemistry, University of Massachusetts,
Amherst, Massachusetts 01003*

(Received December 10, 1990)


Reaction of PhSi(OMe)_3 with KOMe and naphthalenediol led to the first five-coordinated anionic silicate with an oxygen-containing six-membered ring, $[(\text{C}_{10}\text{H}_6\text{O}_2)_2\text{Si}(\text{C}_6\text{H}_5)][\text{K}, 18\text{-c-}6]$ (**1**). Reaction of the tetracoordinate spirocyclic silane $(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{Si}$ with either KO-*t*-Bu or KF in the presence of 18-crown-6 led to the isolation of the first five-coordinated anionic silicates with seven-membered oxygen-containing rings, $[(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{SiR}][\text{K}, 18\text{-c-}6]$, R = O-*t*-Bu (**2**), F (**3**). ²⁹Si NMR spectroscopy shows that the five-coordinated silicates **2** and **3** slowly disproportionate into four-coordinate silicon compounds and organic species lacking any silicon. Syntheses similar to that for **2** and **3** conducted with other ligands containing the isopropoxy group or use of $\text{Bu}_4\text{N}^+\text{F}^-$ in place of potassium 18-crown-6 imparts less stability to the system and leads directly to the formation of anionic organic products consisting of two biphenolate species hydrogen bonded to each other. In the case of $[(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{H}_3][(\text{n-Bu})_4\text{N}]$ (**5**) and $[(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{H}][\text{K}, 18\text{-c-}6]$, (**6**), X-ray studies substantiated these compositions. Although the structure of **6** was disordered, the detailed hydrogen-bonding scheme was revealed in the anionic structure of **5**. A hydrolytic cleavage mechanism is proposed indicating enhanced reactivity of pentacoordinated anionic silicates relative to their tetracoordinated silane precursors. An X-ray structural analysis reveals a trigonal-bipyramidal geometry for **1** with the six-membered rings located in axial-equatorial sites similar to that found for related pentacoordinated anionic oxysilicates with five-membered-ring systems and isoelectronic oxyphosphoranes that have varying ring compositions. Silicate **1** crystallizes in the monoclinic space group $P2_1/n$ with $a = 9.798$ (1) Å, $b = 20.919$ (4) Å, $c = 18.300$ (4) Å, $\beta = 101.12$ (1)° and $Z = 4$. The biphenolate derivative **5** crystallizes in the triclinic space group $P\bar{1}$ with $a = 9.315$ (6) Å, $b = 9.467$ (3) Å, $c = 21.293$ (6) Å, $\alpha = 78.89$ (2)°, $\beta = 84.89$ (4)°, $\gamma = 80.84$ (4)°, and $Z = 2$. The final conventional unweighted residuals are 0.058 (1) and 0.055 (5).

INTRODUCTION


The synthesis and structural characterization of cyclic phosphoranes with five-membered rings has been well developed.^{4,5} More recently, pentacoordinated phosphorus compounds containing larger rings have been reported, particularly members of the cyclic oxyphosphorane class.^{6–9} Development of analogous series of isoelectronic cyclic anionic five-coordinated silicates has lagged, partly due to the increased difficulty in handling these more reactive species^{2,10–12} and partly due to the lack of recognition of the importance of hypervalent silicon species in reaction mechanisms¹⁰ compared to that in phosphorus chemistry.⁵ Recent studies have shown enhanced reactivity of anionic silicates^{13–20} compared to their tetracoordinate counterparts. Also a preliminary report²¹ on cyclic pentacoordinated silicates

Reprinted with permission from Inorg. Chem., 30, 3126 (1991). Copyright 1991 American Chemical Society.


has illustrated the importance of stereochemical control in nucleophilic displacement reactions brought about by variations in ring constraints.




Similar to oxyphosphoranes containing five-membered rings,⁴ it has been shown that monocyclic and spirocyclic oxyphosphoranes with larger rings invariably position such rings in apical-equatorial sites of a trigonal bipyramid (TBP), e.g., A–D.^{6–9} (references are shown as superscripts). Likewise, analogous studies in silicon chemistry have shown that five-membered rings are located in apical-equatorial positions in a TBP framework for the pentaoxysilicate anions $[(OCMe_2CMe_2O)_2SiOR][K, 18-c-6]$, $R = t\text{-Bu}$ (E)^{11b} and $i\text{-Pr}$ (F),^{11a} and $[(OCMe_2CMe_2O)_2SiOEt][n\text{-BuNH}_3]$ (G),^{11b} while $[(OCMe_2CMe_2O)_2SiOMe][n\text{-BuNH}_3] \cdot MeOH$ (H)^{11b} is square pyramidal with hydrogen bonding implicated as a factor responsible for the formation of the square pyramid. Regarding larger size rings for anionic oxysilicates, there appears to be only one study, that of Kira *et al.*,¹⁷ who reported an allyl derivative containing two seven-membered 2,2'-biphenolate rings. However, the substance was not isolated.



In the present study, we report the isolation and characterization of the first five-coordinated anionic silicates $[(C_{10}H_6O_2)_2Si(C_6H_5)]^-$ (anion of **1**) and $[(C_{12}H_8O_2)_2SiR]^-$, $R = O-t-Bu$ (anion of **2**) and F (anion of **3**), as potassium 18-



crown-6 salts with six- and seven-membered oxygen-containing rings, thus extending the range of known cyclic anionic pentaoxysilicate systems. The latter extension is complimentary to our studies of pentacoordinated anionic fluorosilicates,² which show both diequatorial and axial-equatorial orientation in a TBP for carbon-containing five-membered-ring systems **I** and **J**, respectively, and diequatorial placement of carbon-containing six- and seven-membered rings **K** and **L**, respectively. All are potassium 18-crown-6 salts.



Also included in the present study are reactions of biphenyl orthosilicates leading to anionic organic systems presumably via reactive pentacoordinated anionic silicate intermediates. Their compositions, which are composed of hydrogen-bonded biphenolate ions, were characterized by X-ray analysis, elemental analysis, and solution-state NMR spectroscopy.


EXPERIMENTAL

All the manipulations were carried out under an atmosphere of dry nitrogen. The apparatus was evacuated and flame dried before use. All solvents were dried according to standard procedures and stored under nitrogen. 2-Propanol was dried over aluminum isopropoxide. 2,2'-Biphenol, phenyl trimethoxysilane, potassium *tert*-butoxide (as 1.0 M solution in THF), potassium methoxide, potassium

fluoride, and tetrabutylammonium bromide were commercially available (Aldrich Chemical). Potassium isopropoxide was freshly prepared from KH and 2-propanol.

The NMR spectra were recorded on Varian Associates XL-200 (^1H) and XL-300 FT-NMR spectrometers (^{13}C and ^{29}Si) operating in the FT mode at 200.0, 75.43, and 59.59 MHz, respectively, using internal Me_4Si (^1H and ^{29}Si) and CDCl_3 (^{13}C) as standards. The chemical shifts are reported in ppm with shifts downfield to Me_4Si as positive.

2,2'-Biphenoxydichlorosilane²² (^{29}Si NMR (CH_2Cl_2): -48.9), 2,2'-biphenoxydiisopropoxysilane²² (^{29}Si NMR (CH_2Cl_2): -80.3), and 1,8-dihydroxynaphthalene²³ were prepared by literature methods. The numbering scheme for the biphenyldioxy system is as follows:



Synthesis. Potassium 18-Crown-6 Bis(naphthalene-1,8-diolato)phenylsilicate, $[(\text{C}_{10}\text{H}_6\text{O}_2)_2\text{Si}(\text{C}_6\text{H}_5)][\text{K}, 18\text{-c-6}]$ (**1**). To a mixture of phenyltrimethoxysilane (1.35 g, 6.81 mmol), 18-crown-6 91.80 g, 6.81 mmol), and potassium methoxide (0.65 g, 9.3 mmol) in dry toluene (100 mL) was added 1,8-dihydroxynaphthalene (1.09 g, 6.81 mmol), and the mixture was heated at ca. 70 °C for ½ h to remove methanol by azeotropic distillation. The mixture was then filtered under a nitrogen atmosphere, toluene was removed in vacuo, and the residue was recrystallized from dichloromethane (60 mL) to give **1**: mp 275 °C dec; yield 1.50 g, 41%. ^1H NMR (CDCl_3): 3.44 (s, 24 H, OCH_2), 6.55–7.25 (m, 17 H, H(Ar)). ^{29}Si NMR (CDCl_3): -132.35 . Anal. Calcd for $\text{C}_{38}\text{H}_{44}\text{O}_{10}\text{KSi}$: C, 62.97; H, 5.66. Found: C, 62.71; H, 5.71.

Potassium 18-Crown-6 Bis(biphenyl-2,2'-diolato)-*tert*-butoxysilicate, $[(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{Si}(\text{O-}t\text{-Bu})][\text{K}, 18\text{-c-6}]$ (**2**). To a suspension of spirocyclic bis(biphenyl) orthosilicate (1.56 g, 3.93 mmol) in ~15 mL of THF was added potassium *tert*-butoxide (4.0 mL, 1.0 M solution in THF; 4.0 mmol) dropwise. During the addition, solid orthosilicate started going into solution. After 10 min of stirring, a stoichiometric amount of 18-crown-6 (1.04 g, 3.39 mmol) was added and the contents were stirred overnight. After the reaction mixture was filtered, ~5 mL of CH_2Cl_2 was added to the filtrate, which was then layered with ~10–20 mL of hexane and kept at 0 °C overnight. White crystalline material separated out and was filtered off, washed with hexane (2 × 20 mL), and dried: mp ~165 °C dec; yield 2.3 g, 76%. ^1H NMR (CDCl_3): 6.8–7.4 (m, 16 H, aromatic), 3.10 (s, 24 H, $-\text{OCH}_2-$), 1.03 (s, 9 H, $-\text{CH}_3$). ^{13}C NMR (CDCl_3): 31.32 (s, 3 C, $-\text{CH}_3$), 69.19 (s, 12 C, $-\text{OCH}_2-$), 70.50 (s, 1 C, *tert*-C of OBU), 119.18 (s, C4), 123.93 (s, C2), 126.91 (s, C3), 128.08 (s, C5), 131.33 (s, C6), 157.11 (s, C1). ^{29}Si NMR (CH_2Cl_2): -125.84 . Anal. Calcd for $\text{C}_{44}\text{H}_{51}\text{O}_{11}\text{Cl}_2\text{KSi}$: C, 57.29; H, 5.94. Found: C, 58.40; H, 6.2. The formula used for the analysis includes a molecule of CH_2Cl_2 . The latter was not found in the X-ray study of **2**.

Potassium 18-Crown-6 Bis(biphenyl-2,2'-diolato)fluorosilicate, $[(\text{C}_{12}\text{H}_8\text{O}_2)_2\text{SiF}][\text{K}, 18\text{-c-6}]$ (**3**). To a suspension of spirocyclic bis(biphenyl) orthosilicate (2.20 g, 5.5 mmol) in ~20 mL of THF was added potassium fluoride (0.326 g, 5.5 mmol) and 18-crown-6 (1.45 g, 5.5 mmol). The mixture was stirred overnight. The reaction mixture was filtered and the filtrate layered with ~30 mL of hexane and kept at 0 °C for 2 days. White microcrystalline product separated out, which was filtered off, washed with hexane twice, and dried: mp ~135 °C dec; yield 3.1 g, 79%. ^1H NMR (CDCl_3): 6.8–7.45 (m, 16 H, aromatic), 3.24 (s, 24 H, $-\text{OCH}_2-$). ^{13}C NMR (CDCl_3): 69.59 (s, 12 C, $-\text{OCH}_2-$), 120.21 (s, C4), 123.15 (s, C2), 127.39 (s, C3), 128.61 (s, C5), 130.92 (s, C6), 155.92 (s, C1). ^{19}F NMR (CDCl_3): -132.29 . ^{29}Si NMR (CH_2Cl_2): -125.53 (d, $J_{29\text{Si}-19\text{F}} = 183$ Hz). Anal. Calcd for $\text{C}_{36}\text{H}_{40}\text{O}_{10}\text{FSiK}$: C, 60.08; H, 5.56. Found: C, 61.26; H, 6.11.

Reactions. (a) Reaction of Bis(2,2'-biphenoxy)silane with KO-*i*-Pr and 18-Crown-6. To the mixture of bis(2,2'-biphenoxy)silane (1.07 g, 2.7 mmol) and 18-crown-6 (0.71 g, 2.7 mmol) in dry THF was added dropwise potassium isopropoxide (0.26 g, 2.7 mmol) in ~10 mL of THF. Spontaneously white solid compound separated out (**4**). Stirring continued overnight. The solid product was filtered out, washed with dry ether twice (~20 mL) and dried: mp ~190 °C (yield 0.98 g, 54%). ^1H NMR (CDCl_3): 3.29 (s, 24 H, CH_2-O), 6.80–7.41 (m, 16 H, aromatic protons). ^{13}C NMR (CDCl_3): 69.70 ($-\text{CH}_2-\text{O}$), 118.14 (C1/C1'), 118.80 (C3/C3'), 128.27 (C5/C5'), 131.16 (C4/C4', C6/C6'), 156.76 (C2/C2'). No ^{29}Si NMR signal was found. Anal. Calcd for $\text{C}_{36}\text{H}_{43}\text{O}_{10}\text{K}$: C, 64.06; H, 6.31. Found: C, 63.8; H, 6.66.

(b) Reaction of Bis(2,2'-biphenoxy)silane with Tetrabutylammonium Fluoride. To a suspension of bis(2,2'-

biphenoxy)silane (1.7 g, 4.27 mmol) in dry THF (~30 mL) was added dropwise Bu₄NF solution (1.2 mL of 1.0 M solution in THF) at room temperature. Immediately solid silane had gone into solution. After 1 h of stirring at room temperature, the mixture was filtered. The filtrate part was layered with hexane and left at room temperature. Microcrystalline compound **5** that separated was filtered out and washed with hexane (2 × 5 mL): mp 189 °C; yield 2.6 g, 41%. ¹H NMR (CDCl₃): 0.672 (s, 8 H, CH₂), 0.965 (s, 16 H, CH₃), 2.47 (s, 12 H, CH₃), 6.60–7.45 (m, 16 H, aromatic protons). ¹³C NMR (CDCl₃): 13.44 (—CH₃), 19.24 (CH₂—CH₃), 23.36 (CH₂—CH₂—CH₃), 58.08 (N—CH₂), 118.27 (C1/C1'), 118.48 (C3/C3'), 128.35 (C5/C5'), 131.03 (C4/C4', C6/C6'), 157.47 (C2/C2'). No ²⁹Si NMR signal was found. Anal. Calcd for C₃₂H₃₅O₄N: C, 78.30; H, 8.97; N, 2.28. Found: C, 77.84; H, 9.03; N, 2.42.

(c) *Reaction of 2,2'-Biphenoxydiisopropoxysilane with KO-i-Pr and 18-Crown-6.* To freshly prepared KO-i-Pr (0.80 g, 8.1 mmol) in toluene was added 18-crown-6 (2.1 g, 8.0 mmol). The reaction was stirred for ~5 min, and then the silane compound (2.78 g, 8.4 mmol) dissolved in toluene was added dropwise. After ~10 min, white solid started separating out. Stirring was continued overnight. The white powder that separated out was filtered off and washed with ether (2 × 10 mL). Finally the product (**6**) was recrystallized from cold CH₂Cl₂/hexane (1:5): mp 180–182 °C; yield 2.6 g, 40%. ¹H NMR (CDCl₃): 3.21 (s, 72 H, —CH₂—O), 6.58–7.41 (m, 16 H, aromatic protons). ¹³C NMR (CDCl₃): 69.8 (—CH₂—O), 115.15 (C1/C1'), 119.55 (C3/C3'), 127.10 (C5/C5'), 130.11 and 130.31 (C4/C4', C6/C6'), 162.59 (C2/C2'). No ²⁹Si NMR signal was found. Anal. Calcd for C₆₀H₈₀O₂₂K₃: C, 57.10; H, 7.6. Found: C, 58.58; H, 6.90.

X-ray Measurements. The X-ray crystallographic studies were done by using an Enraf-Nonius CAD4 diffractometer and graphite-monochromated molybdenum radiation ($\lambda(K\alpha) = 0.71073 \text{ \AA}$) at an ambient temperature of $23 \pm 2 \text{ }^\circ\text{C}$. Details of the experimental procedures have been described previously.²⁴ Crystals were mounted in thin-walled glass capillaries, which were sealed as a precaution against moisture sensitivity. Data were collected by use of the θ - 2θ scan mode, with $3^\circ \leq 2\theta(\text{Mo } K\alpha) \leq 43^\circ$. No corrections were made for absorption. The structures were solved by use of direct methods and difference Fourier techniques and were refined by full-matrix least squares.²⁵ For [(C₁₀H₆O₂)₂Si(C₆H₅)] [K, 18-c-6] (**1**), of the 50 independent non-hydrogen atoms, 46 were refined anisotropically. Atoms O9 C33, C34, and C35 of the 18-c-6 species were poorly defined and were refined isotropically. The 41 independent hydrogen atoms were included in the refinement in ideal positions as fixed isotropic scatterers.

For [(C₁₂H₈O₂)₂H₃][(n-Bu)₄N] (**5**), the 45 independent non-hydrogen atoms were refined anisotropically. The 52 independent C—H hydrogen atoms were included in the refinement as fixed isotropic scatterers. The three hydroxyl hydrogen atoms of the anion appeared as the most prominent features on a difference Fourier synthesis and were refined isotropically.

Compound **6**, [(C₁₂H₈O₂)₂H][K, 18-c-6]₃, crystallizes in the monoclinic space group *P*2₁/*c* with *a* = 13.858 (5) Å, *b* = 18.790 (6) Å, *c* = 14.193 (5) Å, $\beta = 109.81 (3)^\circ$, and *Z* = 2. The anion and one

TABLE I
Crystallographic Data for [(C₁₀H₆O₂)₂Si(C₆H₅)] [K, 18-c-6] (**1**) and [(C₁₂H₈O₂)₂H₃][(n-Bu)₄N] (**5**)

compd	1	5
formula	C ₃₈ H ₄₁ O ₁₀ SiK	C ₄₀ H ₃₅ O ₄ N
fw	724.92	613.89
cryst color	colorless	colorless
dimens, mm	0.20 × 0.35 × 0.45	0.24 × 0.35 × 0.70
space group	<i>P</i> 2 ₁ / <i>n</i> (No. 14)	<i>P</i> 1̄ (No. 2)
<i>a</i> , Å	9.798 (1)	9.315 (6)
<i>b</i> , Å	20.919 (4)	9.467 (3)
<i>c</i> , Å	18.300 (4)	21.293 (6)
α , deg		78.89 (2)
β , deg	101.12 (1)	84.89 (4)
γ , deg		80.84 (4)
<i>V</i> , Å ³	3680.4	1815.7
<i>Z</i>	4	2
<i>D</i> _{calc} , g/cm ³	1.309	1.123
μ , cm ⁻¹	2.274	0.662
no. of independent reflcs measd	4193 (+ <i>h</i> , + <i>k</i> , ± <i>l</i>)	4142 (+ <i>h</i> , ± <i>k</i> , ± <i>l</i>)
no. of obsd reflcs <i>I</i> ≥ 3 σ _{<i>I</i>}	2212	2387
<i>R</i> (<i>F</i>)	0.058	0.055
<i>R</i> _w (<i>F</i>)	0.070	0.067

of the two independent cations have crystallographic C_i symmetry. Severe disorder prevented a complete solution of the structure.

All computations were performed on a Microvax II computer using the Enraf-Nonius SDP system of programs. Crystallographic data for **1** and **5** are summarized in Table I.²⁶

RESULTS

The atom-labeling scheme for the anion in **1** is shown in the ORTEP plot of Figure 1. Selected atomic coordinates are listed in Table II and selected bond lengths and angles are given in Table III for **1**. Similar information is provided in Tables IV and V for **5**, while Figure 2 shows an ORTEP plot of the anion in **5**. Complete atomic coordinates, anisotropic thermal parameters, additional bond lengths and angles, and hydrogen atom parameters for **1** and **5** are provided as supplementary material.

DISCUSSION

Structural Considerations and Comparisons with Phosphorus Analogues

The X-ray structural analysis reveals that the spirocyclic silicate **1** exists in a trigonal-bipyramidal framework with the six-membered rings positioned at axial–equatorial sites and the phenyl group in an equatorial position. Distortions away from the ideal TBP geometry follow the Berry pseudorotation coordinate,²⁷ which connects this TBP with a rectangular pyramid (RP) having four basal oxygen atoms and C21 of the phenyl group in the apical position. By use of the dihedral angle method²⁸




FIGURE 1 ORTEP plot of $[(C_{10}H_6O_2)_2Si(C_6H_5)]^- [K^+, 18-c-6]$ (**1**) with thermal ellipsoids at the 30% probability level. Atoms of the 18-c-6 species and hydrogen atoms are omitted for clarity.

TABLE II
Selected Atomic Coordinates and B_{equiv} Values in Crystalline
[(C₁₀H₆O₂)₂Si(C₆H₅)] [K, 18-c-6] (1)^a

atom ^a	x	y	z	B_{equiv} , Å ²
K	0.5394 (2)	-0.16846 (8)	1.23624 (9)	4.91 (4)
Si	0.5914 (2)	-0.3032 (1)	1.1052 (1)	4.02 (4)
O1	0.4752 (4)	-0.3672 (2)	1.0816 (2)	4.5 (1)
O2	0.5612 (4)	-0.2988 (2)	1.1934 (2)	4.5 (1)
O3	0.6771 (4)	-0.2295 (2)	1.1314 (2)	4.5 (1)
O4	0.5001 (4)	-0.2722 (2)	1.0252 (2)	4.5 (1)
C1	0.4395 (6)	-0.4130 (3)	1.1258 (4)	4.2 (2)
C2	0.3764 (7)	-0.4680 (3)	1.0955 (4)	5.2 (2)
C3	0.3357 (8)	-0.5158 (4)	1.1411 (4)	6.2 (2)
C4	0.3615 (9)	-0.5091 (4)	1.2164 (4)	6.8 (2)
C5	0.461 (1)	-0.4445 (4)	1.3274 (4)	7.8 (3)
C6	0.527 (1)	-0.3907 (4)	1.3581 (4)	8.1 (3)
C7	0.5635 (8)	-0.3409 (4)	1.3118 (4)	6.2 (2)
C8	0.5308 (7)	-0.3483 (3)	1.2361 (4)	4.2 (2)
C9	0.4673 (7)	-0.4046 (3)	1.2045 (4)	4.2 (2)
C10	0.4920 (7)	-0.4537 (4)	1.2503 (4)	5.5 (2)
C11	0.7223 (6)	-0.1890 (3)	1.0849 (4)	4.6 (2)
C12	0.8311 (7)	-0.1483 (4)	1.1108 (4)	5.9 (2)
C13	0.8705 (8)	-0.1022 (4)	1.0628 (5)	7.5 (2)
C14	0.8023 (8)	-0.0953 (4)	0.9921 (5)	7.2 (2)
C15	0.6067 (9)	-0.1271 (4)	0.8914 (4)	7.3 (2)
C16	0.4925 (9)	-0.1644 (4)	0.8680 (4)	7.3 (2)
C17	0.4542 (8)	-0.2128 (4)	0.9139 (4)	6.0 (2)
C18	0.5330 (7)	-0.2227 (3)	0.9831 (4)	4.4 (2)
C19	0.6482 (7)	-0.1835 (3)	1.0102 (4)	4.7 (2)
C20	0.6870 (7)	-0.1352 (4)	0.9630 (4)	5.9 (2)
C21	0.7519 (6)	-0.3498 (3)	1.0990 (4)	3.9 (2)
C22	0.8770 (7)	-0.3369 (4)	1.1472 (4)	5.5 (2)
C23	0.9979 (8)	-0.3728 (4)	1.1476 (4)	6.6 (2)
C24	0.9971 (7)	-0.4205 (4)	1.0969 (4)	6.2 (2)
C25	0.8781 (8)	-0.4332 (4)	1.0463 (4)	6.6 (2)
C26	0.7563 (7)	-0.3986 (4)	1.0485 (4)	5.7 (2)

^aNumbers in parentheses are estimated standard deviations.

^bAtoms are labeled to agree with Figure 1. ^cEquivalent isotropic thermal parameters are calculated as $(4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$.

TABLE III
Selected Distances (Å) and Angles (deg) for
[(C₁₀H₆O₂)₂Si(C₆H₅)] [K, 18-c-6] (1)^a

Distances			
Si-O1	1.757 (5)	K-O5	2.788 (6)
Si-O2	1.699 (5)	K-O6	2.809 (6)
Si-O3	1.778 (5)	K-O7	2.917 (7)
Si-O4	1.690 (4)	K-O8	2.825 (6)
Si-C21	1.872 (7)	K-O9	2.893 (9)
K-O2	2.856 (5)	K-O10	2.882 (7)
K-O3	2.851 (5)		
Angles			
O1-Si-O2	93.0 (2)	O2-Si-O4	129.9 (2)
O1-Si-O3	167.9 (2)	O2-Si-C21	112.9 (3)
O1-Si-O4	82.5 (2)	O3-Si-O4	92.5 (2)
O1-Si-C21	95.8 (3)	O3-Si-C21	96.3 (2)
O2-Si-O3	81.7 (2)	O4-Si-C21	117.2 (3)

^aEstimated standard deviations in parentheses. The atom-labeling scheme is shown in Figure 1.

TABLE IV
Selected Atomic Coordinates and B_{equiv} Values in Crystalline
[(C₁₂H₈O₂)₂H₃][*n*-Bu₄N] (5)^a

atom ^b	<i>x</i>	<i>y</i>	<i>z</i>	B_{equiv} ^c Å ²
O1	0.9231 (3)	-0.1364 (3)	0.2002 (1)	4.74 (8)
O11	1.0838 (3)	-0.3838 (3)	0.2346 (1)	5.47 (9)
O21	0.7161 (3)	0.0106 (3)	0.2501 (1)	4.72 (8)
O31	0.6122 (3)	0.2640 (3)	0.2625 (1)	5.06 (8)
C1	0.9156 (5)	-0.1457 (5)	0.1376 (2)	4.0 (1)
C2	1.0392 (5)	-0.2053 (4)	0.1050 (2)	3.8 (1)
C3	1.0275 (5)	-0.2110 (5)	0.0407 (2)	5.3 (1)
C4	0.8979 (6)	-0.1581 (6)	0.0098 (2)	6.0 (1)
C5	0.7765 (5)	-0.1012 (5)	0.0436 (2)	5.7 (1)
C6	0.7857 (5)	-0.0947 (5)	0.1073 (2)	4.8 (1)
C11	1.2006 (5)	-0.3476 (5)	0.1936 (2)	4.5 (1)
C12	1.1833 (5)	-0.2621 (4)	0.1329 (2)	4.0 (1)
C13	1.3125 (5)	-0.2355 (5)	0.0951 (2)	5.3 (1)
C14	1.4500 (5)	-0.2953 (6)	0.1166 (3)	6.3 (1)
C15	1.4610 (6)	-0.3822 (6)	0.1756 (3)	6.5 (1)
C16	1.3377 (5)	-0.4073 (5)	0.2149 (2)	5.9 (1)
C21	0.6422 (4)	-0.0594 (5)	0.3011 (2)	4.1 (1)
C22	0.6160 (5)	-0.0039 (5)	0.3583 (2)	4.3 (1)
C23	0.5377 (5)	-0.0834 (5)	0.4097 (2)	5.4 (1)
C24	0.4905 (5)	-0.2129 (6)	0.4050 (3)	6.1 (1)
C25	0.5184 (5)	-0.2636 (5)	0.3484 (3)	6.0 (1)
C26	0.5921 (5)	-0.1881 (5)	0.2971 (2)	5.2 (1)
C31	0.6648 (5)	0.2575 (5)	0.3214 (2)	4.5 (1)
C32	0.6701 (4)	0.1311 (5)	0.3675 (2)	4.2 (1)
C33	0.7235 (5)	0.1338 (6)	0.4263 (2)	6.1 (1)
C34	0.7677 (6)	0.2600 (6)	0.4393 (3)	7.5 (2)
C35	0.7563 (6)	0.3829 (6)	0.3921 (3)	7.7 (2)
C36	0.7062 (5)	0.3839 (5)	0.3329 (3)	6.2 (1)
H1	0.810 (5)	-0.090 (5)	0.223 (2)	5 (1)
H11	1.008 (5)	-0.296 (5)	0.224 (2)	5 (1)
H21	0.656 (5)	0.134 (5)	0.249 (2)	5 (1)

^a Numbers in parentheses are estimated standard deviations.

^b Atoms are labeled to agree with Figure 2. ^c Equivalent isotropic thermal parameters are calculated as $(4/3)[a^2\beta_{11} + b^2\beta_{22} + c^2\beta_{33} + ab(\cos \gamma)\beta_{12} + ac(\cos \beta)\beta_{13} + bc(\cos \alpha)\beta_{23}]$.

TABLE V
Selected Distances (Å) and Angles (deg) for
[(C₁₂H₈O₂)₂H₃][*n*-Bu₄N] (5)^a

Distances			
H1-O1	1.18 (4)	O1-C1	1.362 (5)
H1-O21	1.36 (4)	O11-C11	1.377 (5)
H11-O1	1.60 (4)	O21-C21	1.351 (5)
H11-O11	1.00 (4)	O31-C31	1.372 (6)
H21-O21	1.21 (4)	O1-O11	2.584 (4)
H21-O31	1.31 (5)	O1-O21	2.477 (4)
		O21-O31	2.495 (4)
Angles			
O1-H1-O21	154 (4)	C11-O11-H11	103 (2)
O1-H11-O11	165 (4)	C21-O21-H1	109 (2)
O21-H21-O31	164 (4)	C21-O21-H21	100 (2)
C1-O1-H1	112 (2)	H1-O21-H21	151 (3)
C1-O1-H11	98 (2)	C31-O31-H21	104 (2)
H1-O1-H11	124 (2)		

^a Estimated standard deviations in parentheses. The atom-labeling scheme is shown in Figure 2.





FIGURE 2 ORTEP plot of the anion in $[(C_{12}H_8O_2)_2H_3][n-Bu_4N]$ (5) with thermal ellipsoids at the 30% probability level. Aromatic hydrogen atoms are omitted for clarity. Hydrogen bonds are shown as narrow solid lines.


to assess displacement, the geometry is displaced 29.8% (32.5% with use of unit vectors) from the TBP toward the RP.

The six-membered rings are not planar but rather have the silicon atom displaced out of the plane of the remaining five atoms in a direction toward the phenyl group. Atoms O1, O2, C1, C8, and C9 are coplanar to within ± 0.024 Å, where the Si atom is displaced from this plane by 0.478 Å. For the ring containing O3 and O4 these values are ± 0.065 and 0.532 Å.


This ring arrangement in a TBP having an axial–equatorial disposition is the one commonly observed by X-ray diffraction, independent of ring size, for both pentacoordinate anionic silicates^{2,10–12} and isoelectronic phosphoranes^{4,12j,29} with cyclic components possessing electronegative heteroatoms. In addition to examples of oxygen-containing ring systems illustrated in the Introduction, A–G,^{6,8,11} that follow this pattern, the derivatives, M–O,^{7,30,31} containing sulfur and nitrogen ring



atoms provide further examples. By way of contrast, when electronegativity requirements are relaxed, as in our study of the cyclic fluorosilicate **L**,² depicted in the Introduction, X-ray diffraction reveals that the ring is positioned diequatorially in a TBP. However, thus far no structural reports have appeared showing diequatorial placement in a TBP of any ring size in an anionic pentaoxysilicate or a pentaoxyphosphorane. This includes five-membered rings trans-annulated to six-membered rings for pentaoxyphosphoranes. Structures **P**³¹ and **Q**³² recently have been determined by X-ray analysis.³³




The structure of **1** is very close to that found for **R**,³⁴ both containing equatorial phenyl groups and possessing ring unsaturation in anionic tetraoxysilicate formulations.



The distortion along the Berry coordinate from a TBP to a RP is 29.5%^{12d} for **R** compared to 32.5% for **1** (using unit vectors²⁸). As discussed above, the six-membered ring system for **1** is close to planarity with silicon acting as the flap atom. Although one could expect that ring-strain relief provided by use of a saturated six-membered ring in an oxysilicate might provide some tendency for diequatorial ring formation, the tendency for this to occur apparently is insufficient. With more disparate electronegativity requirements, this situation is known to occur for anionic silicates with five-membered rings containing carbon atoms bonded to silicon.² Thus, **I** and **K**, containing five- and six-membered saturated carbon rings, have TBP structures² with the rings in diequatorial positions in agreement with electronegativity requirements, whereas the solid-state structure of **J** having a five-membered carbon atom unsaturated ring exists as a TBP with the ring-spanning axial-equatorial sites.² This ring arrangement places a carbon atom at an electronically unfavorable axial site.


Somewhat analogous pentacoordinated phosphorus compounds containing ring systems and acyclic ligands similar to **I**–**L** are found in the derivatives **S**–**V**.^{35–37} However, unlike the silicates, **I** and **K**, which are trigonal bipyramidal, the corresponding phosphoranes, **S**³⁵ and **T**,³⁶ are square pyramidal. **S** is displaced 93.3% from the trigonal bipyramid toward the square pyramid, while **T** is displaced 81% on the basis of actual bond distances.²⁸ The presence of the fluorine ligand in the




silicates no doubt contributes to stabilizing the TBP over the SP geometry. A general observation for the formation of square pyramids of main-group five-coordinated elements is that a spirocyclic structure contain an acyclic ligand of low electronegativity.³⁸ The presence of two five-membered rings with unsaturation and having like atoms in each ring bonded to the central atom are additional conditions favoring the formation of square pyramids.^{4,38} These conditions are met for the phosphoranes S and T but not for the silicates I and K. A variable-temperature ^{19}F NMR study³⁷ of the fluorophosphoranes U and V containing the same saturated ring systems as the silicates I and K, respectively, indicates the expected TBP geometry.

Reactions

Synthesis of the first oxygen-containing six-membered ring system in an anionic silicate **1** was accomplished by reaction of phenyltrimethoxysilane with a diol in the presence of potassium methoxide and 18-crown-6 in toluene (eq 1). In contrast,




synthesis of the first five-coordinated anionic silicates with oxygen-containing seven-membered rings **2** and **3** was accomplished by reaction of the spirocyclic bis(biphenyl) orthosilicate precursor with KR in the presence of 18-crown-6 in THF (eq 2).



Unlike **1**, which was subjected to a successful X-ray structural analysis, both **2** and **3**, available as crystalline products, rapidly degraded during data collection, thus preventing a structural determination. Although fairly stable, these spirocyclic derivatives **2** and **3** slowly disproportionate into tetracoordinated silicon compounds and organic entities lacking silicon. Evidence for this process is obtained from ^{29}Si NMR spectroscopy on **2** and **3**. Initially, one signal at ~ -125 ppm, attributable to pentacoordinated silicon, was found along with a less intense signal at ~ -52 ppm in the tetracoordinate region. As time proceeds, the intensity of the high-field signal decreases whereas the intensity of the signal at ~ -52 ppm increases. Attempts to isolate the product with the signal at -52 ppm failed.

In further reactions analogous to that expressed in eq 2 using ligands such as *O*-*i*-Pr and $\text{Bu}_4\text{N}^+\text{F}^-$, less stability was encountered and only the formation of anionic organic systems was established. We postulate that these reactions (eqs 3–5) ini-



tially yield five-coordinated cyclic anionic silicates (by analogy with the formation of **2** and **3**) that then underwent rapid hydrolytic cleavage.


The use of $n\text{-Bu}_4\text{N}^+\text{F}^-$ in place of $\text{K}(\text{O-}i\text{-Pr})$ and 18-crown-6 resulted in the same type of reaction yielding **5**. In both cases, the white solid isolated showed similar elemental analyses and ^1H and ^{13}C NMR spectra indicating the same formulation, **4** = **5**. In the case of **5**, an X-ray study confirmed the formulation and showed that the structure of the anion consisted of two biphenolate units, hydrogen bonded (Figure 2). Three hydrogen atoms are required to maintain electrical neutrality. This composition agrees with the solution-state ^1H NMR spectrum giving a ratio of $(n\text{-Bu})_4\text{N}^+$ protons to aromatic protons of 2.2 (theory, 2.25). For **4**, the ^1H NMR spectrum gave a ratio of the CH_2 protons for K,18-c-6 to the aromatic protons of 1.3 (theory, 1.5).

The detailed structure of the anion in **5** (Figure 2) shows that one of the biphenolate species is monoprotonated and that one is diprotonated. The latter acts as a donor in a hydrogen bond to the former. Within each biphenolate species there is an intramolecular hydrogen bond, which results in the formation of a seven-membered ring. Coplanarity of the two phenyl rings in a given biphenolate species would not be expected due to steric considerations. For the two separate species in **5**, the dihedral angles between bonded phenyl rings are 40.4° (**2**) and 40.0° (**2**). The essential equality of these values most likely reflects constraints imposed by the intramolecular hydrogen bonds. The dihedral angles between phenyl groups of different biphenolate species range from 73.5° (**2**) to 83.0° (**1**).

Elemental analysis of the white solid isolated from the reaction of **X** with $\text{K}(\text{O-}i\text{-Pr})$ agreed with the formulation **6**. A partial X-ray study is consistent with this composition. The 3:1 ratio of K,18-c-6 cations to one hydrogen-bonded bis-(biphenolate) anion for **6** was also indicated by ^1H NMR spectroscopy of a CDCl_3 solution. The integrated intensity of the signal assigned to the CH_2 protons of the 18-c-6 units relative to that for the aromatic protons was 4.5 (theory, 4.5). No silicon was detected by ^{29}Si NMR spectroscopy in CDCl_3 solutions of **4**–**6**. Thus, the compositions of anionic organic species present in the solid state appear to be retained in solution. The 2-propanol and silicic acid compositions were not identified as products but are included in eq 3 and eq 5 for material balance.

It is likely that the slower decomposition of the more stable five-coordinated silicates **2** and **3** also proceeds to yield organic anions similar to **4**–**6**. Either the use of Bu_4N^+ as a cation in place of K,18-c-6 or use of a monocyclic silane reactant in place of spirocyclic derivative lowers the stability of pentacoordinate anionic silicate formation containing the seven-membered biphenolato ring system. In general, use of K,18-c-6 improves hydrolytic stability for pentacoordinated anionic silicates whether they contain cyclic^{2,11,12k} or acyclic^{11a,14,39–42} ligands, as found since its first use by Damrauer.^{39a}

The hydrolysis presumably is initiated by attack at silicon by a water molecule resulting in the formation of a hexacoordinated activated state, which undergoes Si–O bond cleavage and yields a postulated intermediate of the type **Z**. Repetition of the process is envisioned to lead to the organic products. The instability of five-coordinate anionic silicates reported here compared to their four-coordinate silane precursors provides another example of enhanced reactivity^{13–20} for this class of substances. In related work, Kira *et al.*¹⁷ found pentacoordinate anionic allyl de-



rivatives containing the same ring system as is present in **2** and **3** to show a greater reactivity with benzaldehyde than that of the tetracoordinate silicon analogues. Theoretical models have emphasized the lower silicon-ligand bond strengths⁴² and greater positive charge at silicon⁴³ in the pentacoordinated anionic state compared to that in the tetracoordinate state as factors influencing enhanced reactivity of five-coordinated anionic silicon.

ACKNOWLEDGEMENT

The support of this research by the National Science Foundation (Grant CHE 88-19152) is gratefully acknowledged.

Supplementary Material Available: Tables of atomic coordinates, thermal parameters, additional bond lengths and angles, and hydrogen atom parameters (Tables S1–S4 for **1** and Tables S5–S8 for **5**, respectively) (22 pages); tables of calculated and observed structure factors (20 pages). Ordering information is given on any current masthead page.

REFERENCES AND NOTES

- (1) Pentacoordinated Molecules. 87.
- (2) Part 86: Day, R. O.; Sreelatha, C.; Deiters, J. A.; Johnson, S. R.; Holmes, J. M.; Howe, L.; Holmes, R. R. *Organometallics* **1991**, *10*, 1758.
- (3) Present address: School of Chemistry, University of Hyderabad, Hyderabad 500 134, India.
- (4) Holmes, R. R. *Pentacoordinated Phosphorus-Structure and Spectroscopy*; ACS Monograph 175; American Chemical Society: Washington, DC, 1980; Vol. I, Chapter 3 and references cited therein.
- (5) Holmes, R. R. *Pentacoordinated Phosphorus-Reaction Mechanisms*; ACS Monograph 176; American Chemical Society: Washington, DC, 1980; Vol. II, Chapter 2 and references cited therein.
- (6) Burton, S. D.; Kumara Swamy, K. C.; Holmes, J. M.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 6104.
- (7) Kumara Swamy, K. C.; Holmes, J. M.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 6092.
- (8) Kumara Swamy, K. C.; Day, R. O.; Holmes, J. M.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 6095.
- (9) (a) Kumara Swamy, K. C.; Burton, S. D.; Holmes, J. M.; Day, R. O.; Holmes, R. R. *Phosphorus, Sulfur, Silicon* **1990**, *53*, 437. *Ibid.* *49/50*, 367.
- (10) Holmes, R. R. *Chem. Rev.* **1990**, *90*, 17.
- (11) (a) Kumara Swamy, K. C.; Chandrasekhar, V.; Harland, J. J.; Holmes, J. M.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 2341. (b) Holmes, R. R.; Day, R. O.; Payne, J. S. *Phosphorus, Sulfur, Silicon* **1989**, *42*, 1.
- (12) Structural details from X-ray analysis of a few anionic silicates with five-membered rings have been reported. In addition to those cited in ref. 10 and 11, they are: (a) Dixon, D. A.; Hertler, W. R.; Chase, D. B.; Farnham, W. B.; Davidson, F. *Inorg. Chem.* **1988**, *27*, 4012. (b) Kira, M.; Sato, K.; Kabuto, C.; Sakurai, H. *J. Am. Chem. Soc.* **1989**, *111*, 3747. (c) Harland, J. J.; Day, R. O.; Vollano, J. F.; Sau, A. C.; Holmes, R. R. *J. Am. Chem. Soc.* **1981**, *103*, 5269. (d) Holmes, R. R.; Day, R. O.; Harland, J. J.; Sau, A. C.; Holmes, J. M. *Organometallics* **1984**, *3*, 341. (e) Holmes, R. R.; Day, R. O.; Harland, J. J.; Holmes, J. M. *Organometallics* **1984**, *3*, 347. (f) Holmes, R. R.; Day, R. O.; Chandrasekhar, V.; Holmes, J. M. *Inorg. Chem.* **1985**, *24*, 2009. (g) Holmes, R. R.; Day, R. O.; Chandrasekhar, V.; Harland, J. J.; Holmes, J. M. *Inorg. Chem.* **1985**, *24*, 2016. (h) Farnham, W. B.; Harlow, R. L. *J. Am. Chem. Soc.* **1981**, *103*, 4608. (i)

- Stevenson, W. H., III; Wilson, S.; Martin, J. C.; Farnham, W. B. *J. Am. Chem. Soc.* **1985**, *107*, 6340. (j) Holmes, R. R. *Prog. Inorg. Chem.* **1984**, *32*, Chapter 2. (k) Tamao, K.; Hayashi, T.; Ito, Y. *J. Am. Chem. Soc.* **1990**, *112*, 2422.
- (13) Corriu, R. J. P.; Guerin, C.; Henner, B. J. L.; Wong Chi Man, W. W. C. *Organometallics* **1988**, *7*, 237.
- (14) Johnson, S. E.; Deiters, J. A.; Day, R. O.; Holmes, R. R. *J. Am. Chem. Soc.* **1989**, *111*, 3250.
- (15) Cerveau, G.; Chuit, C.; Corriu, R. J. P.; Reye, C. *J. Organomet. Chem.* **1987**, *328*, C17.
- (16) Hosomi, A.; Kohra, S.; Tominaga, Y. *J. Chem. Soc., Chem. Commun.* **1987**, 1517.
- (17) Kira, M.; Sato, K.; Sakurai, H. *J. Am. Chem. Soc.* **1988**, *110*, 4599.
- (18) Kira, M.; Kobayashi, M.; Sakurai, H. *Tetrahedron Lett.* **1987**, *28*, 4081.
- (19) Boyer, J.; Breliere, C.; Corriu, R. J. P.; Kpoton, A.; Poirier, M.; Royo, G. *J. Organomet. Chem.* **1986**, *311*, C39.
- (20) Corrius, R.; Guerin, C.; Henner, B.; Wang, Q. *J. Organomet. Chem.* **1989**, *365*, C7.
- (21) Holmes, R. R.; Day, R. O.; Johnson, S. E.; Kumara Swamy, K. C.; Holmes, J. M.; Deiters, J. A. Presentation at the IXth International Symposium on Organosilicon Chemistry, Edinburgh, Scotland, UK, July, 1990; Abstract B-3.
- (22) Schwarz, R.; Kuchea, W. Z. *Anorg. Allg. Chem.* **1955**, *279*, 84.
- (23) Silox, C. M.; Zuckerman, J. J. *J. Organomet. Chem.* **1966**, *5*, 483.
- (24) Sau, A. C.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1981**, *20*, 3076.
- (25) The function minimized was $\Sigma w(|F_o| - |F_c|)^2$, where $w^{1/2} = 2F_o L_p \sigma_l$.
- (26) $R = \Sigma ||F_o| - |F_c|| / \Sigma |F_o|$ and $R_w = |\Sigma w(|F_o| - |F_c|)|^2 \Sigma w |F_o|^2 |^{1/2}$.
- (27) Berry, R. S. *J. Chem. Phys.* **1960**, *32*, 933.
- (28) Holmes, R. R.; Deiters, J. A. *J. Am. Chem. Soc.* **1977**, *99*, 3318.
- (29) Holmes, R. R. *Acc. Chem. Res.* **1979**, *12*, 257 and references cited therein.
- (30) Day, R. O.; Kumara Swamy, K. C.; Fairchild, L.; Holmes, J. M.; Holmes, R. R. *J. Am. Chem. Soc.* **1991**, *113*, 1627.
- (31) Holmes, R. R.; Kumara Swamy, K. C.; Holmes, J. M.; Day, R. O. *Inorg. Chem.* **1991**, *30*, 1052. Deiters, J. A.; Sreelatha, C.; Day, R. O.; Holmes, R. R. Preliminary account presented at the 200th National Meeting of the American Chemical Society, Washington, DC, August, 1990; INOR 26.
- (32) Yu, J. H.; Sopchik, A. E.; Arif, A. M.; Bentrude, W. G. *J. Org. Chem.* **1990**, *55*, 3444.
- (33) Both P^{31} and Q^{32} have been used as models for intermediates in c-AMP action.
- (34) Boer, F. P.; Flynn, J. J.; Turley, J. W. *J. Am. Chem. Soc.* **1968**, *90*, 6973.
- (35) Devillers, J. R.; Holmes, R. R. *J. Am. Chem. Soc.* **1977**, *99*, 3322.
- (36) Clark, T. E.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1979**, *18*, 1660.
- (37) Muetterties, E. L.; Mahler, W.; Schmutzler, R. *Inorg. Chem.* **1963**, *2*, 613.
- (38) Holmes, R. R. *J. Am. Chem. Soc.* **1975**, *97*, 5379.
- (39) (a) Damrauer, R.; Danahey, S. E. *Organometallics* **1986**, *5*, 1490. (b) Damrauer, R.; O'Connell, B.; Danahey, S. E.; Simon, R. *Organometallics* **1989**, *8*, 1167.
- (40) Johnson, S. E.; Day, R. O.; Holmes, R. R. *Inorg. Chem.* **1989**, *28*, 3182.
- (41) Johnson, S. E.; Payne, J. S.; Day, R. O.; Holmes, J. M.; Holmes, R. R. *Inorg. Chem.* **1989**, *28*, 3190.
- (42) Deiters, J. A.; Holmes, R. R. *J. Am. Chem. Soc.* **1990**, *112*, 7197.
- (43) Baybutt, P. *Mol. Phys.* **1975**, *29*, 389.