

This article was downloaded by:

On: 29 January 2011

Access details: *Access Details: Free Access*

Publisher *Taylor & Francis*

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

DIMETHYL ESTERS OF 1-AMINO-2-ARYLMETHYL CYCLOHEXYL PHOSPHONIC ACIDS. SYNTHESIS AND NMR CHARACTERIZATION

Salvatore Failla^a; Paolo Finocchiaro^a; Mario Latronico^b

^a Istituto Chimico Facoltà di Ingegneria Università di Catania, Catania, Italy ^b Dipartimento di Ingegneria e Fisica dell'Università della Basilicata, Potenza, Italy

To cite this Article Failla, Salvatore , Finocchiaro, Paolo and Latronico, Mario(1995) 'DIMETHYL ESTERS OF 1-AMINO-2-ARYLMETHYL CYCLOHEXYL PHOSPHONIC ACIDS. SYNTHESIS AND NMR CHARACTERIZATION', *Phosphorus, Sulfur, and Silicon and the Related Elements*, 101: 1, 261 – 266

To link to this Article: DOI: 10.1080/10426509508042525

URL: <http://dx.doi.org/10.1080/10426509508042525>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

DIMETHYL ESTERS OF 1-AMINO-2-ARYLMETHYL CYCLOHEXYL PHOSPHONIC ACIDS. SYNTHESIS AND NMR CHARACTERIZATION

SALVATORE FAILLA and PAOLO FINOCCHIARO*

*Istituto Chimico Facoltà di Ingegneria Università di Catania,
Viale A. Doria, 6 I-95125 Catania, Italy*

and

MARIO LATRONICO

*Dipartimento di Ingegneria e Fisica dell'Università della Basilicata,
Via N. Sauro, 85100 Potenza, Italy*

(Received November 15, 1994; in final form November 30, 1994)

Dimethyl esters of amino-arylmethyl phosphonic acids containing a cyclohexyl moiety have been prepared in good yields. All compounds are white crystalline powders very soluble in low boiling point organic solvents. Characterization by ¹H-NMR techniques indicates that ortho substituents shift downfield the CH resonance, whereas ortho-carboxy groups, through hydrogen bonding with the NH group, yield a cyclic structure responsible of the dramatic downfield shift of the NH resonance.


Key words: ¹H-NMR characterization, crystalline, low-melting and very soluble amino-phosphonates, substrates for functionalization at the NH group.

INTRODUCTION


In our continuing efforts aimed at synthesizing new aryl-amino-methyl phosphonates and phosphonic acids mono-alkyl esters^{1–6} we were interested in looking at the reactivity and the chemical properties of phosphonates possessing at least one cyclohexyl moiety linked to the —CH— or —NH— fragment. The potential interest of such new series of compounds can be summarized as follow:

- i) The cycloalkyl moiety should enhance solubility in low boiling organic solvents.
- ii) Reactivity at the —NH— functionality should be increased permitting an easy access to N-substituted amino phosphonates.
- iii) Cycloalkyl groups, generally, impart specific agrochemical activities.

In this paper we wish to report on the synthesis and properties, together with their ¹H-NMR spectroscopic characterization, of different classes of variously substituted amino-arylmethyl cyclohexyl phosphonate esters of general formulas **1** and **2** given below:



In addition, we shall describe the syntheses of compounds of general formula **3** in which one of the aryl rings is bearing a carboxy or carbomethoxy group in order to impart water solubility and derivatives **4** where the benzene ring is functionalized with a trifluoromethyl group and in addition a naphthyl ring is attached to the carbon-atom bearing the phosphonate group.



RESULTS AND DISCUSSION

In our approach to the synthesis of phosphonates **1-4** the best and simplest synthetic route was found in the direct addition (generally at room temperatures) of dimethyl phosphite to the corresponding Schiff bases, which are readily available through

TABLE I
Physical properties of **1**

N.	X	Yield %	m.p.°C	¹ H-NMR		
				δ (CDCl ₃ , TMS)		
				CH ^a	NH ^b	O=P(OMe) ₂ ^c
1a	4-F	29	46-48	4.22	2.30	3.55, 3.77
1b	4-CF ₃	46	87-89	4.31	2.28	3.58, 3.78
1c	2-Cl	28	66-68	4.88	2.26	3.49, 3.86
1d	4-Cl	30	56-59	4.22	2.29	3.56, 3.77
1e	2,3-Cl ₂	45	81-82	4.93	2.25	3.52, 3.87
1f	2,4-Cl ₂	41	86-87	4.81	2.23	3.55, 3.86
1g	3,4-Cl ₂	31	72-73	4.19	2.28	3.62, 3.77
1h [†]	2-OCH ₃	43	79-80	4.85	2.28	3.48, 3.82
1i	3-OCH ₃	80	63-66	4.20	2.34	3.52, 3.77
1j	4-OCH ₃	82	53-55	4.18	2.32	3.52, 3.77
1k	2,5-(OCH ₃) ₂	91	74-76	4.81	2.30	3.52, 3.81
1l	3,4,5-(OCH ₃) ₃	35	55-58	4.17	2.38	3.57, 3.80


^a doublet ($J_{HP} = 22-23$ Hz)

^b broad singlet

^c doublets ($J_{HP} = 10-12$ Hz)

[†] Calc. for C₁₆H₂₆NO₄P: C 58.76, H 7.95, N 4.28; found C 58.68, H 8.02, N 4.31.

TABLE II
 Physical properties of 2



N.	X	Yield %	m.p.°C	¹ H-NMR		
				δ (CDCl ₃ , TMS)		
				CH ^a	NH ^a	O=P(OMe) ₂ ^b
2a	2-CF ₃	11	59-61	3.75	4.50	3.69, 3.74
2b	3-CF ₃	28	116-118	3.71	4.11	3.66, 3.75
2c	2-F	37	52-53	3.70	4.10	3.70, 3.72
2d	4-F	53	116-118	3.60	6.55	3.72, 3.76
2e	2-COOH	29	164-166	3.84	8.20	3.71, 3.83
2f	4-COOH	59	208-209	3.77	4.80	3.66, 3.77
2g	2-COOCH ₃	37	126-130	3.87	8.14	3.72, 3.85
2h	2-Cl	35	53-55	3.64	4.71	3.68, 3.74
2i	3-Cl	33	101-103	3.61	3.97	3.67, 3.74
2j [†]	4-Cl	73	144	3.60	-	3.66, 3.72
2k	2,3-Cl ₂	45	84-85	3.64	4.70	3.68, 3.73
2l	2,5-Cl ₂	23	137-138	3.56	4.58	3.70, 3.74
2m	3,4-Cl ₂	31	133-135	3.56	4.11	3.70, 3.77
2n	3,5-Cl ₂	70	172-173	3.60	4.30	3.68, 3.76
2o	2-OCH ₃	24	52-55	3.62	4.27	3.67, 3.72
2p	2-OCH ₃ , 5-Cl	31	97-98	3.62	4.56	3.67, 3.75
2q	4-N=N-Ph	20	168-170	3.70	4.46	3.67, 3.75

^a ABX system

^b doublets ($J_{HP} = 10+12$ Hz)


[†] Calc. for C₁₅H₂₃ClNO₃P: C 54.30, H 6.94, N 4.22; found C 54.24, H 6.99, N 4.19.

the condensation of primary amines with aldehydes. Most of the aldimine precursors of **1** and **2** were generally oily compounds at ambient temperatures and no attempts were made to purify them. Thus, the dimethyl phosphite was directly added to the crude reaction mixture and then, at the end of the reaction, the desired phosphonate was purified by recrystallisation from ethyl ether or from concentrated solutions of ethylacetate.

In the syntheses of phosphonates **3** and **4**, the Schiff-bases, generally high melting solids, were isolated and recrystallized before proceeding with the addition of dimethyl phosphite. In Tables I–IV are listed the physical properties of the synthesized compounds. All samples are white crystalline materials.

As far as the ¹H-NMR spectra are concerned, first of all, we remark that the aromatic or the cyclohexyl proton chemical shifts are not listed in the Tables because these protons resonate in the expected region of the magnetic field and they generally maintain the multiplicity already present in the precursor Schiff-bases.

The methyne hydrogen of the groups —CH—P(O)(OMe)₂ generally resonate as a sharp doublet with a coupling constant H—P in the range of 22 ÷ 24 Hz, except

TABLE III
 Physical properties of 3



N.	X	Y	Yield %	m.p.°C	¹ H-NMR δ (CDCl ₃ , TMS)		
					CH	NH	O=P(OMe) ₂ ^a
3a	H	2-COOH	67	153-155	4.98 ^b	8.70	3.65, 3.75
3b	H	4-COOH	71	148-150	4.86 ^c	6.13 ^c	3.44, 3.79
3c	H	2-COOCH ₃	46	84-86	4.90 ^c	8.84 ^c	3.65, 3.70
3d [†]	4-OCH ₃	4-COOH	97	110-112	4.83 ^b	6.0	3.47, 3.80
3e	4-OCH ₃	2-COOCH ₃	59	112-114	4.87 ^c	8.77 ^c	3.67, 3.75
3f	4-OCH ₃	4-CF ₃	65	-	4.75 ^b	5.10	3.47, 3.78

^a doublets ($J_{HP} = 10+12$)

^b doublet ($J_{HP} = 24$)

^c ABX system

[†] Calc. for C₁₇H₂₀NO₆P: C 55.89, H 5.48, N 3.83; found C 55.82, H 5.52, N 3.80.

 TABLE IV
 Physical properties of 4


N.	X	Yield %	m.p.°C	¹ H-NMR δ (CDCl ₃ , TMS)		
				CH	NH	O=P(OMe) ₂ ^a
4a	2-CF ₃	30	85-88	5.69 ^b	5.74 ^b	3.25, 3.78
4b	3-CF ₃	43	164-166	5.66 ^c	-	3.13, 3.82
4c	4-CF ₃	50	157-160	5.66 ^b	-	3.12, 3.82
4d [†]	4-OCH ₃	95	147-149	5.60 ^c	-	3.15, 3.82

^a doublets ($J_{HP} = 10+11$ Hz)

^b ABX system.

^c doublet ($J_{HP} = 24$ Hz)

[†] Calc. for C₂₀H₂₂NO₄P: C 64.69, H 5.93, N 3.77; found C 64.63, H 5.98, N 3.73.


in all compounds of series **2** and in some derivatives of **3** where the additional coupling with the NH proton generates an ABX pattern. In all these latter samples the CH resonances give rise at least to a four line pattern with $J_{\text{H-P}}$ in the range of $18 \div 24$ Hz and $J_{\text{H-NH}}$ $8 \div 9$ Hz, whereas the NH protons appear as a triplet with $J_{\text{NH-P}}$ nearly equal to $J_{\text{NH-CH}}$. The observation of an ABX pattern could be due to a slower exchange of the NH protons or to an increase of chemical shift difference due to the strong anisotropic moieties attached to the molecule, as already observed in cognate molecules.⁷

The methyl hydrogens of the methoxy groups always appear as two distinct doublets of equal intensity, due to the close proximity of the stereocenter N—C—P. The alternative hypothesis that the observation of two distinct doublets for the methoxy groups arise from restricted rotation, on the NMR time scale, around the CH—P bond cannot, in principle, be excluded.

The chemical shift difference between the two doublets, as can be inferred from Tables I–IV, is very sensitive to the moiety attached to the methyne carbon bearing the phosphonate group. In particular going from cyclohexyl to 1-naphthalene the $\Delta\nu$ varies from ca. 0.05 ppm as in **2a** to 0.70 ppm as in **4c**. This effect is certainly due to the strong anisotropic ring current experienced by one of the methoxy groups attached to phosphorous which is lying in the close proximity of the aryl ring, whereas the other one is far away from it.

More interesting is the comparison of the CH—P chemical shifts: in all derivatives **1–3** bearing at least one *ortho*-substituent, inspection of Tables I–III reveals that in these compounds the CH resonance is very much shifted downfield when compared with that one of the *para*- or *meta*-substituted cognates. This downfield effect is even more dramatically evident looking at the NH resonance: in phosphonates **2e**, **2g**, **3a**, **3c**, **3e** which possess a carboxy or carbomethoxy group in the *ortho* position, the NH proton resonates below 8.00 ppm, whereas its unperturbed chemical shift ranges from 4.0 to 4.80 ppm for the *para*- or *meta*-derivatives **2** and occurs at 6.13 ppm for **3b**.

No doubt that this exceptional downfield effect is due to the formation of a hydrogen bond between the NH and the *ortho* carbonyl oxygen giving rise to a cyclic structure as indicated below:



EXPERIMENTAL

Amines, aldehydes, dimethylphosphite as well as solvents and all other chemicals used were high purity commercial products from Aldrich. All syntheses were performed under a dry N_2 atmosphere.

$^1\text{H-NMR}$ spectra were recorded in CDCl_3 with Me_4Si as an internal standard using a Bruker AC-250 instrument operating at 250 MHz.

Melting points were determined on a Büchi 530 melting point apparatus and are uncorrected. The Schiff base precursors were all prepared in high yield according to the procedure previously described.³

The compounds listed in Tables I–IV were synthesized by direct reaction between the Schiff bases and the phosphite according to the following general procedure:

To a stirred solution of the Schiff-base precursor (0.1 mol) in dry toluene (50 ml) was added dropwise 14 ml (0.15 mol) of HP(O)(OMe)_2 and a catalytic amount of NaH. After the addition was completed, the mixture was stirred for a few hours. The solvent was then evaporated and the oil residue was solidified by addition of diethyl ether and a few drops of ethylacetate. All the phosphonates synthesized are white solids, except for compound **2q** (Table II) which is red. They were recrystallized from a mixture of diethyl ether and ethylacetate. Microanalytical data are consistent with the molecular formulas and for one compound each of the type **1-4** data are reported as a footnote in Tables I-IV.

ACKNOWLEDGEMENTS

We thank C.N.R. and the Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST) for financial support.

REFERENCES

1. S. Failla, P. Finocchiaro, V. K. Belsky, V. E. Zavodnik and A. N. Sobolev, *Phosphorus, Sulfur and Silicon*, **80**, 75 (1993).
2. S. Caccamese, S. Failla, P. Finocchiaro, G. Hägele and G. Principato, *J. Chem. Research (S)*, 242 (1992).
3. S. Failla, P. Finocchiaro, G. Hägele and R. Rapisardi, *Phosphorus, Sulfur and Silicon*, **82**, 79 (1993).
4. S. Failla and P. Finocchiaro, *Phosphorus, Sulfur and Silicon*, **85**, 65 (1993).
5. S. Failla, P. Finocchiaro, M. Latronico and M. Libertini, *Phosphorus, Sulfur and Silicon*, **88**, 185 (1994).
6. S. Caccamese, G. Principato, U. Gruss, G. Hägele and S. Failla, *Tetrahedron Asymmetry*, **5**, 141 (1994).
7. S. J. Smith, H. Zimmer, E. Fluck and P. Fischer, *Phosphorus and Sulfur*, **35**, 105 (1988).