

This article was downloaded by:

On: 28 January 2011

Access details: Access Details: Free Access

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title~content=t713618290>

### ADDITION of THIO COMPOUNDS to (*R*)-LIMONENE CATALYZED by AMBERLYST-15

Marcio C. S. Mattos<sup>a</sup>; Antonio M. Sanseverino<sup>a</sup>

<sup>a</sup> Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

Online publication date: 16 August 2010

**To cite this Article** Mattos, Marcio C. S. and Sanseverino, Antonio M.(2004) 'ADDITION of THIO COMPOUNDS to (*R*)-LIMONENE CATALYZED by AMBERLYST-15', Phosphorus, Sulfur, and Silicon and the Related Elements, 179: 6, 1203 — 1208

**To link to this Article:** DOI: 10.1080/10426500490459849

**URL:** <http://dx.doi.org/10.1080/10426500490459849>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## ADDITION OF THIO COMPOUNDS TO (R)-LIMONENE CATALYZED BY AMBERLYST-15

Marcio C. S. de Mattos and Antonio M. Sanseverino  
Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil

(Received May 15, 2003; accepted October 15, 2003)

*The addition of thioacids and thiophenol to (R)-limonene catalyzed by Amberlyst-15 produced the corresponding 9-thioester- or 9-thioether-p-menthene chemo- and regioselectively.*

**Keywords:** Amberlyst-15; ion-exchange resin; limonene; thioether; thioester

The catalyzed addition of thio compounds to alkenes can be with Markovnikov (using montmorillonite K-10,<sup>1</sup> Lewis,<sup>2</sup> or protic acids<sup>3</sup>) or anti-Markovnikov (using zeolites<sup>4</sup> or radical conditions<sup>5</sup>) selectivity.

Thio compounds derived from limonene are used as flavors or flotation agents.<sup>6</sup> The uncatalyzed reaction of thioacetic acid with limonene (**1**) is not chemoselective, producing a mixture of addition products to both double bonds along with diaddition products.<sup>7</sup> On the other hand, the reaction of limonene with thiols catalyzed by boron trifluoride produces the 9-thio derivatives,<sup>8</sup> whereas the reaction of thiophenol catalyzed by HClO<sub>4</sub> produces in low yield the 8-thio derivative of difficult purification<sup>9</sup> (Scheme 1).


Continuing our interest in chemo- and regiospecific functionalization of the 8,9-double bond of (*R*)-limonene,<sup>10</sup> we studied the reaction of diverse thio compounds catalyzed by Amberlyst-15, a macroreticular strong acidic sulfonic ion-exchange resin ( $H_0 = -2.2$ ),<sup>11</sup> and present our results here.

## RESULTS AND DISCUSSION

The reaction of (*R*)-limonene (10 mmol) with thioacetic acid (10 mmol) in dichloromethane at room temperature and catalytic Amberlyst-15

We thank Elizabeth R. Lachter for the kindly donation of Amberlyst-15, and W. Bruce Kover and David E. Nicodem for helpful discussions. AMS thanks CNPq for a fellowship.


Address correspondence to Marcio C. S. de Mattos, Departamento de Química, Orgânica, Instituto de Química, Universidade Federal do Rio de Janeiro, Caixa Postal 68545, 21945-970, Rio de Janeiro, Brazil. E-mail: mmattos@iq.ufrj.br



SCHEME 1

(ca. 5% in weight) led to a product **2a** along with some unreacted substrate (determined by high-resolution gas chromatography). After low-pressure distillation, **2a** was isolated in 73% yield and its MS spectra indicated incorporation of one unit of thioacetic acid to limonene ( $m/z$  212 and 214). The IR spectra of the product showed characteristic signals of thioesters ( $1693$ ,  $1135$ , and  $1109\text{ cm}^{-1}$ ) and the disappearance of the signal of the disubstituted double bond of limonene ( $890\text{ cm}^{-1}$ ), indicating a chemoselectivity to the acyclic double bond of the substrate. The  $^1\text{H}$  NMR spectra showed two multiplets at  $2.76$  and  $3.04\text{ ppm}$  (1H each) of a  $\text{CH}_2$  linked to thioester, indicating both an anti-Markovnikov addition and a diastereoisomeric mixture of products. The fact the **2a** is a pair of diastereoisomers (ca. 1:1) is confirmed by two signals of methyl at  $1.70$  and  $1.71\text{ ppm}$ . The  $^{13}\text{C}$  NMR spectrum showed a mixture of diastereoisomers too, as there were many signals with very close shifts (e.g., ring CH at  $120.6$  and  $120.8\text{ ppm}$ ,  $\text{CH}_3$  of isopropyl at  $15.9$  and  $16.3\text{ ppm}$ ). Based on these informations the proposed structure for **2a** is shown in Figure 1.

To our surprise, an anti-Markovnikov product was obtained and it was unexpected, because its formation is consistent with a radical process.<sup>5</sup> Furthermore, the addition of alcohols to limonene catalyzed by Amberlyst-15 produced regioselectively the 8-substituted terpinyl



**FIGURE 1** Selected NMR data for **2a** (mixture of diastereoisomers).

ethers.<sup>12</sup> Control experiments with freshly distilled limonene,<sup>13,\*</sup> rigorous exclusion of air, or reaction in the presence of free radicals inhibitors<sup>14</sup> led to the same product **2a** in similar yield. On the other hand, in the absence of the catalyst, the reaction of limonene with thioacetic acid was incomplete after several hours, producing the thioester **2a** in low yield along with unreacted substrate (in our hands no mixture of several products was detected as previously described<sup>7</sup>).


A possible explanation for the formation of the anti-Markovnikov product could be an initial complexation of limonene with the resin. In analogy to the case with zeolites,<sup>4</sup> due to steric constraints the bulkier Markovnikov adduct could not be retained in the resin and the attack of the thioacid occurs at the less hindered primary carbon of the complex, producing the anti-Markovnikov product.

Based on the above results, it was decided to demonstrate the generality and utility of the Amberlyst-15 catalyzed reaction of (*R*)-limonene with other thio compounds. In all cases the corresponding 9-thio-p-menthenes were obtained as a ca. 1:1 mixture of diastereoisomers (Table I summarizes the results).

The reaction of (*R*)-limonene with thiobenzoic acid produced the thioester **2b** in 65% isolated yield after 20 h [methylene adjacent to thioester at 2.73 and 3.03 ppm (m, 1H each)]. The reaction of (*R*)-limonene with thiophenol produced the thioether **2c** in 51% isolated yield [methylene adjacent to thioester at 2.75 and 3.06 ppm (m, 1H each)] along with unreacted limonene (ca. 30% recovered), diphenyldisulphide (ca. 5%) and diaddition products (ca. 10%).

On the other hand, the reactions of (*R*)-limonene with 1-propanethiol and 2-propanethiol were disappointing as the substrate was recovered

\*It is known that limonene produces peroxides quickly.

**TABLE I** Yields of **2** Obtained


Entry	R	Time (t/h)	Yield of <b>2</b> (%) <sup>a</sup>
a	CH <sub>3</sub> C(O)	18	73
b	PhC(O)	20	65
c	Ph	48	51
d	Pr	>60	— <sup>b</sup>
e	<i>i</i> -Pr	>48	— <sup>b</sup>

<sup>a</sup>Yield of pure product based on limonene.<sup>b</sup>Limonene recovered.

and no product was detected by the analytical techniques employed (high-resolution gas chromatography, <sup>1</sup>H and <sup>13</sup>C NMR spectroscopy). Several attempts to perform the reaction changing the conditions (reaction time, temperature, and addition of more catalyst) proved unsuccessful too.

The fact that thiophenol added to limonene while alkylthiols failed could be explained in terms of the proton affinity of these thio compounds. The proton affinity of alkylthiols is 3 kcal/mol higher than thiophenol (189.3 *vs* 186.3 kcal/mol<sup>15</sup>) and consequently the more basic\* alkylthiols could be protonated by the acidic resin ( $H_0 = -2.2^{11}$ ). Hence, a nucleophilic attack of thiophenol to the limonene-resin complex is possible while protonated alkylthiols do not react at all.

In summary, we have developed a very attractive route to 9-thio-menthanes. As in the case of related monoterpene sulfides,<sup>16</sup> the products offer great potentials as starting materials for multistep preparation of bisabolanes.

## EXPERIMENTAL

(*R*)-Limonene (Dierberger) was distilled at 177°C (lit.<sup>17</sup> 178°C) and stored in a freezer. Amberlyst-15 (Rohm & Haas), thio compounds

\*As the solvent (CH<sub>2</sub>Cl<sub>2</sub>) does not efficiently solvate the thiols, the basicity in the gas phase can be assumed to be the same as in solution.

(Aldrich) and other chemicals were used without further purification.  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were acquired on a Bruker AC-200 (200 MHz and 50 MHz respectively) spectrometer in  $\text{CDCl}_3$  solutions with TMS as internal standard. IR spectra were recorded on a Perkin-Elmer 1600 FT-IR spectrometer (KBr film). MS were obtained on a Hewlett-Packard HP 5896-A HRGC-MS using electron impact (70 eV). Analyses by high-resolution gas chromatography were performed on a HP 5890-II gas chromatograph with FID by using a 25 m (length), 0.53 mm (ID) and 0.50  $\mu\text{m}$  (phase thickness) RTX-5 silica capillary column (Restek Co.) and He (flow rate 50 cm/s) as carrier gas (split 1:20).

### General Procedure for the Addition of Thio Compounds to Limonene Catalyzed by Amberlyst-15

To a stirred solution of (*R*)-limonene (10 mmol) and the thio compound (10 mmol) in  $\text{CH}_2\text{Cl}_2$  (25 ml) was added Amberlyst-15 (70 mg) at room temperature. After the time shown in Table I, the catalyst was filtered off and washed with  $\text{CH}_2\text{Cl}_2$  (10 ml). The combined liquid was washed with water (10 ml), 5%  $\text{NaHCO}_3$  (10 ml), water (10 ml) and then dried ( $\text{Na}_2\text{SO}_4$ ). After evaporation of the solvent, the residue was distilled at reduced pressure.

#### (2*RS*, 3'*R*)-1-[4'-Methylcyclohex-3'-enyl]propylthio]ethan-1-one (2a) ca. 1:1 Mixture of Diastereoisomers

It was obtained 1.54 g (73%), b.p. 120–122°C/1 Torr.  $\delta_{\text{H}}$  0.92 (m, 3H), 1.25 (m), 1.70 (s,  $\text{CH}_3$ ), 1.71 (s,  $\text{CH}_3$ ), 2.35 (s), 2.76 (m, 1H), 3.04 (m, 1H), 5.37 (broad s, 1H) ppm.  $\delta_{\text{C}}$  15.9, 16.3, 23.6, 25.5, 27.3, 27.8, 29.4, 30.4, 30.6, 30.8, 34.1, 34.3, 37.8, 37.9, 38.0, 120.5, 120.6, 134.2, 196.3 ppm.  $\nu_{\text{max}}$  2961, 2924, 2836, 1693, 1451, 1354, 1135, 1109, 955, 800, 629  $\text{cm}^{-1}$ .  $m/z$  214 ( $\text{M}^+ + 2$ ), 212 ( $\text{M}^+$ ), 173, 171, 136 (100%), 123, 121, 107, 94, 79.

#### (2*RS*, 3'*R*)-2-(4'-Methylcyclohex-3'-enyl)propylthio Phenyl Ketone (2b) ca. 1:1 Mixture of Diastereoisomers

It was obtained 1.75 g (65%), b.p. 185–187 °C/1 Torr.  $\delta_{\text{H}}$  1.03 (m, 3H), 1.61–1.84 (m), 1.70 (s,  $\text{CH}_3$ ), 1.92–1.96 (m), 2.73 (m, 1H), 3.03 (m, 1H), 5.37 (broad s, 1H), 7.45–8.00 (m) ppm.  $\delta_{\text{C}}$  16.1, 23.6, 30.4, 30.6, 31.6, 34.2, 38.2, 38.3, 38.4, 41.4, 120.2, 120.3, 128.7, 129.7, 131.1, 137.3, 137.7, 139.3, 197.0 ppm.

**(2R, 3'R)-2-(4'-Methylcyclohex-3'-enyl)propylthio]benzene (2c) ca. 1:1 Mixture of Diastereoisomers**

It was obtained 1.24 g (51%), b.p. 150–153°C/1 Torr.  $\delta_{\text{H}}$  1.00 (dd), 1.63 (s, CH<sub>3</sub>), 1.77 (m), 2.75 (m, 1H), 3.06 (m, 1H), 5.36 (broad s, 1H), 7.29 (m) ppm.  $\delta_{\text{C}}$  15.9, 16.4, 23.6, 25.5, 29.7, 30.8, 30.9, 37.3, 37.4, 39.0, 39.1, 120.8, 127.3, 129.0, 129.2, 137.2, 137.7, 196.2 ppm. m/z 248 (M<sup>+</sup> + 2), 246 (M<sup>+</sup>), 136 (100%), 123, 107, 94, 79.

**REFERENCES**

- [1] S. Kanagasabapathy, A. Sudalai, and B. C. Benicewicz, *Tetrahedron Lett.*, **42**, 3791 (2001).
- [2] M. Belley and R. Zamboni, *J. Org. Chem.*, **54**, 1230 (1989).
- [3] C. G. Screttas and M. Micha-Screttas, *J. Org. Chem.*, **44**, 713 (1979).
- [4] P. Kumar, R. K. Pandey, and V. R. Hedge, *Synlett*, 1921 (1999).
- [5] K. Griesbaum, *Angew. Chem. Int. Ed. Engl.*, **9**, 273 (1970).
- [6] A. F. Thomas and Y. Bessiere, *Nat. Prod. Rep.*, **6**, 291 (1989).
- [7] G. R. Davletina, E. K. Kazakova, and S. G. Vul'fson, *Zh. Org. Khim.*, **29**, 1632 (1993).
- [8] V. A. Morgunova, L. E. Nikitina, V. V. Plemenkov, M. G. Fazylaeva, and Y. B. Chugunov, *Russ. J. Org. Chem.*, **36**, 489 (2000).
- [9] J. D. Fourneron, L. M. Harwood, and M. Julia, *Tetrahedron*, **38**, 693 (1982).
- [10] a) M. C. S. de Mattos, R. B. Coelho, and A. M. Sanseverino, *Synth. Commun.*, **34**, 541 (2004); b) M. C. S. de Mattos, S. P. L. de Souza, and S. M. Elias, *Heterocycl. Commun.*, **9**, 247 (2003); c) M. C. S. de Mattos, S. P. L. de Souza, and S. M. Elias, *Heterocycl. Commun.*, **7**, 249 (2001); d) A. M. Sanseverino, M. C. S. de Mattos, and W. B. Kover, *J. Chem. Res. (S)*, 346 (2000).
- [11] J. Tejero, F. Cunill, M. Iborra, J. F. Izquierdo, and C. Fité, *J. Mol. Catal. A: Chem.*, **182–183**, 541 (2002).
- [12] M. J. L. Rao, S. Nagarajan, and K. N. Gurudutt, *J. Sci. Ind. Res.*, **58**, 682 (1999).
- [13] W. F. Newhall, *J. Org. Chem.*, **23**, 1274 (1958).
- [14] E. T. Denisov and I. V. Khudyakov, *Chem. Rev.*, **87**, 1313 (1987).
- [15] G. A. DiLabio, D. A. Pratt, A. D. LoFaro, and J. S. Wright, *J. Phys. Chem. A*, **103**, 1653 (1999).
- [16] O. Smitt and H.-E. Hogberg, *J. Chem. Res. (S)*, 527 (2002).
- [17] R. C. Weast (Ed.), *Handbook of Chemistry and Physics* (CRC Press, Boca Raton, 1985), 65th ed.