Erratum to: Human Pancreatic Polypeptide in a Phospholipid Based Micellar Formulation

Amrita Banerjee • Hayat Onyuksel

Published online: 27 March 2012

© Springer Science+Business Media, LLC 2012

Erratum to: Pharm Res DOI:10.1007/s11095-012-0718-4

Incorrect legends for Figures 7 and 8 were published in the original article. The correct legends appear below.

Legend for Fig 7:

Particle size distribution of PP-SSM before and after lyophilization. Representative Nicomp figures showing particle size distribution of PP-SSM at lipid: peptide molar ratio of 35: 1 (a) before and (b) after freeze drying. (A) Samples prepared in phosphate buffered saline and reconstituted using sterile water; (B) samples prepared in phosphate buffer and reconstituted using normal saline. Lipid and peptide

concentration kept constant at 5 mM and 143 μM for all samples.

Legend for Fig 8:

Fluorescence emission of PP-SSM before and after lyophilization. Fluorescence emission of PP-SSM before and after freeze drying (FD) at 35: 1 lipid: peptide molar ratios; (**A**) samples prepared in phosphate buffered saline (pH 7.4) and reconstituted in sterile water; (**B**) samples prepared in phosphate buffer (pH 7.4) and reconstituted using normal saline (n=3). Lipid and peptide concentration kept constant at 5 mM and 143 μ M respectively for all samples. All results are mean \pm S.D. *p<0.05 in comparison to fluorescence emission of PP-SSM before lyophilization (n=3/group).

The online version of the original article can be found at http://dx.doi.org/ 10.1007/s11095-012-0718-4.

A. Banerjee · H. Onyuksel Department of Biopharmaceutical Sciences University of Illinois at Chicago, Chicago, Illinois 60612, USA

H. Onyuksel Department of Bioengineering, University of Illinois at Chicago Chicago, Illinois 60612, USA

H. Onyuksel (☑)
Department of Biopharmaceutical Sciences (M/C 865)
College of Pharmacy University of Illinois at Chicago, 833 South Wood
Street, Chicago, Illinois 60612-7231, USA
e-mail: hayat@uic.edu

