

ASSOCIATE EDITOR: DIANNE M. PEREZ

Purinergic Signaling in the Airways

Geoffrey Burnstock, Inge Brouns, Dirk Adriaensen, and Jean-Pierre Timmermans

Autonomic Neuroscience Centre, University College Medical School, Royal Free Campus, Rowland Hill Street, London, United Kingdom (G.B.); and University of Antwerp, Groenenborgercampus, Department of Veterinary Sciences, Laboratory of Cell Biology & Histology/Core Facility Biomedical Microscopic Imaging, Antwerpen, Belgium (I.B., D.A., J.-P.T.)

Abstract	834
I. Introduction	835
II. Lung and intrapulmonary airways	835
A. Epithelial lining	835
1. Alveolar and transport epithelium	835
2. Ciliated cells	838
3. Goblet cells	839
4. Neuroepithelial bodies	839
B. Immune cells	840
1. Alveolar macrophages	840
2. Lung dendritic cells	842
3. Mast cells	842
C. Airway smooth muscle	842
D. Innervation	843
1. Nonadrenergic/noncholinergic neurotransmission	843
2. Sensory nerves, reflex activity, and pain	843
E. Vasculature	843
III. Trachea	847
A. Epithelial cells	847
B. Smooth muscle and its innervation	847
IV. Nasal mucosa	848
V. Central control of respiration	848
VI. Diseases of the airways	850
A. Asthma	850
B. Chronic obstructive pulmonary disease	852
C. Airway infections	853
D. Lung injury	854
E. Cystic fibrosis	855
F. Lung cancer	857
G. Rhinosinusitis	857
H. Pulmonary hypertension	858
VII. Concluding comments	858
References	858

Abstract—Evidence for a significant role and impact of purinergic signaling in normal and diseased

Address correspondence to: Geoffrey Burnstock, Autonomic Neuroscience Centre, University College Medical School, Royal Free Campus, Rowland Hill Street, London NW3 2PF, UK. E-mail: g.burnstock@ucl.ac.uk

This article is available online at <http://pharmrev.aspetjournals.org>.
<http://dx.doi.org/10.1124/pr.111.005389>.

airways is now beyond dispute. The present review intends to provide the current state of knowledge of the involvement of purinergic pathways in the upper and lower airways and lungs, thereby differentiating the involvement of different tissues, such as the epithelial lining, immune cells, airway smooth muscle, vasculature, peripheral and central innervation, and neuroendocrine system. In addition to the vast number of well illustrated functions for purinergic signaling in the healthy respiratory tract,

increasing data pointing to enhanced levels of ATP and/or adenosine in airway secretions of patients with airway damage and respiratory diseases corroborates the emerging view that purines act as clinically important mediators resulting in either proinflammatory or protective responses. Purinergic signaling has been implicated in lung injury and in the pathogenesis of a wide range of respiratory disorders and diseases, including asthma, chronic ob-

structive pulmonary disease, inflammation, cystic fibrosis, lung cancer, and pulmonary hypertension. These ostensibly enigmatic actions are based on widely different mechanisms, which are influenced by the cellular microenvironment, but especially the subtypes of purine receptors involved and the activity of distinct members of the ectonucleotidase family, the latter being potential protein targets for therapeutic implementation.

I. Introduction

A seminal article describing the potent actions of adenosine compounds on the heart and blood vessels was published by Drury and Szent-Györgyi in 1929. Many years later, ATP was proposed as the transmitter responsible for nonadrenergic, noncholinergic neurotransmission in the gut and bladder and the term 'purinergic' was introduced by Burnstock in 1972. Early resistance to this concept seemed to stem from the fact that ATP was recognized first for its important intracellular roles in many biochemical processes, and the intuitive feeling was that such a ubiquitous and simple compound was unlikely to be used as an extracellular messenger.

Implicit in the concept of purinergic signaling was the existence of purinergic receptors, mediating potent actions of extracellular ATP on many different cell types (Burnstock and Knight, 2004). Purinergic receptors were first defined in 1976; 2 years later, a basis for distinguishing two types of purinoceptor, identified as P1 and P2 (for adenosine and ATP/ADP, respectively), was proposed (Burnstock, 1978). At approximately the same time, two subtypes of the P1 (adenosine) receptor were recognized, but it was not until 1985 that a pharmacological basis for distinguishing two types of P2 receptor (P2X and P2Y) was found (Burnstock and Kennedy, 1985). In 1993, the first G protein-coupled P2 receptors were cloned, and a year later, two ion-gated receptors were cloned. In 1994, Abbracchio and Burnstock, on the basis of molecular structure and transduction mechanisms, proposed that purinoceptors should belong to two major families: a P2X family of ligand-gated ion channel receptors and a P2Y family of G protein-coupled purinoceptors. This nomenclature has been widely adopted, and seven P2X subtypes and eight P2Y receptor subtypes are currently recognized, including receptors that are sensitive to pyrimidines as well as purines (Ralevic and Burnstock, 1998; Burnstock, 2007b).

It was assumed for some time that the only source of extracellular ATP acting on purinoceptors was damaged or dying cells, but it is now recognized that ATP release from healthy cells is a physiological mechanism (see Burnstock, 1999). ATP is released as a cotransmitter from both peripheral and central neurons but also from many non-neuronal cell types during mechanical deformation in response to shear stress, stretch, or osmotic

swelling, as well as hypoxia and stimulation by various agents. The precise transport mechanism(s) involved in ATP release remain under debate. There is compelling evidence for Ca^{2+} -dependent exocytotic vesicular release of ATP from nerves, but for ATP release from non-neuronal cells, various transport mechanisms have been proposed, including ATP-binding cassette transporters, connexin or pannexin hemichannels, or possibly plasmalemmal voltage-dependent anion channels, as well as vesicular release (see Bodin and Burnstock, 2001; Lazarowski, 2012). ATP released from cells is regulated by a number of proteins for which the catalytic site is located on the outer side of the plasma membrane (Yegutkin et al., 2008). The CD39 family consists of ectonucleoside triphosphate diphosphohydrolases that hydrolyze nucleoside 5'-tri- and diphosphates. Another family of enzymes with three subtypes includes ectonucleotide pyrophosphatase/phosphodiesterases with a broad substrate specificity. Alkaline phosphatases are also used, and ecto-5'-nucleotidase (CD73), which breaks down AMP to adenosine. ATP-induced Ca^{2+} wave propagation occurs widely in glial cells, osteoclasts, chondrocytes, epithelial cells, fibroblasts, and cancer cells and usually involves P2Y receptors (see, for example, Gallagher and Salter, 2003). Selective agonists and antagonists to many of the P1, P2X, and P2Y receptor subtypes are now available (see Burnstock, 2012).

A number of reviews on different aspects of purinergic signaling in the airways are available: extracellular purinergic regulation of airways secretion and ciliary activity (Braiman et al., 2000; Rooney, 2001; Shimura, 2001; Leipziger, 2003; Schwiebert and Zsemberly, 2003; Davis and Lazarowski, 2008; Barth and Kasper, 2009; Lazarowski and Boucher, 2009; Dietl et al., 2010); nervous control of the airways (Burnstock, 1988; Wegner, 2001; Brouns et al., 2006; Taylor-Clark and Udem, 2006; Funk et al., 2008); purinergic signaling in airway diseases (Ahmad et al., 2006; Spicuzza et al., 2006; Ackland et al., 2007); and pulmonary vasculature (Liu and Barnes, 1994). Here, we intend to provide a more comprehensive overview of airway-related purinergic signaling.

II. Lung and Intrapulmonary Airways

A. Epithelial Lining

1. *Alveolar and Transport Epithelium.* Evidence for regulation of surfactant secretion from alveolar type II

(ATII¹) cells by ATP and ADP acting on P2 receptors was presented in 1986 (Rice and Singleton, 1986). Later it was suggested that P2Y receptors mediated the mobilization of $[Ca^{2+}]_i$ involved in surfactant secretion (Rice and Singleton, 1987, 1989) but that A₂ receptors were also implicated after breakdown of ATP to adenosine (Griese et al., 1991). Plasma membrane ADP activity was identified in rat lung (Dawson et al., 1986). ATP, ADP, AMP, and adenosine were shown to stimulate phosphatidylcholine secretion in primary cultures of ATII pneumocytes (Gilfillan and Rooney, 1987). Both P1 and P2 receptor signal transduction was identified in ATII cells (Warburton et al., 1989a), and ATP-induced surfactant secretion seemed to be mediated via activation of protein kinase C (Chander et al., 1995). ATP was shown to be present in bronchoalveolar lavage in concentrations sufficient to mediate surfactant phospholipid secretion in vivo (Rice et al., 1989). Pulmonary surfactant is a complex lipoprotein mixture, and ATP was shown to stimulate the synthesis of surfactant protein A in rat lung (Wali et al., 1993). Luminal ATP activates an apical P2 receptor that stimulates K⁺ secretion across cultured human airways epithelium (Clarke et al., 1997). Multiple modes of regulation of Cl⁻ secretion from airway epithelial cells by extracellular ATP, UTP, and adenosine were identified (Stutts et al., 1994; Barker and Gatzky, 1998; Poulsen et al., 2006). Ca²⁺-dependent release of arachidonic acid from airway epithelium in response to P2U (i.e., P2Y₂ and/or P2Y₄) receptor activation by UTP and ATP was reported (Lazarewski et al., 1994). A P2U receptor was cloned from rat ATII cells and shown to be functional by expression in an unrelated cell line (Rice et al., 1995). Besides the potential involvement in secretion by directly increasing $[Ca^{2+}]_i$ via P2Y receptors, extracellular ATP was also shown to lead to membrane depolarization and activation of L-type Ca²⁺ channels in the alveolar epithelial cell line L2 (Dietl et al., 1995). It was suggested that the resulting "near membrane $[Ca^{2+}]_i$ hotspots" might play an important role in surfactant exocytosis. Depending on the substrate conditions, P2Y₂ receptors were identi-

fied in distal lung epithelial cells isolated from fetal rats (Clunes et al., 1998).

Release of ATP from lung epithelial cells was detected with bioluminescence (Taylor et al., 1998). ATP release to mechanical and osmotic stimuli were quantitated using a capillary cell culture system with Calu-3, a human lung cell line (Guyot and Hanrahan, 2002). Both elevating apical flow rate and reducing osmolarity strongly increased ATP release. Nucleotide release provides a mechanism for airway surface liquid homeostasis (Lazarewski et al., 2004). Mechanical stress results in the local release of ATP, which produces a wave of increasing $[Ca^{2+}]_i$ that spreads to other cells, mediated via P2Y receptors (Hansen et al., 1993; Homolya et al., 2000). Alveolar type I (ATI) epithelial cells release 4-fold more ATP in response to stretch than ATII cells, and it was concluded that ATI cells are mechanosensors in the lung and that paracrine stimulation of ATI cells by ATP released from these cells plays a role in regulating surfactant secretion (Patel et al., 2005). Cell deformation by surface tension forces in the proximity of the air-liquid interface causes Ca²⁺-dependent exocytotic ATP release (Ramsingh et al., 2011), which may provide a simple sensing mechanisms to detect low content of surface liquid or surfactant, leading via ATP to compensatory secretion of fluid, mucus, and surfactant. Released ATP exerts an autocrine regulation of epithelial NaCl absorption, mainly by inhibiting the amiloride-sensitive epithelial Na⁺ channel (ENaC) and ion conductance via a P2Y receptor-dependent increase in Ca²⁺ (Poulsen et al., 2005; Bao et al., 2007). Evidence that the released ATP reached sufficient concentrations at the airway surface to activate P2Y₂ receptors has been provided, supporting the view that this is a physiological mechanism for epithelial cell volume regulation (Okada et al., 2006). UTP, as well as ATP, is released from ATII-like A549 epithelial cells, probably by exocytosis, because the secretion is Ca²⁺-dependent (Tatur et al., 2008). Thrombin promotes release of ATP from lung A549 epithelial cells through the coordinated activation of Rho- and Ca²⁺-dependent signaling pathways; the ATP release was inhibited by connexin hemichannel blockers (Seminario-Vidal et al., 2009). The contribution of pannexin 1 to ATP release from airway epithelium has also been suggested (Ransford et al., 2009). Van Scott et al. (1995) claimed that P2U receptors (i.e., P2Y₂ and/or P2Y₄) mediate regulation of ion transport across nonciliated bronchiolar epithelial (Clara) cells.

Mouse airway epithelium was reported to express P2Y₂ protein, where the immunostaining was most prominent in Clara cells (De Proost et al., 2009; Brouns et al., 2012; Fig. 1). mRNA for P2Y₂, P2Y₄, and P2Y₆ receptors were coexpressed in epithelial cell lines (IHAEo, 16HBE14o, and A549) derived from human lungs (Communi et al., 1999). P2Y₂ receptor-mediated inhibition of ion transport in distal lung epithelial cells has been reported (Ramminger et al., 1999). Extracellular

¹Abbreviations: α,β -meATP, α,β -methylene ATP; 5-HT, 5-hydroxytryptamine; Ap₄A, diadenosine pentaphosphate; ARDS, acute respiratory stress syndrome; ATII, alveolar type II; ATP γ S, adenosine-5'-(γ -thio)-triphosphate; BALF, bronchoalveolar lavage fluid; CF, cystic fibrosis; CFTR, cystic fibrosis transmembrane conductance regulator; CGS 21680, 2-[p-(2-carboxyethyl)phenethylamino]-5'-N-ethylcarboxamidoadenosine; COPD, chronic obstructive pulmonary disease; DRG, dorsal root ganglion; ENaC, epithelial Na⁺ channel; IL, interleukin; INS37217, P¹-(uridine 5')-P⁴-(2'-deoxycytidine 5')tetraphosphate; LPS, lipopolysaccharide; MRS2365, (N)-methanocarba-2-methylthio-ADP; NA, noradrenaline; NANC, nonadrenergic/noncholinergic; NBD, nucleotide-binding domain; NEB, neuroepithelial body; NTS, nucleus tractus solitarius; PBC, pre-Bötzinger complex; PECAM, platelet endothelial cell adhesion molecule; PMN, polymorphonuclear leukocyte; RT-PCR, reverse transcription polymerase chain reaction; SMAR, smooth muscle-associated airway receptor.




FIG. 1. Double immunostaining for P2Y₂ receptors (red fluorescence) and Clara cell-specific protein (CCSP; green fluorescence) in a mouse intrapulmonary airway. Left, P2Y₂ expression is seen in most cells of the airway epithelial layer (E) but with varying intensities. Center, CCSP staining of Clara cells. Right, combination of the red and green channels reveals the most prominent P2Y₂ expression in Clara cells. L, airway lumen.

nucleotides induce cyclooxygenase-2 up-regulation and prostaglandin E₂ production in A549 cells (Marcat et al., 2007a). Luminal P2Y₂ receptors on epithelial cells in distal bronchi of pigs regulate Cl⁻ secretion and Na⁺ absorption (Inglis et al., 1999). ATP, acting through P2Y receptors, regulates the secretion of ions, mucin, and surfactant phospholipids by rat respiratory epithelium (Kishore et al., 2000). Nucleotides induce interleukin (IL)-6 release from human airway epithelia via P2Y₂ receptors and p38 mitogen-activated protein kinase-dependent pathways (Doullet et al., 2006). Diadenosine pentaphosphate (Ap₄A) increased [Ca²⁺]_i in HBE1 cells derived from human bronchial epithelial cells, probably acting via P2Y₂ receptors (Laubinger et al., 2001). Multiple P2Y receptor subtypes seem to be present on both apical and basolateral membranes of human airway-derived Calu-3 cells (Chambers et al., 2002), including P2Y₁ receptors (Son et al., 2004). UTP stimulates alveolar fluid clearance in the distal airspaces of rat lungs (Sakuma et al., 2004). P2Y₁₄ receptor mRNA was shown to be expressed in human lung epithelial cells and in the lung epithelial cell lines A549 and BEA5-2B (Müller et al., 2005). Ap₄A, ATP, and UDP induce arachidonic acid release and nitric-oxide synthase expression in A549 lung epithelial cells via P2Y receptors (Laubinger et al., 2006). Expression and functional P2Y₆ receptors were demonstrated in primary cultures of normal human bronchial epithelial cells; they were shown to be located only on the apical membranes, and stimulation with UDP-induced Cl⁻ secretion (Dulong et al., 2007). 16HBE14o cells were shown to express P2Y₁, P2Y₂, P2Y₄, and P2Y₆ receptors under different culture conditions and P2Y₆ receptors were present on both apical and basolateral membranes that regulate Cl⁻ secretion (Wong et al., 2009). However, the P2Y₆ receptors seem to be differentially coupled to different downstream signaling pathways at the two sites.

A₁, A_{2A}, A_{2B}, and A₃ receptors are expressed in both ATI and ATII cells in rats and mice, and further findings suggested that physiological concentrations of adenosine

allow the alveolar epithelium to counterbalance active Na⁺ absorption with Cl⁻ efflux, via A₁ receptors (Factor et al., 2007). Adenosine increased phosphatidylcholine secretion in rat-cultured ATII pneumocytes via A₂ receptors (Gilfillon and Rooney, 1987) and resulted in A₁ receptor-mediated inhibition of surfactant secretion (Gobran and Rooney, 1990). A_{2B} receptors are highly expressed on murine ATII epithelial cells (Cagnina et al., 2009). Apical adenosine regulates basolateral Ca²⁺-activated K⁺ channels in Calu-3 cells (Wang et al., 2008). It was claimed that volume-sensitive anion channels (K⁺ and Na⁺) are regulated by an autocrine mechanism involving swelling-induced ATP release and then hydrolysis to adenosine to act on P1 receptors (Musante et al., 1999; Szkotak et al., 2001). Control of cystic fibrosis transmembrane conductance regulator (CFTR)-mediated airway surface ligand secretion by adenosine and ENaC has been claimed (Tarran, 2007). Adenosine promotion of cellular migration in bronchial epithelial cells is mediated by A_{2A} receptors via activation of protein kinase A (Allen-Gipson et al., 2007). Adenosine induces fibronectin expression in A549 lung epithelial cells, and it was suggested that this might be involved in tissue remodeling (Roman et al., 2006). Elevated lung adenosine can lead to abnormal alveolar development by interfering with cell proliferation and apoptosis (Banerjee et al., 2004). Histamine-induced bronchospasm is potentiated by adenosine (Nieri et al., 1997), possibly via A₁ receptors (Breschi et al., 1998).

Electrophysiological data and mRNA analysis of primary cultures of human and mouse pulmonary epithelia revealed the presence of P2X receptors on both the apical and basolateral membranes of well differentiated cells, occupation of which stimulated Cl⁻ transport (Taylor et al., 1999). Expression of P2X4 and P2X6 receptor isoforms in bronchial epithelium has been shown by in situ hybridization (Buell et al., 1996; Collo et al., 1996). Biochemical evidence was presented to show that P2X4 receptors on human 16HBE14o airway epithelial

cells mediated ATP-gated calcium entry channels that induced a sustained increase in $[Ca^{2+}]_i$ (Zsembery et al., 2003). Molecular and biochemical data show coexpression of P2X4 and P2X6 subtypes in 16HBE14o and CF (IB3-1) human bronchial epithelial cells (Liang et al., 2005). They additionally showed that novel transient lipid transfection-mediated delivery of small-interference RNA fragments specific to P2X4 and P2X6 receptors into IB3-1 cells inhibited extracellular ATP-induced Ca^{2+} entry. P2X4 receptors have been shown to be expressed on exocytotic vesicles known as lamellar bodies in ATII cells, and when lamellar bodies fuse with the plasma membrane, activated P2X4 receptor-mediated local Ca^{2+} signals facilitate fusion pore expansion and subsequent surfactant secretion (Miklavc et al., 2011). P2X7 receptors were found to be expressed on the ATI-like E10 cell line (Barth et al., 2007). There seems to be an interaction between P2X7 receptor protein and Caveolin-1 in lipid rafts in E10 cells (Barth et al., 2007, 2008). Reverse transcription polymerase chain reaction (RT-PCR) studies revealed that NT-1 and A549 cells expressed P2X4, P2X5, and P2X6 receptor mRNA, whereas P2X7 receptors were detected only on NT-1 cells (Théâtre et al., 2009). Activation of P2X7 receptors increases surfactant secretion by releasing ATP from ATII cells and subsequently stimulating P2Y₂ receptors on the ATII cells (Mishra et al., 2011). In mouse lungs, P2X7 protein seems to be expressed on ATI cells (unpublished observations) (Fig. 2).

Hydrogen peroxide is found in exhaled breath and is produced by airway epithelia. Apical production of hy-

drogen peroxide by human epithelial cells was stimulated by ATP but not ADP (Forteza et al., 2005).

A novel type of ATP-diphosphohydrolase was identified in bovine lung (Picher et al., 1993; Sévigny et al., 1997). Adenylate kinase and nucleoside diphosphokinase contribute to nucleotide metabolism on human airway surfaces (Donaldson et al., 2002).

2. Ciliated Cells. Mucociliary clearance serves as a primary host defense mechanism for removing dust and microorganisms from the lungs. ATP released mechanically from airway epithelial cells has a ciliostimulatory action in animal and human respiratory mucosa (Saano et al., 1991). In human and ovine ciliated airway epithelial cells, purinergic stimulation increases both $[Ca^{2+}]_i$ and ciliary beat frequency, and the regulator of G protein signaling protein 2 modulates these actions (Nlend et al., 2002). Extracellular ATP was shown to directly gate a cation-selective channel in rabbit ciliated epithelial cells (Korngreen et al., 1998), indicating the presence of P2X receptors. P2Y₂ receptor agonists increased lung mucociliary clearance in sheep (Sabater et al., 1999). The number of immunohistochemically positive cells for ATPase strongly correlated with ciliary motility in vitro, and cultures without ciliary activity did not exhibit ATPase staining (Schütz et al., 2002). Purinergic stimulation of ovine airway epithelial cells for 1 min transiently increased $[Ca^{2+}]_i$ and ciliary beat frequency, but stimulation of human epithelial cells led to an increase in ciliary beat frequency that outlasted the calcium transient by at least 20 min, perhaps indicating that ATP is triggering a signaling cascade (Lieb et al.,




FIG. 2. Immunostaining for P2X7 receptors (red fluorescence) in an alveolar region of the mouse lung, revealing a clear P2X7 expression on the surface membrane of alveolar type I cells. A, alveolar spaces.

2002). cAMP and cGMP augment ATP-driven ciliary beat frequency increases at the level of isolated axonemes (Wyatt et al., 2005).

Airway ciliated epithelial cells have been shown to express the calcium-sensitive transmembrane adenylyl cyclase isoforms AC1 and AC8 on the apical membrane, suggesting that they are involved in the purinergic regulation of ciliary beating (Nlend et al., 2007). Extracellular sodium regulates airway ciliary motility by inhibiting P2X receptor activation (Ma et al., 1999). In addition to ectonucleotidases that degrade ATP mechanically released from epithelial cells, ecto adenylyl kinase mediates adenosine reformation into nucleotides, thereby prolonging the effects of ATP and ADP on airway epithelial surfaces (Picher and Boucher, 2003).

3. *Goblet Cells.* Mucin release from airway goblet cells is stimulated by ATP acting via P2 receptors (Kim and Lee, 1991). P2U (i.e., P2Y₂ and/or P2Y₄) receptors were later implicated in ATP and UTP regulation of mucin secretion in SPOC1 cells, an airway goblet cell line (Abdullah et al., 1996, 2003), and in differentiated human bronchial cells (Kemp et al., 2004). Inhaled ATP causes mucin release from goblet cells in rats (Shin et al., 2000) and humans (Kemp et al., 2004). UTP can enhance both mucin secretion and mucin gene expression through different signaling pathways in cultures of human tracheobronchial tissues (Chen et al., 2001). P2Y₂ receptors provide a major pathway for the regulation of mucin secretion by ATP and UTP acting on the apical membranes of goblet cells (Huang et al., 2001) via phospholipase C pathways (Conway et al., 2003). The increase in [Ca²⁺]_i is due to mobilization of intracellular stores by activation of P2Y receptors (Rossi et al., 2007). ATP is secreted from goblet airway cells during Ca²⁺-regulated mucin exocytosis, suggesting that ATP is released from mucin granules to signal neighboring ciliated cells to promote ion/water transport, perhaps via adenosine A_{2B} receptors, for hydration of secreted mucins into the airway lumen (Kreda et al., 2008). This group later reported that mucin granules contained ADP and AMP levels that exceeded ATP levels by nearly 10-fold (Kreda et al., 2010). Okada et al. (2011) published in vitro data suggesting that increased goblet cell nucleotide release and resultant adenosine accumulation might provide compensatory mechanisms to hydrate mucins by paracrine stimulation of ciliated cell ion and water secretion and to modulate inflammatory responses.

4. *Neuroepithelial Bodies.* Pulmonary neuroepithelial bodies (NEBs) are densely innervated groups of neuroendocrine cells that are invariably surrounded by a population of presumed airway epithelial stem cells, referred to as Clara-like cells. NEBs have been suggested to have several functions in the regulation of physiological processes in the lungs during prenatal, perinatal, and postnatal life (Adriaensen et al., 2003; Linnoila, 2006), but the strongest evidence to date im-

plicates them in airway oxygen sensing (Cutz et al., 2003) involved in the autonomic regulation of breathing.

Vagal sensory nerve terminals, arising from the nodose ganglion, in rat pulmonary NEBs were first shown to express P2X3 receptors by Brouns et al. (2000). These vagal nerve fibers are myelinated but lose their myelin sheaths just before branching and protruding intraepithelially between NEB cells (Brouns et al., 2003b). They also showed that these fibers were not activated by capsaicin and were immunoreactive to calbindin. A second sensory nerve population, arising from dorsal root ganglia (DRG), innervated NEBs and expressed calcitonin gene-related peptide and substance P but not P2X3 receptors. In a developmental study, vagal nodose sensory nerve endings expressing P2X3 receptors were the first to selectively innervate NEBs, from fetal stage GD16 onward. This suggested a role for purinergic regulation of NEBs during intrauterine life, when the density of NEBs was high, suggesting that NEBs may act as chemoreceptors for hypoxia before the carotid body chemoreceptors, which are known to be relatively immature at birth (Brouns et al., 2003a). This group later showed that approximately 40% of NEBs were innervated by vagal sensory fibers, approximately 50% by calcitonin gene-related peptide-immunoreactive DRG sensory fibers, and approximately 10% by intrinsic pulmonary nitrenergic nerves (Van Genechten et al., 2004). Another research group demonstrated that both P2X2 and P2X3 receptor immunoreactivity on hamster NEBs was identified with 5-hydroxytryptamine (5-HT), a marker for NEBs (Fu et al., 2004). They further showed 1) that ATP induced a nondesensitizing inward current typical of P2X2 receptor-mediated responses, whereas α,β -methylene ATP (α,β -meATP), an agonist for P2X3 receptors, produced a slowly desensitizing inward current, perhaps indicating involvement of a P2X2/3 heteromultimer, and 2) that hypoxia and ATP induced 5-HT release from NEB cells.

In rodent studies of the neurochemical characterization of sensory receptor terminals in NEBs, it was shown that subpopulations of myelinated vagal sensory fibers expressing P2X2/3 receptors supplied NEBs (Fig. 3), and schematics were presented (Brouns et al., 2009, 2012; Fig. 4).

Quinacrine histochemistry indicated high amounts of vesicular ATP in NEB cells (Brouns et al., 2000; De Proost et al., 2009), and live cell imaging, using "reporter patching" with human embryonic kidney 293 cells overexpressing P2X2 receptors, indicated quantal ATP release after depolarization and paracrine interactions between NEB and Clara-like cells. It was concluded that in addition to ATP acting on P2X3 receptors on vagal sensory nerve endings in NEBs, local paracrine purinergic signaling in the NEB microenvironment, in which ATP released from activated NEB cells induces a secondary activation of the surrounding stem cell-like Clara-like cells via P2Y₂ receptors, may be important for airway




FIG. 3. Triple immunostaining for P2X2 receptors (green fluorescence), P2X3 receptors (red fluorescence), and PGP9.5 (blue fluorescence; marker for nerve fibers and NEB cells) in a mouse intrapulmonary airway. P2X2 and P2X3 receptors are invariably coexpressed in vagal sensory nerve terminals in the NEB (arrowheads).

epithelial regeneration after injury and/or the pathogenesis of small-cell lung carcinomas.

B. Immune Cells

Extracellular ATP serves as a danger signal to alert the immune system of tissue damage. P2 receptors are widely expressed in cells of the immune system, as illustrated mainly during the last decade for many species, and are now believed to underlie various functions in health and disease. In addition, adenosine is a signaling nucleoside that has been implicated in the regulation of immune function and may serve both pro- and anti-inflammatory functions in pulmonary disease.

1. Alveolar Macrophages. Extracellular adenine nucleotides have been known for many years to stimulate a Ca^{2+} -dependent respiratory burst (superoxide production) in alveolar macrophages that is mediated via purinergic receptors (Murphy et al., 1993). P2Y receptors mediate up-regulation of inducible nitric-oxide synthase mRNA and protein in alveolar macrophages (Greenberg et al., 1997).

Bovine alveolar macrophages, as well as B and T lymphocytes, express P2X7 receptors (Smith et al., 2001). Rat alveolar macrophages (NR8383) express functional P2X4 and phospholipase C-coupled P2Y₁ and P2Y₂ receptors (Bowler et al., 2003), and treatment with ATP results in an increased reactive oxygen species-mediated stress response and secretion of proinflammatory cytokines (Cruz et al., 2007). ATP can release cytokines and/or chemokines from rat alveolar macrophages (NR8383), which lack the P2X7 receptor, via P2Y₂ receptors (Stokes and Surprenant, 2007). In rat alveolar macrophages, freshly isolated from bronchoalveolar lavage fluid (BALF), the ATP-induced Ca^{2+} -release from intracellular stores seems to be reduced by nicotine via nicotinic acetylcholine receptors (Mikulski et al., 2010).

Although RT-PCR revealed transcripts for P2X1, P2X3, P2X4, P2X5, and P2X7 receptors in mouse alveolar macrophages, functional and pharmacological data point to a major participation of P2X4 and P2X7 recep-

tors in pulmonary immune function (Kessler et al., 2011). Protein expression was confirmed for P2X1 and P2X4 receptors in control mouse lungs (unpublished observation) (Fig. 5).

Primary human alveolar macrophages obtained from BALF were shown to express mRNA for a large panel of purinoreceptors, namely P2X1, P2X4, P2X5, and P2X7, and P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁, P2Y₁₃, and P2Y₁₄. Moreover, extracellular ATP induces Ca^{2+} transients and increased IL-1 β and IL-6 secretion via P2X receptors, supporting the hypothesis that purinoreceptors may be involved in the pathogenesis of inflammatory lung diseases (see section VI) (Myrtek et al., 2008).

P2X4 receptor expression of human and rat alveolar macrophages is regulated by physiological stimuli, whereby P2X4 receptors become functional in response to initial phagocytic activity, but return to nonfunctional during sustained classic macrophage activation (Stokes and Surprenant, 2009). Endogenous extracellular ATP induces an up-regulation of purinergic receptors on neutrophils and alveolar macrophages. ATP also stimulates the recruitment of neutrophils to the lungs via P2Y₂ receptors and, as such, contributes to smoke-induced lung inflammation and development of emphysema (Cicko et al., 2010; Lommatzsch et al., 2010; see further on). P2X7 receptors were reported on macrophages in human BALF from patients with chronic obstructive pulmonary disease (COPD) (see section VI) (Lommatzsch et al., 2010).

Extracellular ATP, via the up-regulation of P2X7 receptors—as observed on alveolar macrophages in BALF and airway and blood neutrophils and eosinophils—has been implicated in the pathogenesis of airway inflammation (Müller et al., 2011), smoke-induced lung inflammation and emphysema (Lucatelli et al., 2011) in both rodents and humans. In mouse lungs, A₁ adenosine receptors were reported to be predominantly expressed on alveolar macrophages and to play an anti-inflammatory and/or protective role in adenosine-dependent pulmonary injury (Sun et al., 2005). Although mRNA for all




FIG. 4. Schematic representation of the main innervation of airway smooth muscle and of the sensory innervation of complex NEB receptors in rat lungs. Known characteristics of the represented neuronal populations are included in the scheme in the same color as the respective structures. The pulmonary NEB cells (yellow) express ATP, calcitonin gene-related peptide (CGRP), calbindin (CB), 5-HT, calcitonin, vesicular acetylcholine transporter (VChT), P2X₂ purinergic receptors, and Ca_v2.1 voltage-gated calcium channels. Clara-like cells (dark gray) can be distinguished by their location and immunoreactivity for Clara cell-specific protein, P2Y₂, and Ca_v2.3. The lower part of the scheme shows airway smooth muscle that receives nerve terminals from postganglionic parasympathetic neurons located in an airway ganglion, and laminar nerve terminals of vagal SMARs (dark purple) that intercalate between the smooth muscle cells. The center of the scheme represents a pulmonary NEB and its extensive interactions with sensory nerve terminals. The upper left part shows the vagal nodose afferent, and the upper right part the dorsal root afferents. CRT, calretinin; MB, myelin basic protein; nNOS, neuronal nitric-oxide synthase; SMA, smooth muscle actin; TRPV, transient receptor potential vanilloid; VGLUT, vesicular glutamate transporter; VIP, vasoactive intestinal peptide. [Reproduced from Brouns I, Pintelon I, Timmermans JP, and Adriaenssens D (2012) Novel insights in the neurochemistry and function of pulmonary sensory receptors. *Adv Anat Embryol Cell Biol* 211:1–115, vii. Copyright © 2012 Springer. Used with permission.]




FIG. 5. Double immunostaining for P2 receptors (red fluorescence) and CD169 (green fluorescence), as a marker for alveolar macrophages (arrows) in control mouse lungs—macrophages were not activated. Left, P2X1 expression in CD169 positive alveolar macrophages. Right, P2X4 staining of a macrophage; note that other alveolar cells, presumably alveolar type II cells (arrowheads), appear to express P2X4. A, alveolar spaces.

adenosine receptors (A_1 , A_{2A} , A_{2B} , A_3) could be detected in human alveolar macrophages, only A_{2A} receptor expression was strongly up-regulated after lipopolysaccharide (LPS) treatment. In addition, adenosine-mediated inhibition of tumor necrosis factor- α and some chemokines seemed to be mediated mainly via A_{2A} adenosine receptors (Buenestado et al., 2010).

2. Lung Dendritic Cells. P2 receptors are expressed by human and mouse airway dendritic cells, and purinergic signaling is now believed to have a key role in allergen-driven airway inflammation (e.g., asthma) (see section VI) as a result of the recruitment and activation of lung myeloid dendritic cells that induce Th2 responses (Idzko et al., 2007). ATP was shown to trigger 1) migration to the airways of monocyte-derived dendritic cells and eosinophils and 2) production of reactive oxygen species involved in asthmatic airway inflammation via P2Y₂ receptors (Müller et al., 2010).

3. Mast Cells. Adenosine was reported to both inhibit and potentiate IgE-dependent histamine and 5-HT release from lung mast cells; A_2 receptors are involved (Hughes et al., 1984; Ott et al., 1992). Immunological identification of A_{2B} receptors was reported in human lung mast cells (Feoktistov et al., 2003). Adenosine closes the K^+ channel in human lung mast cells and inhibits their migration via A_{2A} receptors (Duffy et al., 2007). A_3 receptors were shown to play an important role in adenosine-mediated lung mast cell degranulation (Reeves et al., 1997; Zhong et al., 2003). ATP mediates anti-IgE-induced release of histamine from human lung mast cells, probably via P2Y₁ and/or P2Y₂ receptors (Schulman et al., 1999), but they have also been reported to express functional P2X1, P2X4, and P2X7 receptors (Wareham et al., 2009).

C. Airway Smooth Muscle

ATP produced strong contractions of human lung bronchiolar segments, whereas adenosine elicited only very weak contractions (Finney et al., 1985). ATP stimulated Ca^{2+} oscillations and contraction of airway smooth muscle of mouse lung slices, probably via P2Y₂ or P2Y₄ receptors (Bergner and Sanderson, 2002). In a later article, a more detailed analysis of the actions of ATP on human and rat bronchial smooth muscle led to the conclusion that ATP induces a transient contractile response mainly mediated by P2X4 receptors, but P2Y receptors were also present (Mounkaïla et al., 2005). RT-PCR and Western blot analyses carried out on primary cultures of human airway smooth muscle showed expression of P2Y₁, P2Y₂, P2Y₄, and P2Y₆ receptors and ATP, UTP, ADP, and UDP all produced a significant increase in $[Ca^{2+}]_i$ and contraction (Govindaraju et al., 2005).

Adenosine receptor antagonists induced bronchoconstriction in rats (Salonen et al., 1982), whereas enprofylline, another xanthine, produced bronchodilation, apparently without adenosine receptor antagonism (Lunell et al., 1983). Adenosine caused bronchoconstriction in a rat model, but analysis suggested that adenosine acts by stimulating postjunctional vagal nerve endings and mast cells (Pauwels and Van der Straeten, 1987). Inflammatory mediators inhibited by indomethacin may be involved in adenosine bronchoconstriction (Crimi et al., 1989). In perfused lungs sensitized with ovalbumin, adenosine produced bronchoconstriction, but at high concentrations, it had bronchodilator actions (Thorne and Broadley, 1992). The ADP receptor antagonist 8-phenyltheophylline blocked relaxant but not contractile responses. It has been claimed that 5-HT may mediate adenosine-induced airway contraction (Matera et

al., 1995). Adenosine-mediated bronchoconstriction was characterized to be mediated by A_1 , A_{2B} , and A_3 but not A_{2A} receptor subtypes (Pauwels and Joos, 1995). A_1 receptors mediated mobilization of Ca^{2+} in human bronchial smooth muscle cells (Ethier and Madison, 2006). Other studies suggested that xanthine-resistant A_3 receptors dominated (Thorne et al., 1996). Experiments on rabbit airway smooth muscle suggested that adenosine causes relaxation via an A_2 receptor that releases nitric oxide from the epithelium (Ali et al., 1997). Adenosine enhanced endothelium-induced bronchoconstriction in guinea pigs (Kanazawa et al., 1997). A_{2B} receptors mediate increase in cytokine release from bronchial smooth muscle cells (Zhong et al., 2004) and mediate the relaxing effects of adenosine on guinea pig airways (Breschi et al., 2007).

D. Innervation

In addition to sympathetic, parasympathetic, and sensory nerves, many neurons in intrinsic ganglia are derived from a separate population of neural crest cells (Freem et al., 2010), which suggests that, as in the gut and heart, intrinsic neurons may allow local reflex activity independent of the central nervous system (see Burnstock, 2009a).

1. *Nonadrenergic/Noncholinergic Neurotransmission.* NANC neurotransmission was recognized early to participate in autonomic control of pulmonary smooth muscle (Koelle, 1975; Diamond and O'Donnell, 1980). It is now well established that ATP is a cotransmitter both with noradrenaline (NA) in sympathetic nerves and with acetylcholine in parasympathetic nerves (see Burnstock, 2007a). ATP has been considered a NANC transmitter in inhibitory vagal nerves supplying the airways (Irvin et al., 1980, 1983; Souhrada et al., 1980; Satchell, 1982; Pauwels et al., 1990). Adenosine inhibits NANC excitatory transmission in guinea pig trachea (Kamikawa and Shimo, 1989), in retrospect probably by acting on P_1 (A_1) receptors on nerve terminal varicosities to inhibit transmitter release (Schmidt et al., 1995).

2. *Sensory Nerves, Reflex Activity, and Pain.* In canine lungs, ATP activates pulmonary vagal C fiber sensory nerve terminals, mediated by P2X receptors (Pelleg and Hurt, 1996), and causes reflex bronchoconstriction (Katchanov et al., 1998). In another study, activation of P2X receptors on vagal afferent terminals evoked Bezold-Jarisch depressor cardiorespiratory reflexes in anesthetized rats (McQueen et al., 1998). Because α,β -meATP initiated the reflex in this study, it is likely that P2X1 and/or P2X3 receptors were involved. ATP stimulated both capsaicin-sensitive and -insensitive vagal bronchopulmonary C fibers in the mouse via P2X receptors (Kollarik et al., 2003). Vagal C fibers innervating the pulmonary system are derived from cell bodies situated in both jugular (neural crest-derived) and nodose (placodal) vagal sensory ganglia. Only the nodose C-fiber population responds to P2X receptor activation (Udem

et al., 2004; Nassenstein et al., 2010). In addition, intrapulmonary myelinated nodose airway mechanosensors (A-fiber stretch receptors) have been reported to be activated by ATP via P2X receptors (Lee et al., 2004). ATP evoked an inward current in 34 of 39 pulmonary sensory neurons in nodose ganglia via P2X receptors (Ni and Lee, 2008). Evidence has been presented for both A_1 and A_{2A} receptors on vagal sensory C fibers in guinea pig lungs (Chuaychoo et al., 2006). The sensory transduction of pulmonary reactive oxygen species by capsaicin-sensitive vagal lung afferent fibers was found to be mediated by both transient receptor potential vanilloid 1 and P2X receptors (Ruan et al., 2005).

In rodents, subpopulations of myelinated vagal sensory fibers expressing P2X2/3 receptors were shown to terminate both in NEBs (see section II.4) (Brouns et al., 2000, 2009, 2012) (Figs. 3 and 4) and in airway smooth muscle as so-called smooth muscle-associated airway receptors (SMARs) (Brouns et al., 2006, 2012) (Figs. 4 and 6). Neuronal tracing from the airways, combined with immunocytochemistry, calcium imaging, and electrophysiological recording on dissociated vagal nodose airway-related neurons revealed that a considerable part of the traced neurons coexpress P2X3 and the cold-sensitive channel transient receptor potential melastatin 8 (Xing et al., 2008).

The visceral pleura has historically been considered insensitive to painful stimuli. However, more recent studies have revealed sensory fibers with complex laminar terminals that express P2X3 receptors embedded in the elastic fiber network of the rodent visceral pleura (Pintelon et al., 2007; Brouns et al., 2012; Fig. 7). These fibers probably originate from nerve cell bodies in DRGs and have been reported to mediate nociception in other visceral organs (see Burnstock, 2009b).

E. Vasculature

Evidence for release of NA and ATP as cotransmitters from sympathetic nerves supplying rabbit pulmonary arteries was reported 30 years ago (Katsuragi and Su, 1982). An important advance was made when it was shown that excitatory junctional potentials recorded in the smooth muscle of rat intrapulmonary arteries in response to sympathetic nerve stimulation were mediated by ATP (Inoue and Kannan, 1988). Adenosine, acting prejunctionally, mediates release of transmitter from sympathetic nerves supplying the rabbit pulmonary artery (Husted and Nedergaard, 1985). Perivascular nerve stimulation of rabbit pulmonary artery elicits release of NA and ATP (Mohri et al., 1993). It has been claimed that inhibitory purinergic neurotransmission is endothelium-dependent in pulmonary vessels (Liu and Barnes, 1994).

Porcine pulmonary vessels express P2 receptors that mediate the synthesis and release of prostacyclin (Hellewell and Pearson, 1984). In the isolated, blood-




FIG. 6. Triple immunostaining for P2X2 receptors (red fluorescence), P2X3 receptors (green fluorescence), and PGP9.5 (blue fluorescence) in a mouse intrapulmonary airway. P2X2/3 stained laminar terminals of a SMAR are intermingled with PGP9.5 labeled nerve fibers that are present in the airway smooth muscle bundle right beneath the epithelium (E). L, airway lumen.

perfused, and ventilated rat lung, P2X receptors activated by α,β -meATP caused vasoconstriction, whereas P2Y receptors, probably located largely on the endothelium, mediated a small vasodilator response (Liu et al., 1989; McCormack et al., 1989a). Vasoconstriction of the rat pulmonary vascular bed by UTP and ATP is resistant to suramin antagonism (Rubino and Burnstock, 1996) and is probably mediated by P2Y₄ receptors. ATP- and ADP-induced increase in $[Ca^{2+}]_i$ in rat pulmonary artery myocytes was found to be mediated by P2X and P2U (P2Y₂ and/or P2Y₄) receptors (Guibert et al., 1996; Hartley and Kozlowski, 1997). In endothelium-denuded rat pulmonary arteries, P2Y₂ and a UDP-sensitive receptor (presumably P2Y₆) were identified on myocytes (Hartley et al., 1998). Although ATP was a potent vasoconstrictor in perfused rat lung (Rubino and Burnstock, 1996), it was only a weak agonist in isolated rat pulmonary artery (Rubino et al., 1999). UTP and UDP are equipotent in inducing vasoconstriction of rat intrapulmonary arteries (Rubino et al., 1999). There is regional variation in P2 receptor expression mediating constriction in the rat pulmonary arterial circulation; P2Y re-

ceptor agonist potencies were similar in large and small vessels, but P2X receptor agonists were more potent in small arteries (Chootip et al., 2002). ATP was reported to produce vasoconstriction via P2X1 receptors in the feline pulmonary vascular bed (Neely et al., 1991). In mouse airways, arterial smooth muscle seems to express P2X1 (unpublished observations) (Fig. 8) and P2Y₁₁ protein (unpublished observations) (Fig. 9) with regional selectivity, whereas airway smooth muscle is negative.

An early article described hydrolysis of ATP to adenosine by cultures of porcine pulmonary endothelial cells and uptake of adenosine to be rapidly phosphorylated by adenosine kinase (Dieterle et al., 1978). Inhibition of ecto-ATPase by adenosine-5'-(γ -thio)-triphosphate (ATP γ S), α,β -meATP, and adenylylimidodiphosphate in bovine pulmonary artery endothelial cells has been reported (Chen and Lin, 1997). Adenosine was shown to mediate vasodilation via P1 receptors but apparently does not mediate the pulmonary vasodilator response to ATP in the feline pulmonary vascular bed (Neely et al., 1989). As mentioned above, ATP produces vasoconstriction via P2X1 recep-




FIG. 7. P2X3 receptor expression (green fluorescence) in a typically structured sensory visceral pleura receptor (VPR) of a mouse lung. PGP9.5 immunostaining (red fluorescence) labels all nerve fibers and terminals.




FIG. 8. Double immunostaining for P2X1 receptors (red fluorescence) and PECAM (green fluorescence; marker for endothelial cells) in small and terminal airways of mouse lungs. Left, P2X1 expression in smooth muscle cells of a small (resistance) artery. Note the abrupt end of the expression in a small vessel that branches off (arrows). Right, additional PECAM staining clearly reveals P2X1 expression in arterial smooth muscle cells, but not in the endothelial cells in this volume projection of confocal images of another small artery. In addition, the airway smooth muscle, which is invisible in these images but known to be invariably located just beneath the epithelial layer (E), appears P2X1-negative. A, alveolar spaces; L, airway lumen.

tors but also in part by the action of adenosine on A_1 -like receptors after its breakdown in pulmonary arteries of cats (Neely et al., 1991), guinea pigs (Wiklund et al., 1987), and sheep (Biaggioni et al., 1989). However, P1 receptors mediate relaxation of bovine bronchial arteries (Alexander and Eyre, 1985). In human pulmonary arter-

ies, the adenosine vasodilator effects are mediated by A_2 receptors (McCormack et al., 1989b). In juvenile rabbit pulmonary arteries and veins, the relaxations to adenosine were found to be endothelium-dependent (Steinhorn et al., 1994). Adenosine, acting via A_2 receptors, mediates vasodilation in the perfused adult rabbit lung




FIG. 9. Multiple immunostaining for P2 receptors (red fluorescence), PECAM (green fluorescence; marker for endothelial cells), and PGP9.5 (blue fluorescence; marker for nerve fibers and NEBs) in a mouse intrapulmonary airway. P2Y₁₁ expression is strong in smooth muscle cells of a large artery in the airway wall but absent from the airway smooth muscle that is invisible in this image but known to be invariably located just beneath the epithelial layer (E). Note that also the NEB (open arrowhead), nerve fibers, and endothelial cells are P2Y₁₁ negative. L, airway lumen.

(Pearl, 1994), and it was asserted that this dilation was due to direct action on the vascular smooth muscle (El-Kashef et al., 1999). Adenosine mediates both contractile and relaxant effects on the main pulmonary artery of guinea pigs via A_1 and A_{2B} receptors, respectively (Szentmiklósi et al., 1995). In addition, in the feline pulmonary vascular bed, adenosine causes vasoconstriction via A_1 receptors and vasodilation via A_2 receptors but apparently not by releasing NO (Cheng et al., 1996). LPS binds to and activates A_1 receptors on human pulmonary artery endothelial cells to induce the release of IL-6 and thromboxane A_2 (Wilson and Batra, 2002) and may contribute to the pathophysiology of acute lung injury.

Endothelium-dependent relaxation of human pulmonary arteries by ATP has been demonstrated (Greenberg et al., 1987; Dinh Xuan et al., 1990), probably via $P2Y_2$ receptors (Hasséssian and Burnstock, 1995). ATP and UTP, acting via endothelial $P2Y_1$, $P2Y_2$, and/or $P2Y_4$ receptors, elicits vasodilation of bovine and rabbit pulmonary arteries, partly via mobilization of intracellular Ca^{2+} and subsequent prostacyclin release (Lustig et al., 1992; Chen et al., 1996; Qasabian et al., 1997). Shear stress evokes release of ATP from endothelial cells, which causes NO release and pulmonary vasodilation (Hasséssian et al., 1993; Kieffmann et al., 2009). ATP release from endothelial cells induced by NA was greater in the distal (smaller) arteries compared with proximal pulmonary arteries of the rabbit (Takeuchi et al., 1995). In addition to ATP released from endothelial cells, ATP release from erythrocytes contributes to release of NO and subsequent vasodilation (Sprague et al., 1996, 2003; Kotsis and Spence, 2003). Protein kinase C mediates inhibition of endothelial $P2Y_1$ and $P2Y_2$ receptor-mediated phosphoinositide turnover in bovine pulmonary artery (Chen and Lin, 1999). Intercellular communication via Ca^{2+} waves upon mechanical stimulation of calf pulmonary artery endothelial cells is mediated by nucleotides (Moerenhout et al., 2001). Shear stress stimulates human pulmonary artery endothelial cells to release ATP, which then activates Ca^{2+} influx in endothelial cells via $P2X_4$ receptors (Yamamoto et al., 2003). $P2Y_4$ receptor knockout mice do not exhibit normal endothelial cell responses; increased $[Ca^{2+}]_i$, subsequent production of NO, and adaptive vascular remodeling are lost (Ando and Yamamoto, 2009). ATP causes NO-mediated pulmonary vasodilation predominantly via endothelial $P2Y_2$ receptors in newborn rabbits (Konduri et al., 2004).

Uridine adenosine tetraphosphate has been identified as an endothelium-derived vasoconstrictor (Jankowski et al., 2005) that stimulates endothelium-independent contraction of the isolated rat pulmonary artery (Gui et al., 2008). Luminal ATP-induced contraction of rabbit pulmonary arteries has been reported (Baek et al., 2008).

Pulmonary artery rings from fetal sheep show diminished endothelium-mediated relaxation to ADP compared with postnatal sheep (Abman et al., 1991). ATP and adenosine increase pulmonary blood flow in perinatal lamb via $P1$ and $P2$ receptors that are independent of prostacyclin synthesis (Konduri et al., 1992), later identified as A_{2A} and $P2Y_1$ receptors (Konduri et al., 2000), and shown to act via NO (Konduri and Mital, 2000). An increase in ATP release during oxygen exposure may contribute to birth-related pulmonary vasodilation in lambs (Konduri and Mattei, 2002).

Distension of the main pulmonary artery in anesthetized dogs stimulates vagal afferent activity (Moore et al., 2004), possibly via purinergic mechanosensory transduction, where ATP released in response to stretch acts on $P2X_3$ receptors expressed on sensory nerve terminals in the vessel wall (see Burnstock, 2007a).

It has long been known that, unlike in most other vascular beds, hypoxia induces pulmonary vasoconstriction (von Euler and Liljestrand, 1946), leading to pulmonary hypertension. However, hypoxia can also result in pulmonary vasodilation, which, in contrast to the systemic circulation, was shown not to be mediated by adenosine (Gottlieb et al., 1984). The pulmonary hypertensive response to endotoxin was first reported by Hinshaw et al. (1957), and damage to the endothelium may play a role in the development of pulmonary hypertension in humans (Greenberg et al., 1987). ATP-MgCl₂ (the naturally occurring form of ATP in the body is bound to magnesium) has been shown to reduce the vasoconstriction associated with hypoxic pulmonary hypertension (Paidas et al., 1988). On the basis of studies in lambs, it was subsequently proposed that ATP and adenosine may have a beneficial role in the management of pulmonary hypertension in children (Konduri, 1994). Adenosine can decrease pulmonary artery pressure and vascular resistance in patients with primary pulmonary hypertension who respond to calcium-channel blockers (Inbar et al., 1993).

Extracellular nucleotides enhance leukocyte adherence to pulmonary artery endothelial cells via $P2Y$ receptors. Because adherence of leukocytes is an important early step in acute vascular injuries, it was speculated that nucleotide-induced leukocyte adherence could be essential in mediating vascular injuries in such conditions as respiratory distress syndrome and septic shock (Dawicki et al., 1995).

Extracellular ATP is a proangiogenic factor for pulmonary artery vasa vasorum endothelial cells, potentiating the effect of both vascular endothelial growth factor and β -fibroblast growth factor (Gerasimovskaya et al., 2008). Hypoxia, through hypoxia-inducible transcription factor-1 α and -2 α , induces angiogenesis by up-regulating cytokines. The adenosine A_{2A} receptor is an angiogenic target of hypoxia-inducible transcription factor-2 α in human pulmonary endothelial cells and mediates increases in cell proliferation, cell migration, and tube

formation (Ahmad et al., 2009). The authors further show that there is increased expression of A_{2A} receptors in human lung cancer.

Permeability edema is a life-threatening complication accompanying acute lung injury. Extracellular ATP, produced during inflammation, induces a rapid and dose-dependent increase in transendothelial electrical resistance, and restoration of endothelial barrier integrity is achieved via ATP and adenosine acting on endothelial P_{2Y} and A₂ receptors (Lucas et al., 2009).

Dormant pulmonary vein conduction can be provoked by ATP after extensive encircling pulmonary vein isolation, and it has been claimed that radio frequency application for provoked ATP-reconnection may reduce atrial fibrillation recurrence (Hachiya et al., 2007). Adenosine restores atriovenous conduction after ostial isolation of pulmonary veins (Tritto et al., 2004).

III. Trachea

A. Epithelial Cells

Intravenous ATP prevented the decrease in tracheal ciliary beat frequency that occurs in the incised trachea in control anesthetized rats after 70 min (Saano et al., 1990). Luminal administration of ATP increased tracheal ciliary beat frequency, whereas adenosine reduced the ATP effects (Wong and Yeates, 1992). ATP, acting via an epithelial P₂ receptor, stimulated eicosanoid metabolism in rabbit trachea, leading to the production of prostaglandin E₂ (Aksoy et al., 1995). ATP evoked Cl⁻ secretion across tracheal epithelium (Satoh et al., 1995) via multiple P_{2Y} receptors on both apical and basolateral regions (Hwang et al., 1996). ATP caused a rapid increase in both [Ca²⁺]_i and ciliary beat frequency in cultured rabbit tracheal epithelial cells (Korngreen and Priel, 1996; Uzmaner and Priel, 1999). ATP and UTP induced mucin release from primary hamster tracheal surface epithelial cells via P_{2U} receptors (Kim et al., 1996). ATP directly gates a cation-selective channel via P_{2X} receptors in rabbit tracheal ciliated epithelial cells (Korngreen et al., 1998). From P_{2Y₂} receptor knockout studies, it was concluded that the P_{2Y₂} receptor was the dominant P_{2Y} receptor subtype that regulates epithelial Cl⁻ transport in the mouse trachea (Cressman et al., 1998, 1999).

RT-PCR and pharmacological studies identified P_{2Y₁}, P_{2Y₂}, P_{2X₄}, and P_{2X₇} receptors that regulate the increase in [Ca²⁺]_i in freshly isolated rat tracheal epithelial cells (Marino et al., 1999). The P_{2X} receptors on rabbit tracheal ciliated cells were also identified as P_{2X₄} and P_{2X₇} subtypes (Ma et al., 2006). Mechanically induced release of ATP resulted in Ca²⁺ waves in rabbit tracheal epithelial cells (Evans and Sanderson, 1999; Woodruff et al., 1999) and enhanced ciliary beat frequency (Winters et al., 2007). Oscillations in ciliary beat frequency and [Ca²⁺]_i in rabbit tracheal epithelial cells were induced by ATP (Zhang and Sanderson, 2003).

Hypo-osmotic stress stimulated ATP release from tracheal ciliary cells, leading to increase in ciliary beat frequency (Kawakami et al., 2004). Activation of P₂ receptors inhibited ENaC in mouse tracheal epithelial cells via hydrolysis of phosphatidylinositol-bisphosphate (Kunzelmann et al., 2005). P_{2Y₆} receptors increase Ca²⁺ and CFTR-dependent Cl⁻ secretion in mouse trachea (Schreiber and Kunzelmann, 2005). ATP and ADP stimulated goblet cells in the superfused epithelium to secrete mucin; P₂ receptors were present on both apical and basolateral membranes (Davis et al., 1992).

ATP evoked an increase in [Ca²⁺]_i and mucus glycoprotein secretion from tracheal submucosal glands in cats (Shimura et al., 1994), chloride secretion in humans (Yamaya et al., 1996), and [Ca²⁺]_i in pigs (Zhang and Roomans, 1999) via P_{2U} receptors. Both P_{2Y₂} and P_{2Y₄} receptors were characterized in human tracheal gland cells (Merten et al., 1998). It was claimed that a receptor for Ap₄A different from P₂ receptors was present on the human tracheal gland cells (Saleh et al., 1999b).

Adenosine evoked mucus secretion in canine trachea that was blocked by P₁ receptor antagonists (Johnson and McNee, 1985). Mucosal adenosine also stimulated Cl⁻ secretion in canine tracheal epithelium (Pratt et al., 1986). Adenosine induced cAMP-dependent inhibition of ciliary activity in rabbit tracheal epithelium (Tamaoki et al., 1989). Adenosine inhibits endothelin-1 production and secretion in guinea pig tracheal epithelial cells via A_{2B} receptors and cAMP formation (Pelletier et al., 2000). A₃ receptor-mediated potentiation of mucociliary transport, probably through Ca²⁺-mediated stimulation of airway epithelial ciliary motility, has been reported (Taira et al., 2002).

B. Smooth Muscle and Its Innervation

NANC inhibitory neurons supplying the guinea pig, dog, and human trachea were recognized early (Coburn and Tomita, 1973; Coleman and Levy, 1974; Richardson and Bouchard, 1975; Richardson and Béland, 1976; Kalenberg and Satchell, 1979; Russell, 1980), and the inhibitory response to field stimulation was claimed to show features of purinergic neurotransmission (Coburn and Tomita, 1973; Coleman and Levy, 1974).

A₁ and A₃ receptor agonists were shown to inhibit NANC relaxation in isolated guinea pig trachea (Dellabianca et al., 2009). Adenosine deaminase reduced the inhibitory response to adenosine and NANC inhibitory nerve stimulation in guinea pig trachea (Satchell, 1984b). ATP and adenosine caused relaxations of the guinea pig trachea, although adenosine was more potent than ATP (Farmer and Farrar, 1976; Mizrahi et al., 1982; Small, 1982), and Satchell (1984a) suggested that ATP acts on adenosine receptors after breakdown. However, in contradiction, Welford and Anderson (1988) later claimed that ATP itself can cause relaxation without being metabolized to adenosine. Evidence for more than one type of purine receptor was presented for iso-

lated guinea pig trachea, mediating relaxation with a potency order of adenosine > 5'-AMP > ATP > ADP (Jones et al., 1980). Excitatory junctional potentials and contraction, which were reported to be blocked by atropine, were recorded in guinea pig tracheal smooth muscle in response to nerve stimulation, followed by prolonged relaxations that were blocked by indomethacin, suggesting mediation by prostaglandins (Kamikawa and Shimo, 1976). Later, EP₂ prostaglandin receptors were shown to mediate tracheal relaxation to ATP (Fortner et al., 2001). Others reported that neither vasoactive intestinal peptide nor ATP mediated NANC inhibitory neurotransmission in the guinea pig trachea (Karlsson and Persson, 1984). α,β -MeATP contracted guinea pig tracheal smooth muscle, suggesting that a P2X receptor was involved (Candenas et al., 1992), in retrospect probably a P2X1 or P2X3 receptor. ATP and UTP contracted the isolated, perfused trachea; removal of the epithelium decreased it, suggesting that factors released from the respiratory epithelium might be contributing (Fedan et al., 1993a). In a sister article, the authors showed that concentrations of ATP higher than those causing contraction of the perfused trachea relaxed the smooth muscle via a P2, but not P1, receptor (Fedan et al., 1993b). Propofol attenuated ATP-induced contraction of rat trachea through inhibition of P2 receptor binding (Yamaguchi et al., 2007). It was claimed in a more recent article that ATP caused contractions and increased $[Ca^{2+}]_i$ in single smooth muscle cells from porcine trachea via P2X4 receptors (Nagaoka et al., 2009). An ecto-ATP-diphosphohydrolase similar to that described in lung was shown to be localized in the membranes of nonvascular smooth muscle cells of the bovine trachea (Picher et al., 1994).

Adenosine caused contraction of spiral strips and relaxation of transverse strips of guinea pig trachea (Satchell and Smith, 1984). A₂ receptors were identified in tracheal smooth muscle by pharmacological analysis (Brackett et al., 1990; Losinski and Alexander, 1995). Ghai et al. (1987) presented evidence for both A₁ and A₂ receptors on guinea pig tracheal smooth muscle. Adenosine relaxed guinea pig trachea via an A_{2B} receptor (Brown and Collis, 1982; Karlsson and Persson, 1984). Relaxation of the trachea is also produced by xanthines, but this is independent of adenosine antagonism (Darmani and Broadley, 1986; Small et al., 1988). Advenier et al. (1988) suggested that epithelium had an influence on the mechanical response of guinea pig trachea to adenosine, but a later article (Thorne and Broadley, 2001) did not support this possibility. However, treatment with caffeine for 10 weeks led to changes in adenosine-induced relaxation that did seem to involve the epithelium (Brugós et al., 2007). Adenosine induced a cholinergic tracheal reflex contraction in guinea pig in vivo via an A₁ receptor-dependent mechanism (Reynolds et al., 2008). A₂ receptor deficiency led to impaired tra-

cheal relaxation via the NADPH oxidase pathway in allergic mice (Nadeem et al., 2009).

IV. Nasal Mucosa

Both basolateral and apical application of ATP induces Cl⁻ secretion in human nasal epithelium (Clarke and Boucher, 1992). Extracellular ATP increases fluid transport by human nasal epithelial cells in culture (Benali et al., 1994). Evidence was presented to suggest that ATP acts as a cotransmitter with NA in sympathetic nervous control of blood vessels of the nasal mucosa of cats and dogs (Lacroix et al., 1994). UDP activates a receptor, distinct from the P2Y₂ receptor, on human nasal epithelial cells (Lazarowski et al., 1997), which in retrospect was a P2Y₆ receptor. Extracellular sodium regulates ciliary motility in epithelial cells isolated from surgically excised nasal polyps by inhibiting P2X receptor-mediated responses (Ma et al., 1999). ATP, UTP, UDP, and adenosine-stimulated ciliary activity of human nasal epithelial cells via P2Y₂, P2Y₆, and A_{2B} receptors, whereas P2Y₁, P2Y₄, and P2Y₁₁ receptors were not involved (Morse et al., 2001). ATPase expression in nasal epithelial cells showed correlation with ciliary activity (Schütz et al., 2002). UTP and ATP γ S were shown to induce mucin secretion via Ca²⁺-dependent pathways in human nasal epithelial cells (Choi et al., 2003). Purinergic stimulation via P2 receptors induces Ca²⁺-dependent activation of Na⁺-K⁺-2Cl⁻ cotransporter in human nasal epithelium (Shin et al., 2004). RT-PCR studies of human nasal epithelial cells showed expression of mRNA for P2X3, P2X4, P2X7, P2Y₁, P2Y₂, P2Y₄, P2Y₆, P2Y₁₁, and P2Y₁₂, and functional studies showed that luminal membranes express P2Y₂, P2Y₆, and probably P2Y₁₁ receptors, whereas basolateral membranes express P2Y₂ receptors; P2X3, P2X4, P2X7, P2Y₄, and P2Y₁₂ did not seem to be functional in normal epithelial cells (Kim et al., 2004). Immunohistochemical studies of P2 receptor proteins showed P2X3 receptors on neurons of the rat nasal mucosa, P2X5 and P2X7 receptors on respiratory epithelial cells, P2Y₁ receptors on respiratory epithelial cells and submucosal glands, and P2Y₂ receptors on mucous secretory cells (Gayle and Burnstock, 2005). Zinc increased ciliary beat frequency in a calcium-dependent way in mouse nasal septal epithelial cultures through activation of P2X receptors and may thus be a useful agent for stimulating mucociliary clearance (Woodworth et al., 2010).

V. Central Control of Respiration

The ventrolateral medulla contains a network of respiratory neurons that are responsible for the generation and shaping of respiratory rhythm; it also functions as a chemoreceptive area mediating the ventilatory response to hypercapnia. Evidence has been presented that ATP acting on P2X2 receptors expressed in ventrolateral medulla neurons influences these functions

(Gourine et al., 2003). A potentially important role for P2 receptor synaptic signaling in respiratory motor control is suggested by the multiple physiological effects of ATP in hypoglossal activity associated with the presence of P2X2, P2X4, and P2X6 receptor mRNA in the nucleus ambiguus and the hypoglossal nucleus (Collo et al., 1996; Funk et al., 1997), and microinjection of ATP into the caudal nucleus tractus solitarius (NTS) in the working heart-brainstem preparation of rats produces respiratory responses (Antunes et al., 2005). P2X receptors have been shown to be involved in the histamine-induced enhancement of the cough reflex severity in guinea pigs by mediating direct activation of rapidly adapting receptors, whereas P2Y receptors mediate increase in histamine release, indirectly increasing the cough reflex sensitivity (Kamei and Takahashi, 2006).

Evidence showing P2X receptor subtypes (P2X1, P2X2, P2X5, and P2X6) within areas of the ventrolateral medulla in the brainstem has been presented, and it was shown that activation of these receptors was implicated in respiratory networks including the Bötzing and pre-Bötzing complex (PBC) (Thomas et al., 2001). Recent data indicated that ionotropic glutamate and P2X receptors in the rostral ventrolateral medulla of the Bötzing complex play roles in opposite directions in the control of respiratory responses to chemoreflex activation in awake rats (Moreaes et al., 2011). It has been shown in rodents that the ATP-adenosine balance is a determining factor in the purinergic modulation of the PBC inspiratory rhythm and that the underlying pathways seem to differ between mice and rats (Zwicker et al., 2011). Another article showed that the chemosensitive regions of the ventrolateral medulla were immunoreactive for the P2X2 receptor, which mediates ATP potentiation of respiratory discharge frequency (Lorier et al., 2004). P2X receptors mediate central inspiratory facilitation by low-frequency vagal afferent inputs in the rabbit (Takano et al., 2004). Microinjection of ATP into different regions of the NTS produces different respiratory responses. ATP injection into the intermediate NTS produced apnea (Bianchi et al., 1995); injected into the caudal NTS, it caused excitation of neurons involved in ventilatory pathways activated by the chemoreflex (Antunes et al., 2005). Increase of pCO₂ in the arterial blood triggers the release of ATP from three chemosensitive regions on the ventral surface of the medulla oblongata, and blockade of ATP receptors at these sites diminishes the chemosensory control of breathing (Gourine et al., 2005). Occupation of P2Y₁ receptors in the PBC, a key site of inspiratory rhythm generation, led to increase in frequency (Lorier et al., 2007). In a developmental study, ATP applied locally into PBC slices potently increased activity as early as embryonic day 19, the youngest rats tested (Huxtable et al., 2007). Local application of the P2Y₁ agonists 2-methylthio-ADP and (*N*)-methanocarba-2-methylthio-ADP (MRS2365) to neurons in the XII nucleus evoked a tonic discharge, although UTP also

had an effect (de Souza Alvares et al., 2007), perhaps indicating involvement also of P2Y₂ receptors. Further experiments led to the conclusion that ATP is a potent excitatory modulator of the inspiratory network with a dynamic interaction between the actions of ATP at P2 receptors and ectonucleotidases that degrade it to metabolites that act on P2Y and P1 receptors (Huxtable et al., 2009). Connexin hemichannel-mediated CO₂-dependent release of ATP in the medulla oblongata contributes to central respiratory chemosensitivity (Huckstepp et al., 2010). Astrocytes in the brainstem chemoreceptor area are highly chemosensitive; they respond to decreases in pH with increases in [Ca²⁺]_i and release of ATP, which led to activation of chemosensitive neurons and induced adaptive increases in breathing (Gourine et al., 2010). It was demonstrated that PBC glia also respond to ATP, with increased [Ca²⁺]_i and glutamate release, and may therefore contribute to the ATP sensitivity of PBC networks and possibly to the hypoxic ventilatory response (Huxtable et al., 2010).

Adenosine mechanisms were originally implicated in the central nervous regulation of breathing in several species (Hedner et al., 1982; Mueller et al., 1984; Wessberg et al., 1984; Eldridge and Millhorn, 1987; Norsted et al., 1987; Bissonette et al., 1991; Barros and Branco, 2000). Adenosine injected into the caudal nucleus of the NTS decreased respiratory frequency by an action on A_{2A} receptors expressed on glutamatergic nerve terminals (Phillis et al., 1997). A medial thalamic sector was shown to be involved in adenosine-induced apnea (Kos et al., 2000). Intracerebroventricular administration of P2 receptor antagonists reduced the minimum alveolar concentration of inhaled volatile anesthetics (Masaki et al., 2000).

In a study of perinatal development in rats, it was concluded that respiration is already modulated by adenosine acting on A₁ receptors in the medulla oblongata in the period around birth (Herlenius et al., 2002). A_{2A} receptors interact with GABAergic pathways in the medulla oblongata to modulate respiration in neonatal piglets (Wilson et al., 2004), whereas A₁ receptors were reported to modulate the activities of both inspiratory and expiratory neurons in the medial region of the nucleus retrofacialis of neonatal rats (Wang et al., 2005). Activation of central A_{2A} receptors enhances superior laryngeal nerve stimulation-induced apnea in piglets via a GABAergic pathway (Abu-Shaweesh, 2007). Adenosine release from NTS does not seem to mediate hypoxia-induced respiratory depression in rats (Gourine et al., 2002). The reduced hypoxic ventilatory response observed in obese Zucker rats was attributed to depressed adenosinergic peripheral excitatory mechanisms and to enhanced central depression mechanisms (Lee et al., 2005).

Reviews concerned with purinergic signaling in the brain stem associated with airways functions are available (Spyer and Thomas, 2000; Gourine and Spyer,

2003; Gourine et al., 2004; Gourine, 2005; Spyer and Gourine, 2009; Spyer, 2009).

VI. Diseases of the Airways

Inflammation is a feature of most diseases of the airways, including asthma, COPD, CF, dyspnea, allergy, infection and injury. Mucous hyperproduction in the airways is also characteristic of all pulmonary obstructive diseases, including chronic bronchitis, asthma, COPD, CF and primary ciliary dyskinesia.

The release of purine nucleotides from airway epithelial cells is often dramatically increased during inflammatory processes and is believed to play an important role in the pathophysiology of chronic lung disease (see Adriaensen and Timmermans, 2004). Measurement of adenosine, ATP, and AMP in sputum or exhaled breath condensate has been used as a noninvasive method to track airway inflammation (Huszár et al., 2002; Esther et al., 2008). Extracellular ATP was reported to sensitize airway smooth muscle cells to contractile agonists (e.g., metacholine) via P2X receptor activation, without enhancing the induced $[Ca^{2+}]_i$ rise, and the resulting hyper-responsiveness was referred to as “ Ca^{2+} sensitization” (Oguma et al., 2007). Erythromycin therapy has been used to decrease airway secretion in chronic inflammatory airway diseases (Kondo et al., 1998), perhaps by selectively inhibiting Ca^{2+} influx through P2X receptor channels (Zhao et al., 2000). Uridine and 4-thiouridine have anti-inflammatory effects, reducing edema formation, leukocyte infiltration, and tumor necrosis factor in a lung inflammation model (Evaldsson et al., 2009). Alveolar macrophages (also see section II.B.1) play a pivotal role in the development of chronic lung inflammatory reactions such as idiopathic pulmonary fibrosis, silicosis, asbestosis, hypersensitivity pneumonitis, sarcoidosis, and mycobacterium tuberculosis. P2X7 receptors are expressed in alveolar macrophages, which upon stimulation activate the proinflammatory IL-1 to IL-5 cytokine cascade and the formation of multinucleated giant cells, a hallmark of granulomatous reactions (Lemaire and Leduc, 2004). P2X7 receptors may be a relevant target for therapeutic intervention in lung hypersensitivity reactions associated with chronic inflammatory responses. ATP also promotes tumor necrosis factor- α -elicited IL-8 expression via P2X ion channel-triggered Ca^{2+} entry (Théâtre et al., 2009). Adenosine promotes IL-6 release from airway epithelium during inflammation (Sun et al., 2008). There is a role for the vagal parasympathetic reflex bronchoconstriction in dyspnea, which seems to be exaggerated in inflammatory airway disease (Udem and Nassenstein, 2009). Paruchuri et al. (2009) reported that leukotrienes E_4 -induced pulmonary inflammation is mediated by the P2Y₁₂ receptor.

Anti-inflammatory effects of adenosine have been described previously (Schrier et al., 1990; Vass and Horváth, 2008). Adenosine, probably via A₁ receptors, sig-

nificantly elevated the sensitivity of C-fiber afferents in rat lung (Gu et al., 2003). Interaction between adenosine and IL-13 evoked a stimulation that may contribute to the nature and severity of airway inflammation and fibrosis (Blackburn et al., 2003). Mast cell involvement in adenosine-mediated airway hyper-reactivity was reported in a murine model of ovalbumin-induced lung inflammation (Wyss et al., 2005). A₃ receptor signaling is needed for migration of eosinophils into the airways, and influences airway inflammation, mucus production, and fibrosis (Young et al., 2004; Morschl et al., 2008). A_{2B} receptors are reportedly involved in airway inflammation via induction of IL-19 release from human bronchial epithelial cells, resulting in tumor necrosis factor- α release from monocytes, which in turn up-regulates A_{2B} receptor expression (Zhong et al., 2006). An A_{2B} receptor antagonist has been proposed as a clinical candidate for chronic inflammatory airway disease (Elzein et al., 2008). A protective role for A_{2B} receptor signaling during the acute stages of lung inflammation has been proposed (Zhou et al., 2009b). CD39 and CD73 ectonucleotidases are critical mediators of pulmonary neutrophil transmigration in LPS-induced lung inflammation (Reutershan et al., 2009).

A. Asthma

Asthma is a chronic inflammatory disease of the airways, characterized by recurring bronchospasm, expressed by wheezing, coughing, chest tightness, and shortage of breath.

The involvement of adenosine in asthma has been the subject of considerable attention over the years, after it was shown that it produced powerful bronchoconstriction in asthma, although it had little effect in healthy lungs; xanthines, antagonists of adenosine receptors, were widely used for treatment of asthma (for reviews, see Persson, 1982; Kawasaki et al., 1983; Holgate et al., 1987, 1990; Feoktistov and Biaggioni, 1996; Fozard and Hannon, 2000; Meade et al., 2001; Fozard and McCarthy, 2002; Holgate, 2002, 2005; Polosa et al., 2002; Fozard, 2003; Lee et al., 2003; Livingston et al., 2004; Caruso et al., 2006, 2009; Russo et al., 2006; Brown et al., 2008b; Wilson, 2008). It was suggested that the hyper-responsive effects of adenosine in isolated bronchi was due to indirect actions by liberation of histamine and leukotrienes (Björck et al., 1992).

Although theophylline and caffeine are well established antagonists to adenosine receptors, their bronchodilator action in patients with asthma does not seem to be due to this mechanism of action; for example, enprofylline is 4 to 5 times more potent than theophylline although it was reported to be devoid of adenosine receptor antagonism (Persson et al., 1981). Adenosine, acting through A₂ receptors, was reported to both stimulate and inhibit histamine release from mast cells in the lung (Hughes et al., 1984; Ott et al., 1992). Inhaled adenosine, but not guanosine, is a potent bronchocon-

strictor in asthma (Cushley et al., 1983). Antagonism of adenosine-induced bronchoconstriction in asthma by oral theophylline was described previously (Mann and Holgate, 1985). Data were presented to suggest that adenosine-induced bronchoconstriction was not mediated by parasympathetic or sympathetic reflexes (Mann et al., 1985). Inhalation of dipyridamole, which inhibits adenosine uptake, increased adenosine-induced bronchospasm in patients with asthma (Crimi et al., 1988). Patients with asthma have neutrophils that exhibit diminished responsiveness to adenosine and it was suggested that this may be responsible for the severity and persistence of inflammation in the airways of patients with asthma (Sustiel et al., 1989). Superoxide generation and its modulation by adenosine in neutrophils from patients with asthma was noted (Meltzer et al., 1989).

There is elevated expression of A_1 receptors in bronchial biopsy specimens from patients with asthma (Brown et al., 2008a). Administration of an aerosolized phosphorothioate antisense oligodeoxynucleotide targeting the A_1 receptor desensitized the animals for subsequent challenge with either adenosine or dust-mite allergen, suggesting that adenosine is an important mediator of both airway inflammation and obstruction (Nyce and Metzger, 1997). Lowering adenosine levels with adenosine deaminase therapy had striking effects on gene expression analyzed with cDNA arrays in inflammatory airway disease associated with elevated adenosine (Banerjee et al., 2002). It has been claimed that airway bronchoconstrictor responses to aerosolized adenosine in mice occur largely through A_3 receptor activation and that mast cells contribute significantly to these responses (Tilley et al., 2003). The anti-inflammatory properties of theophylline in asthma include an inhibition of circulating progenitor cells (Wang et al., 2003). Adenosine cooperates with inflammatory cytokines to stimulate mucin production in the asthmatic airway (McNamara et al., 2004). Intravenous adenosine in healthy subjects does not induce bronchospasm, but causes dyspnea, most likely by an effect on vagal C fibers in the lungs (Burki et al., 2005). A more recent article claimed that all four P1 receptor subtypes play various roles in the airway responses to inhaled 5'-AMP in sensitized guinea pigs (Smith and Broadley, 2008). Genetic polymorphisms of A_1 and A_{2A} receptors were associated with aspirin-intolerant asthma (Kim et al., 2009).

In summary, several widely different mechanisms have been proposed to account for the potent bronchoconstriction produced by adenosine in patients with asthma (but not healthy patients):

1. Inhibition of release of histamine from lung mast cells via A_{2B} receptors (Forsythe and Ennis, 1999) and/or A_3 receptors (Hua et al., 2008). An increase in mast cell degranulation was reported in adenosine deaminase-deficient mice (Zhong et al., 2001).

Adenosine also potentiates the immediate response to allergen in patients with asthma.

2. Enhancement of the potent bronchoconstrictor effects of endothelin 1 released from epithelial cells (Kanazawa et al., 1997).
3. Proinflammatory effects of adenosine in patients with asthma (see Rorke et al., 2002; Caruso et al., 2009).
4. A_1 receptors are up-regulated in bronchial smooth muscle and epithelial cells (Brown et al., 2008a).
5. Adenosine production is increased by hypoxia, inflammatory cell activation, and allergen challenge and is elevated in patients with asthma. It is also increased in exercise-induced bronchoconstriction in asthma (Vizi et al., 2002).
6. Increased incidence of adenosine-induced dyspnea in asthma may be due to adenosine sensitization of vagal C fibers (Burki et al., 2006) via A_1 receptors (Hua et al., 2007).

However, which of the above or other mechanisms are involved still needs to be resolved.

Although the early emphasis was on the roles of adenosine in asthma, there have been an increasing number of studies concerned with the roles of nucleotides in this disease. For example, data were presented to suggest that ATP plays an important modulatory role in histamine release from human lung mast cells, and it was suggested that it may be involved in allergic/asthma reactions (Schulman et al., 1999). ATP was shown to be a more potent bronchoconstrictor and to have greater effects on dyspnea and other symptoms than AMP in patients with asthma (Basoglu et al., 2005). P2X7 receptor function is altered in asthma; P2X7 pore activity is associated with change in nasal lavage neutrophil counts during the development of asthma symptoms (Denlinger et al., 2009).

In allergic patients with asthma, inflammation is triggered by specific inhalation of allergens, such as house dust-mite allergen and pollen spores or nonspecific triggers such as air pollution and viral infection. Activation of dendritic cells is essential for the immune response to allergens. In allergic patients with asthma, inhaled adenosine-induced bronchoconstriction was claimed to be mediated by A_1 receptors (Marmo et al., 1985). This was also demonstrated in allergic rabbits; a role for A_3 receptors was not found (el-Hashim et al., 1996; Abebe and Mustafa, 1998). Challenge of sensitized Brown Norway rats with ovalbumin induced a marked airway hyperresponsiveness to adenosine, which, in contrast to several other findings (mentioned previously), was believed to be largely due to an increased release of histamine and 5-HT from mast cells (Hannon et al., 2001). Theophylline attenuated adenosine-enhanced airway inflammation but could not reverse allergen-induced airway inflammation (Fan and Mustafa, 2002). A selective A_{2A} receptor agonist, 2-[p-(2-carboxyethyl)phenethylamino]-

5'-*N*-ethylcarboxamidoadenosine (CGS 21680), inhibited allergic airway inflammation in the rat (Fozard et al., 2002). In contrast, A_{2B} and A₃ receptors were reported to play an important role in adenosine-induced bronchoconstriction in an allergic mouse model (Fan et al., 2003). In other mouse allergy models, A_{2B} receptors were identified as the main receptor subtype involved in increases in ion transport (Kornerup et al., 2005; Mustafa et al., 2007). Mast cells were shown to be involved in adenosine-mediated bronchoconstriction and inflammation in an allergic mouse model (Oldenburg and Mustafa, 2005). Mast cells were claimed to be activated after adenosine challenge in allergic, but not nonallergic, patients with asthma (Bochenek et al., 2008). AMP and an A₁ receptor-selective agonist induced airway obstruction in sensitized guinea pigs by a mechanism unrelated to release of histamine from mast cells but probably through stimulation of sensory nerves and reflex activity (Keir et al., 2006). A₃ receptor signaling contributed to airway mucin secretion after allergen challenge in mice (Young et al., 2006). Allergen inhalation decreased adenosine receptor expression in sputum and blood of patients with asthma (Versluis et al., 2008). A_{2A} receptor deficiency led to impaired tracheal relaxation in allergic mice (Nadeem et al., 2009). Allergen-induced airway hyper-responsiveness was absent in ecto-5'-nucleotidase (CD73)-deficient mice (Schreiber et al., 2008).

Allergen challenge caused short-term accumulation of ATP in the airways of subjects with asthma and mice with experimentally induced asthma; when ATP levels were neutralized using apyrase or, in mice, treated with P2 receptor antagonists, the features of asthma were reduced, including eosinophilic inflammation, Th2 cytokine production and bronchial hyper-reactivity (Idzko et al., 2007). The authors concluded that the adjuvant effects of ATP were due to the recruitment and activation of lung myeloid dendritic cells that induced Th2 responses. ATP contributes to the induction of inflammation in allergic asthma by activation and recruitment of various inflammatory cells (Willart and Lambrecht, 2009). Evidence has been presented that P2Y₁ receptors are involved in allergic airway inflammation, which could be at least partly explained by its role in the regulation of dendritic cell maturation (Maitre et al., 2008). Extracellular adenine nucleotides inhibit the release of major monocyte recruiters by human monocyte-derived dendritic cells and pharmacological data support the involvement of P2Y₁₁ and P2Y₁ purinergic receptors (Horckmans et al., 2006). In a recent article, it is proposed that purinergic signaling via P2Y₂ receptors in the respiratory epithelium is a critical sensor for airway exposure to airborne allergens, which may provide novel opportunities to dampen hypersensitivity in the Th2-type immune responses in asthma (Kouzaki et al., 2011). In a study from a different group, it was also claimed that ATP acts as a "danger signal" by inducing inflammation via P2Y₂ receptors, via recruitment of

neutrophils to the lungs; P2Y₂ receptor-deficient mice showed reduced pulmonary inflammation and development of emphysema after short-term smoke exposure (Cicko et al., 2010). It has also been demonstrated that P2X7 receptors are implicated in the pathophysiology of allergy-induced lung inflammation (Faffe et al., 2009). For example, low-intensity aerobic exercise attenuates the epithelial response in allergic asthma in a mouse model, reducing P2X7 receptor expression, synthesis of Th2 cytokines, chemokines, adhesion molecules, and proteases that regulate inflammation, remodeling, and hyper-responsiveness (Vieira et al., 2011). It has also been proposed that targeting P2X7 receptors on hematopoietic cells, such as dendritic cells or eosinophils, might be a novel therapeutic option for the treatment of allergic asthma (Müller et al., 2011).

B. Chronic Obstructive Pulmonary Disease

COPD refers to chronic bronchitis and emphysema, diseases in which the airways become narrowed, causing shortness of breath. In contrast to asthma, the limited airflow is poorly reversible and usually gets progressively worse over time. COPD is caused by noxious particles or gas, most usually from tobacco smoking, which triggers an inflammatory response in the lungs. The inflammatory response in the larger airways is known as chronic bronchitis, whereas inflammation in the alveoli is known as emphysema.

As for asthma, the main purinergic emphasis has been on the role of adenosine receptors in COPD (Polosa et al., 2002; Polosa, 2002; see reviews by Peleg and Schulman, 2002; Blackburn, 2003; van den Berge et al., 2004, 2007; Caruso et al., 2006; Mohsenin and Blackburn, 2006; Trevethick et al., 2008; Polosa and Blackburn, 2009; Zhou et al., 2009b, 2010). Enprofylline, a xanthine lacking adenosine receptor antagonism, was shown to be a potent bronchodilator in patients with COPD (Lunell et al., 1982). A_{2B} receptors on human mast cells are a strategic target for COPD (Spicuzza et al., 2003). The affinity of A₁, A_{2A}, and A₃ receptors expressed in different cells in peripheral lung parenchyma was significantly decreased in patients with COPD, whereas their density was increased; the affinity of A_{2B} receptors was not altered, but their density was significantly decreased (Varani et al., 2006). There were no significant differences in mast cell numbers between patients with COPD or asthma and age-matched healthy control subjects (Liesker et al., 2007). Activation of A_{2A} receptors inhibited inflammatory cell activation in COPD (Bonneau et al., 2006). Inhaled A_{2A} receptor agonists have been used for the treatment of COPD (Mantell et al., 2008; Trevethick et al., 2008; Thomas et al., 2008). It has been shown that after a 1-year smoking cessation airway inflammation in COPD persisted or was aggravated and that this may be due to an increase in neutrophils expressing A₃ receptors and macrophages expressing A₁ receptors (Versluis et al., 2009). Dyspnea, a side effect of

adenosine infusion, is not correlated with impaired respiratory resistance in patients with COPD (Fricke et al., 2008). It has been reported that oxidative/nitrosative stress selectively altered A_{2B} receptors in COPD (Varani et al., 2010). Mice lacking the A_{2B} receptor revealed enhanced airway inflammation and remodeling (Zhou et al., 2009a).

ATP seems to be implicated in COPD and asthma as well as adenosine (Adriaensen and Timmermans, 2004; Mortaz et al., 2010). However, ATP in exhaled breath condensate was reported to be the same for healthy people and patients with COPD (Lázár et al., 2008). ATP-induced pulmonary vasodilation has been described in patients with COPD (Gaba et al., 1986; Gaba and Préfaut, 1990). Activation of purinergic signaling by cigarette smoke seems to be involved in the pathogenesis of emphysema (see Fig. 10); cigarette smoke induces ATP release from neutrophils, which plays a crucial role in release of the chemokine CXCL8 that is largely responsible for neutrophil recruitment into the sites of inflammation, and of elastase that leads to the destruction of lung tissue (Mortaz et al., 2009). Mortaz et al. (2010) later suggest that P2X7 receptors might be involved. In murine model experiments focused on emphysema, changes in lung innate immune cell recruitment were associated with both P2Y₂ and P2X7 receptors (Cicko et al., 2010; Lucattelli et al., 2011).

In summary, adenosine-induced bronchoconstriction in both COPD and asthma involves mast cell degranu-

lation and neural pathways, whereas the direct effects on smooth muscle are less clear. Production of inflammatory cytokines via A_{2B} receptors is also involved. A_{2A} agonists and A_1 , A_{2B} , and A_3 antagonists have been used for the treatment of COPD.

C. Airway Infections

Erythromycin is an antibiotic used widely for the treatment of upper and lower respiratory tract infections. One of the most conspicuous effects of erythromycin is the suppression of fluid secretion from bronchial epithelial cells in the treatment of bronchitis. Erythromycin has been shown to block the P2X receptor-mediated Ca^{2+} influx and may represent one mechanism by which it exerts its antisecretory effects in the treatment of chronic respiratory tract infections (Zhao et al., 2000). During bacterial infections of the lung, the airway epithelial cells play a central role in activating the innate immune response (i.e., release of proinflammatory cytokines and subsequent recruitment of neutrophils) to fight the infection. *Pseudomonas aeruginosa* produces a common bacterial pathogen that causes chronic infections in immunocompromised hosts, such as patients with CF. In mouse airways, luminal exposure to the protein flagellin, a structural component of this pathogen, led to inhibition of Na^+ absorption by the ENaC, probably because of the release of ATP and occupation of P2Y receptors, which activate the mitogen-activated protein kinase pathway leading to inhibition of ENaC




FIG. 10. Diagrammatic representation of the possible mechanism of action of ATP in pathogenesis of lung emphysema. MMP, matrix metalloproteinase; NALP3, Nacht, leucine-rich repeat and PYD domains-containing protein 3; NF- κ B, nuclear factor- κ B; TLR, Toll-like receptor; TNF, tumor necrosis factor. [Reproduced from Mortaz E, Folkerts G, Nijkamp FP, and Henricks PA (2010) ATP and the pathogenesis of COPD. *Eur J Pharmacol* 638:1–4. Copyright © 2010 Elsevier. Used with permission.]

(Kunzelmann et al., 2006). In another article, bacterial flagellin was shown to interact with Toll-like receptors, and the cell surface glycolipid, asialoGM1, was shown to activate an innate immune response. The release of ATP induced by flagellin was dependent on a Toll signaling cascade, revealing a role for autocrine extracellular ATP in Toll-like receptor signaling (McNamara et al., 2006). ATP-induced increases in $[Ca^{2+}]_i$ in airway epithelial cells synergized with *P. aeruginosa* flagellin in activating the immune response (Fu et al., 2007). Extracellular nucleotides regulate expression and release of the antimicrobial chemokine CCL20 in primary human airway epithelial cells; P2Y₂ and P2Y₆ agonists increased CCL20 release (Marcet et al., 2007b). It was concluded that nucleotides modulate chemokine release and leukocyte recruitment in inflamed airways by acting on P2Y receptors on both epithelial and immune cells.

Mucociliary clearance constitutes the first line of defense against airways infection. Inhaled pathogens are trapped by a mucus layer positioned above ciliated epithelia. In the airways, adenine nucleotides support a complex network mediating host defenses. ATP released by the epithelium into the airway surface liquid layer regulates mucus clearance via P2 receptors and, after breakdown to adenosine, via P1 receptors (Zuo et al., 2008). ATP-induced activation of the NADPH oxidase homolog dual oxidase 1 mediated airway epithelial responses to bacterial stimuli (Boots et al., 2009). They showed further that epithelial IL-8 production in response to bacterial stimuli involved ATP release and dual oxidase 1 activation. Patients with pulmonary tuberculosis had significantly higher adenosine deaminase activity in bronchoalveolar lavage fluid (Orphanidou et al., 1998) and in the sputum (Dimakou et al., 2009). ATP is a modulator of lung inflammation in pneumonia in cattle; it is released by activated macrophages and damaged cells (McClenahan et al., 2009). It was shown in this study that ATP caused striking morphological and permeability changes in bovine pulmonary epithelial cells and that both epithelial cells and pulmonary microvascular endothelial cells expressed mRNA for P2X7 receptors, with a P2X7 receptor antagonist reversing the changes in permeability induced by ATP.

The pneumovirus respiratory syncytial virus is the most common cause of lower respiratory tract disease in infants and children. It has a detrimental inhibitory effect on alveolar clearance, an effect that appears to be mediated by UTP, perhaps released by the bronchoalveolar epithelium in response to infection (Davis et al., 2004). This suggests that P2Y₂ receptor antagonists may be useful for the treatment of severe respiratory syncytial virus bronchiolitis. Respiratory syncytial virus infection causes fluid accumulation and decreased Na⁺ transfer across respiratory epithelial cells, probably via UTP released from epithelial cells (Chen et al., 2009). In addition, influenza A virus inhibits alveolar fluid clearance in BALB/C mice (Wolk et al., 2008).

D. Lung Injury

Acute lung injury and acute respiratory stress syndrome (ARDS) are major causes of respiratory failure, characterized by edema, neutrophil infiltration with hemorrhage, and increased production of inflammatory cytokines. In rats treated with α -naphthylthiourea to induce acute lung injury, particularly to the endothelium, as demonstrated by pulmonary edema, the ectoenzymes catalyzing ADP and AMP hydrolysis were little effected (Grantham and Bakhle, 1988). Sublethal oxidant injury inhibited ATP-induced surfactant secretion (Warburton et al., 1989b). ARDS can be the result of trauma (as well as smoke inhalation, aspiration, and sepsis). The need to support the failing lung with mechanical ventilation is potentially lifesaving; unfortunately, however, alveolar overdistension and pulmonary shear stress may itself cause lung injury (ventilator-induced lung injury), increasing bronchoalveolar leakage leading to lung edema. It has been suggested that ventilator-induced lung injury may involve stretch-associated release of ATP from epithelial cells or pulmonary neuroendocrine cells in NEBs (Brouns et al., 2000, 2003a,b; Rich et al., 2003), which may therefore be a therapeutic target for this condition. ATP-dependent calcium signaling during ventilator-induced lung injury is amplified by hypercapnia (Briva et al., 2011). The protective effect of ATP-MgCl₂ in ischemia-reperfusion lung injury seems to require the presence of leukocytes (Chen et al., 2003).

Pulmonary fibrosis can be caused by injury, although it may also be secondary to other lung disorders or may even be idiopathic. Lung injury induced by bleomycin, an intercalating agent that causes DNA strand breaks, causes lung fibrosis, and it was reported that suramin, a P2 receptor antagonist, did not affect transforming growth factor β activity, which is presumed to play a role in pulmonary fibrosis (Lossos et al., 2000). However, it has been proposed that ATP, released from airways after administration of bleomycin, constitutes a major endogenous danger signal that engages the P2X7 receptor pannexin-1 axis, leading to IL-1 β maturation and lung fibrosis (Riteau et al., 2010). In patients with idiopathic pulmonary fibrosis, alterations in adenosine metabolism were described and this study also presented evidence that A_{2B} receptor signaling could promote the production of inflammatory and fibrotic mediators in patients with these disorders (Zhou et al., 2010). Chronic adenosine elevations are associated with pulmonary fibrosis in adenosine deaminase-deficient mice, suggesting that adenosine functions as a profibrotic signal in the lung (Chunn et al., 2005).

Lung epithelial cells release ATP during ozone exposure, which protects against ozone toxicity by inhibiting apoptotic and necrotic cell death; P2 receptor antagonists diminished this protective action (Ahmad et al., 2005). Lung injury produced by the gaseous contami-

nant nitrogen dioxide resulted in a loss of ATP and a reduction in cell viability (Bakand et al., 2006). The cytotoxic effects of gaseous airborne contaminants NO₂, SO₂, and NH₃ were compared, and ATP levels were shown to be reduced with all three gases (Bakand et al., 2007). ATP-mediated activation of dual oxidase 1, an NADPH oxidase homolog within the tracheobronchial epithelium, results in epithelial cell migration and wound repair after injury by environmental pollutants (Wesley et al., 2007). Inhalation of industrial environmental arsenic has long been known to lead to respiratory compromise, including lung cancer. In a recent study, it was shown that short-term exposure to low concentrations of sodium arsenide reduced the ability of lung epithelial cells to respond to wound-induced purinergic signaling via both P2X and P2Y receptors (Sherwood et al., 2011). The P2 receptor agonist ATP γ S had a protective effect against acute lung injury induced in a murine model by the bacterial endotoxin LPS (Kolosova et al., 2008). ATP released during mechanical ventilation caused lung injury via P2Y receptors (Matsuyama et al., 2008). LPS and mechanical ventilation modify purinoceptor expression in the lung. LPS caused expression of A₁, A_{2A}, and A_{2B} and P2Y receptors to be increased, and A₃ receptors to be decreased, whereas high-volume ventilation reduced P2Y₄ mRNA levels (Riesenman et al., 2008). On the basis of a study of LPS administration in isolated and ventilated rabbit lungs, it was suggested that adenosine A₂ receptor-mediated stimulation might be beneficial during acute lung injury (Heller et al., 2007). Both A_{2A} receptors (Reutershan et al., 2007) and A_{2B} receptors (Eckle et al., 2008) have been claimed to be involved in the attenuation of acute lung injury.

Adenosine prevented phorbol myristate acetate-induced canine lung injury via an A₂ receptor mechanism (Adkins et al., 1993). A_{2A} receptors mediated adenosine-stimulated wound healing of bronchial epithelial cells (Allen-Gipson et al., 2006). A_{2A} receptor activation reduced lung injury in trauma/hemorrhagic shock (Haskó et al., 2006). Purinergic sensory transduction pathways contribute to activation of the nasal sensory irritation response to inspired irritant vapors are present in the mouse; the involved nerve terminals may serve as sensors of airway cell damage (Vaughan et al., 2006). Patients with ARDS and high altitude pulmonary edema build up excess lung fluid that leads to alveolar hypoxia. The decrease in fluid clearance in these conditions is due in part to down-regulation of plasma membrane adenosine triphosphatase (Zhou et al., 2008). The migration of polymorphonuclear leukocytes (PMN) into the lung plays a critical role in the development of LPS-induced acute lung injury; A₃ receptors have been claimed to inhibit this effect (Wagner et al., 2010). The protective effects of A₃ receptor activation in attenuating reperfusion lung injury are mediated in part through up-regulation of phosphorylated extracellular signal-regulated

kinase (Matot et al., 2006). Chemoattractants induce the release of ATP from migrating PMN, which is rapidly hydrolyzed to adenosine; PMN express A_{2A} and A₃ receptors that mediate suppression and stimulation of chemotaxis, respectively, and removal of adenosine or inhibition of A₃ receptors blocks PMN migration (Junger et al., 2006).

Recent reviews about adenosine receptors as promising targets for acute lung injury and ARDS are available (Schepp and Reutershan, 2008; Eckle et al., 2009).

E. Cystic Fibrosis

CF (also known as mucoviscidosis) is a hereditary disease caused by a mutation in the gene for the CFTR protein. It affects cells in the lung as well as in the intestine and in exocrine glands. CFTR is essential for maintaining water balance by enabling the transport of Cl⁻ ions across cell membranes, and its absence in CF results in the production of thick mucus. Mucociliary clearance in CF lung is limited by airway dehydration, leading to persistent bacterial infection and inflammation in the airways.

There are some outstanding reviews concerned with purinergic signaling in CF (Schultz et al., 1995; Ramsey, 1996; Weisman et al., 1998; Yerxa, 2001; Boucher, 2002, 2004, 2007; Kellerman et al., 2002; Bucheimer and Linden, 2004; Marcet and Boeynaems, 2006; Tarran et al., 2006; Clunes and Boucher, 2008).

Cloning and characterization of the *CFTR* gene was reported in 1989 (Riordan et al., 1989). Evidence for the regulation of ion transport by ATP and UTP in normal and CF human airway epithelium was described previously (Mason et al., 1991), in retrospect probably via P2Y₂ and/or P2Y₄ receptors. In particular, ATP and UTP activated Cl⁻ secretion in the nasal epithelia of patients with CF (Knowles et al., 1991). P2 receptor purinergic compounds are being explored for the treatment of CF, to bypass the defective function of CFTR, and to restore Cl⁻ secretion and/or inhibit Na⁺ absorption (Brown et al., 1991; Stutts and Boucher, 1999; Mall et al., 2000; Yerxa et al., 2002). The long-lasting P2Y₂ receptor agonist P¹-(uridine 5')-P⁴-(2'-deoxycytidine 5')tetraphosphate (INS37217) increases the duration of mucociliary clearance stimulation and therefore has significant advantages over other P2Y₂ agonists for the treatment of CF (Kellerman et al., 2002; Yerxa et al., 2002; Goralski et al., 2010). ATP regulates Cl⁻ channels in both normal and CF cultured human airway epithelia (Stutts et al., 1992) and in cultured human tracheal gland cells (Merten et al., 1993). Control of CFTR Cl⁻ conductance by ATP is through nonhydrolytic binding (Quinton and Reddy, 1992). Nucleotide activation of CFTR is by a cAMP-independent mechanism (Cantiello et al., 1994). Evidence has been provided that the anti-inflammatory mediator lipoxin A₄ is decreased in CF. In vitro experiments showed that lipoxin A₄ enhances the airway surface liquid layer height in CF and non-CF

epithelia, and this effect is inhibited by, among others, the P2Y₁₁ receptor antagonist NF340 (Urbach et al., 2012). The same authors also showed that lipoxin A₄ stimulates a carbenoxolone-sensitive apical ATP release.

A₁ receptor agonists mobilize [Ca²⁺]_i and activate K⁺ and Cl⁻ currents in normal and CF airway epithelial cells (Rugolo et al., 1993). A_{2A} receptors participate in regulation of Cl⁻ secretion via a Ca²⁺-dependent signaling pathway and appears to be at least partially conserved in CF airway epithelial cells (Chao et al., 1994). Evidence has also been presented for the presence and function of A₁ receptors in modulating Cl⁻ secretion in normal and CF airway epithelial cells (McCoy et al., 1995). A_{2B} receptors mediate activation of CFTR as well as of the clinically important CFTR mutation R117H (Clancy et al., 1999). Single-nucleotide polymorphisms of the A_{2B} receptor gene are much more frequent in African Americans than in whites, but none of the polymorphisms identified are likely to be modifiers in CF (Tang et al., 2005). Bronchoalveolar lavage from patients with CF exhibited high adenosine concentrations, which was correlated with an increase in 5'-nucleotidase and a decrease in adenosine deaminase activity (Picher et al., 2006). Regulation of purinergic signaling by NTPDase1 (CD39) and NTPDase3 is remodelled in CF (Fausther et al., 2010). A_{2B} receptors preserve epithelial barrier integrity in the lung (Eckle et al., 2008), suggesting that the colocalization of NTPDase1 and ecto 5'-nucleotidase on basolateral surfaces may contribute to the defense mechanism by locally enhancing adenosine availability. Adenosine regulates CFTR via A_{2B} receptors and prostanoids (Li et al., 2006).

An important advance was the cloning and expression of a human P2U nucleotide receptor, a target for CF pharmacotherapy (Parr et al., 1994). A murine P2U receptor cDNA was cloned from neuroblastoma-glioma hybrid (NG108-15) cells (Lustig et al., 1993). Gating of CFTR Cl⁻ channels is regulated by phosphorylation and ATP hydrolysis (Hwang et al., 1994). It was argued that incremental phosphorylation differentially regulates the interactions between nucleotides and the two nucleotide-binding domains (NBD1 and NBD2) of CFTR, and that ATP hydrolysis at one NBD controls channel opening, whereas ATP hydrolysis at the other regulates channel closing (Gunderson and Kopito, 1994; Zhou et al., 2006). NBD1 and NBD2 function as adenylate kinases, but not ATPases (Gross et al., 2006). Uridine-5'-(γ -thio)-triphosphate, a potent hydrolysis-resistant agonist at P2U receptors, was developed as a promising compound for treating CF (Lazarowski et al., 1996). Purified CFTR does not function as an ATP channel (Li et al., 1996). CF is caused by a single deletion mutation (Δ Phe508) within the first nucleotide binding fold (NBF1) of the CFTR protein (Ko et al., 1997). ATP depletion induces loss of respiratory epithelial functional integrity and down-regulates CFTR expression (Brézil-

lon et al., 1997). An ATP-activated Na⁺/H⁺ exchanger was shown to be located at the basolateral surface of nasal epithelial cells from patients with CF (Paradiso, 1997).

A characteristic feature of CF is chronic bacterial infection by *P. aeruginosa*, which accumulates diffusible *N*-acylhomoserine lactone signal molecules that activate virulence factor genes. *N*-Acylhomoserine lactone down-regulates expression of P2Y₂ and P2Y₄ receptors, altering the responses to ATP and UTP (Saleh et al., 1999a). Flagellin of *P. aeruginosa* inhibits ENaC expression in airway epithelium, reducing Na⁺ absorption to enhance local mucociliary clearance, a mechanism that is attenuated in CF (Kunzelmann et al., 2006). The action of LPS on ENaC was mediated in part by the protein kinase C and phospholipase C pathways and by purinergic signaling (Boncoeur et al., 2010). In intact distal bronchi of porcine lungs, UTP stimulated Cl⁻ secretion by a Ca²⁺-independent mechanism and inhibited Na⁺ absorption by a Ca²⁺-dependent mechanism; both effects are likely to favor increased hydration of the airway surface and may therefore be beneficial in CF (Inglis et al., 1999). Inhibition of amiloride-sensitive epithelial Na⁺ absorption by nucleotides in human normal and CF airways was reported (Mall et al., 2000). In CF airway epithelium, unlike normal epithelium, the mucosal ATP/UTP-dependent anion secretory response was mediated exclusively by [Ca²⁺]_i (Paradiso et al., 2001). Low doses of glucocorticoids, such as dexamethasone, are used for basal immunosuppressive and anti-inflammatory treatment of asthma and CF. Inhibition of ATP-mediated Cl⁻ secretion by dexamethasone in human bronchial epithelium has been reported (Urbach et al., 2002).

Extracellular zinc as well as ATP restores Cl⁻ secretion across CF airway epithelia by triggering calcium entry via P2X receptors (Zsembery et al., 2004; Hargitai et al., 2010). Synergistic stimulation with zinc and ATP suggested a significant role for the P2X₄ receptor. However, possible roles for P2X₅ and P2X₆ receptors were not excluded, and in a later article from this group, it was suggested that P2X_{4/6} heteromultimers might be involved (Liang et al., 2005). The activity of CFTR is required for deformation-induced ATP release from erythrocytes; in CF, where CFTR is normally reduced or absent, ATP was not released from erythrocytes by mechanical deformation (Sprague et al., 1998). Cyclic compressive stress, mimicking normal tidal breathing, results in ATP release and thereby regulates airway surface liquid in normal lungs and improves clearance in the lungs of patients with CF (Button et al., 2007; Button and Boucher, 2008). Dysregulation of extracellular pH regulation is characteristic of CF and involves purinergic signaling (Kaunitz and Akiba, 2009). Maneuvers directed at increasing motion-induced nucleotide release may be therapeutic for patients with CF (Tarran et al., 2005).

Women with CF exhibit reduced survival compared with men. It was suggested that estrogen may reduce the breathing-induced ATP release and ATP receptor-mediated increase in $[Ca^{2+}]_i$ that results in Cl^- secretion (Coakley et al., 2008). Antiestrogens may therefore be beneficial in the treatment of CF because they increase Cl^- secretion. Bestrophines are up-regulated during inflammation and augment Ca^{2+} transients elicited via $P2Y_2$ receptors, influencing CF (Barro-Soria et al., 2010).

F. Lung Cancer

A549 human lung epithelial-like adenocarcinoma cells express $P2U$ (i.e., $P2Y_2$ and/or $P2Y_4$) receptors, which, when occupied, lead to increases in $[Ca^{2+}]_i$ (Clunes and Kemp, 1996) but do not inhibit forskolin-evoked cAMP accumulation in these cells (Remsbury et al., 1996). Calcium-dependent release of ATP and UTP (with subsequent increase in adenosine levels) from A549 cells has been reported (Tatur et al., 2008).

A phase II study of intravenous ATP in patients with previously untreated non-small-cell lung cancer led to the conclusion that ATP, at least at the dose and administration schedule employed, was an inactive agent in patients with advanced non-small-cell lung cancer (Haskell et al., 1998).

Erythromycin is widely used in the treatment of respiratory tract infections. It has also been shown to selectively inhibit the Ca^{2+} influx induced through $P2X_4$ receptor activation in A549 lung tumor cells (Zhao et al., 2000). In this study, it was also shown with RT-PCR that A549 cells express $P2Y_2$, $P2Y_4$, and $P2Y_6$ as well as $P2X_4$ receptors.

Cachexia is a common feature of patients with lung cancer and is associated with metabolic alterations, including elevated lipolysis, proteolysis, and gluconeogenesis. An increase in glucose turnover during high-dose ATP infusion in patients with advanced non-small-cell lung cancer was reported, perhaps contributing to the reported beneficial effects of ATP on body weight in patients with advanced lung cancer (Agteresch et al., 2000a). Later randomized clinical trials led to the conclusion that ATP has beneficial effects on weight, muscle strength, and quality of life in patients with advanced non-small-cell lung cancer, as well as on enhancing median survival from 3.5 to 9.3 months (Agteresch et al., 2000b; Dagnelie and Agteresch, 2004; Beijer et al., 2009). ATP infusion restores hepatic energy levels in patients with advanced lung cancer, especially in patients who are losing weight (Leij-Halfwerk et al., 2002). ATP has been claimed to reduce radiation-induced damage (Swennen et al., 2008a), and clinical trials are under way to assess the effect of concurrent ATP and radiotherapy treatment on outcome in patients with non-small-cell lung cancer (Arts et al., 2008).

ATP induced a significant dose-dependent growth inhibition of five different cell lines: human large-cell lung

carcinoma (H460), human papillary lung adenocarcinoma (H441), human squamous cell lung carcinoma (H520), human small cell lung carcinoma (GLC4), and human mesothelioma (MER082) (Agteresch et al., 2003). ATP also had cytotoxic effects on the PC14 lung adenocarcinoma cell line and enhanced the antitumor effect of etoposide (VP16) in both PC14 and A549 cell lines (Hatta et al., 2004). Calu-3 cells are derived from pleural effusion associated with human lung adenocarcinoma, and both mucin and ATP were released from these cells, probably in response to $P2Y_2$ receptor activation (Kreda et al., 2007). It has been claimed that extracellular ATP, UTP, and UDP stimulate proliferation of A549 lung tumor cells via $P2Y_2$ and $P2Y_6$ receptors, as well as an ADP-sensitive receptor that was not the $P2Y_1$ subtype (Schafer et al., 2003).

ATP-based chemotherapy response assay has been used to guide the outcome of platinum-based drug chemotherapy for nonresectable non-small-cell lung cancer (Moon et al., 2007, 2009). It has been suggested that tumor-infiltrating immune cells can benefit the tumor by producing factors that promote angiogenesis and suppress immunity; adenosine levels are high in tumors. A_{2B} receptors on host immune cells may participate in these effects, and A_{2B} receptor knockout mice exhibited significantly attenuated growth in a Lewis lung carcinoma isograft model (Ryzhov et al., 2008). Cisplatin is a widely used anticancer agent for the treatment of lung cancer. ATP increased the cytotoxicity of cisplatin in a human large-cell lung carcinoma cell line (H460) (Swennen et al., 2008b, 2010).

Exposure of human lung cancer cell lines A549 and H1299 to 8-chloro-adenosine induced cell arrest at the G_2/M phase and mitotic catastrophe, followed by apoptosis (Zhang et al., 2004; Gu et al., 2006; Li et al., 2009). 3'-Deoxyadenosine (cordycepin) exerted an inhibitory effect on the growth of the mouse Lewis lung carcinoma cell line by stimulating A_3 receptors (Nakamura et al., 2006). The A_3 receptor agonist thio-2-chloro- N^6 -(3-iodobenzyl)-5'- N -methylcarboxamidoadenosine inhibited cell proliferation through cell-cycle arrest and apoptosis of A549 human lung carcinoma cells (Kim et al., 2008).

G. Rhinosinusitis

Rhinosinusitis refers to inflammation of the nasal sinuses due to injury, allergy, or autoimmunity. Inhaled AMP caused airway narrowing in a significantly higher proportion of subjects with allergic rhinitis (Prieto et al., 2001). A randomized, double-blind trial concluded that novel agonists and antagonists to A_{2A} and A_3 receptors had limited clinical benefit in both early- and late-phase responses to intranasal allergen challenge (Rimmer et al., 2007). Reversal of chronic rhinosinusitis-associated sinonasal ciliary dysfunction, as the result of stimulation by ATP, was reported (Chen et al., 2007).

H. Pulmonary Hypertension

See also section I.E. Adenosine is an effective vasodilator in patients with pulmonary hypertension (Schrader et al., 1992). In a recent article, it has been shown that in rats with hypobaric hypoxia-induced pulmonary hypertension, there is a significant increase in P2X4 receptor mRNA and protein expression and in P2X1 receptor protein in the right ventricle (Ohata et al., 2011).

VII. Concluding Comments

In airway epithelia, extracellular ATP contributes to mucous ciliary clearance, a critical process required to maintain the airways clear of inhaled particles or pathogens. It mediates Cl^- , K^+ , and fluid secretion; inhibits Na^+ absorption; increases ciliary beat frequency; potentiates regulatory volume decreases after hypotonic cell swelling; and triggers mucin release. These responses involve P2Y₂ and P2X4 and P2X7 receptors. ATP seems to be involved in signal transduction in a multitude of sensory lung receptors (NEBs, SMARs, visceral pleura receptors) via P2X3 receptors on vagal and spinal sensory nerves. Considering the stem cell characteristics of Clara-like cells in the NEB microenvironment, local purinergic signaling may be of great importance for normal airway function, for airway regeneration after injury, and/or for the pathogenesis of small-cell lung carcinomas.

As should be clear from the preceding extensive lists of examples, purinergic signaling, via different subtypes of P1 and P2 purinoceptors, seems to be involved in the physiological regulation of normal functions in virtually all cell and tissue types in the airways. In addition, the reported expression profiles of the different purinoceptor types may be up-regulated and/or changed as a consequence of pathological modifications.

Regarding the potential sources of extracellular purines involved in these signaling pathways, the situation also turned out to be complex. Besides a considerable number of cell types that release ATP as a transmitter via stimulated and strictly regulated exocytosis, many cells and tissues have been shown to release ATP via different mechanisms, such as connexin or pannexin hemichannels, voltage-dependent anion and P2X7 receptor channels, and vesicular release, and as a response to physiological stimuli (e.g., mechanical, hypoxia, osmotic).

In normal physiological situations, functionally released ATP is rapidly degraded by strategically located ectonucleotidases. In pathophysiological situations, however, the extracellular environment often seems to contain aberrant concentrations of ATP, not only because of enhanced stimulation and release by "normal sources" and/or insufficient breakdown but also because most cells harbor considerable cytoplasmic amounts of ATP that may be massively released in a noncontrolla-

ble way when cells or tissues are damaged (e.g., inflammation, physical damage).

All together, the ubiquitous expression of the many subtypes of purinoceptors in all cells and tissues that function in the multitude of normal physiological roles of extracellular ATP, in combination with the often enhanced and uncontrolled availability of extracellular ATP, unfortunately represents a huge potential for pathological purinergic "cross-signaling" between different cell and tissue types. On the other hand, thorough knowledge of the receptor subtypes and the pathways involved may also create interesting possibilities for targeted therapeutic interventions.

Authorship Contributions

Wrote or contributed to the writing of the manuscript: Burnstock, Brouns, Adriaensen, and Timmermans.

References

- Abbraccio MP and Burnstock G (1994) Purinoceptors: are there families of P_{2X} and P_{2Y} purinoceptors? *Pharmacol Ther* **64**:445–475.
- Abdullah LH, Bundy JT, Ehre C, and Davis CW (2003) Mucin secretion and PKC isoforms in SPOC1 goblet cells: differential activation by purinergic agonist and PMA. *Am J Physiol Lung Cell Mol Physiol* **285**:L149–L160.
- Abdullah LH, Davis SW, Burch L, Yamauchi M, Randell SH, Nettesheim P, and Davis CW (1996) P_{2U} purinoceptor regulation of mucin secretion in SPOC1 cells, a goblet cell line from the airways. *Biochem J* **316**:943–951.
- Abebe W and Mustafa SJ (1998) A₁ adenosine receptor-mediated Ins(1,4,5)P₃ generation in allergic rabbit airway smooth muscle. *Am J Physiol* **275**:L990–L997.
- Abman SH, Chatfield BA, Rodman DM, Hall SL, and McMurtry IF (1991) Maturation changes in endothelium-derived relaxing factor activity of ovine pulmonary arteries in vitro. *Am J Physiol* **260**:L280–L285.
- Abu-Shaweesh JM (2007) Activation of central adenosine A_{2A} receptors enhances superior laryngeal nerve stimulation-induced apnea in piglets via a GABAergic pathway. *J Appl Physiol* **103**:1205–1211.
- Ackland GL, Kasymov V, and Gourine AV (2007) Physiological and pathophysiological roles of extracellular ATP in chemosensory control of breathing. *Biochem Soc Trans* **35**:1264–1268.
- Adkins WK, Barnard JW, Moore TM, Allison RC, Prasad VR, and Taylor AE (1993) Adenosine prevents PMA-induced lung injury via an A₂ receptor mechanism. *J Appl Physiol* **74**:982–988.
- Adriaensen D and Timmermans JP (2004) Purinergic signalling in the lung: important in asthma and COPD? *Curr Opin Pharmacol* **4**:207–214.
- Advenier C, Devillier P, Matran R, and Naline E (1988) Influence of epithelium on the responsiveness of guinea-pig isolated trachea to adenosine. *Br J Pharmacol* **93**:295–302.
- Agteresch HJ, Dagnelie PC, van der Gaast A, Stijnen T, and Wilson JH (2000a) Randomized clinical trial of adenosine 5'-triphosphate in patients with advanced non-small-cell lung cancer. *J Natl Cancer Inst* **92**:321–328.
- Agteresch HJ, Leij-Halfwerk S, Van Den Berg JW, Hordijk-Luijk CH, Wilson JH, and Dagnelie PC (2000b) Effects of ATP infusion on glucose turnover and gluconeogenesis in patients with advanced non-small-cell lung cancer. *Clin Sci (Lond)* **98**:689–695.
- Agteresch HJ, van Rooijen MHC, van den Berg JW, Minderman-Voortman GJ, Wilson JH, and Dagnelie PC (2003) Growth inhibition of lung cancer cells by adenosine 5-triphosphate. *Drug Dev Res* **60**:196–203.
- Ahmad A, Ahmad S, Glover L, Miller SM, Shannon JM, Guo X, Franklin WA, Bridges JP, Schaack JB, Colgan SP, et al. (2009) Adenosine A_{2A} receptor is a unique angiogenic target of HIF-2 α in pulmonary endothelial cells. *Proc Natl Acad Sci USA* **106**:10684–10689.
- Ahmad S, Ahmad A, McConville G, Schneider BK, Allen CB, Manzer R, Mason RJ, and White CW (2005) Lung epithelial cells release ATP during ozone exposure: signaling for cell survival. *Free Radic Biol Med* **39**:213–226.
- Ahmad S, Ahmad A, and White CW (2006) Purinergic signaling and kinase activation for survival in pulmonary oxidative stress and disease. *Free Radic Biol Med* **41**:29–40.
- Aksoy MO, Borenstein M, Li XX, and Kelsen SG (1995) Eicosanoid production in rabbit tracheal epithelium by adenine nucleotides: mediation by P₂-purinoceptors. *Am J Respir Cell Mol Biol* **13**:410–417.
- Alexander I and Eyre P (1985) P1-purinoceptors mediate relaxation of the bovine bronchial artery. *Eur J Pharmacol* **107**:359–362.
- Ali S, Metzger WJ, Olanrewaju HA, and Mustafa SJ (1997) Adenosine receptor-mediated relaxation of rabbit airway smooth muscle: a role for nitric oxide. *Am J Physiol* **273**:L581–L587.
- Allen-Gipson DS, Wong J, Spurzem JR, Sisson JH, and Wyatt TA (2006) Adenosine A_{2A} receptors promote adenosine-stimulated wound healing in bronchial epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **290**:L849–L855.
- Allen-Gipson DS, Spurzem K, Kolm N, Spurzem JR, and Wyatt TA (2007) Adenosine promotion of cellular migration in bronchial epithelial cells is mediated by the activation of cyclic adenosine monophosphate-dependent protein kinase A. *J Invest Med* **55**:378–385.

- Ando J and Yamamoto K (2009) Vascular mechanobiology: endothelial cell responses to fluid shear stress. *Circ J* **73**:1983–1992.
- Antunes VR, Braga VA, and Machado BH (2005) Autonomic and respiratory responses to microinjection of ATP into the intermediate or caudal nucleus tractus solitarius in the working heart-brainstem preparation of the rat. *Clin Exp Pharmacol Physiol* **32**:467–472.
- Arts IC, Swennen EL, Vandeurzen KG, and Dagnelie PC (2008) Effect of ATP on survival, tumour response, nutritional status and quality of life in lung cancer patients: a multicentre, double-blind randomized trial. *Purinergic Signal* **4**:S194–S195.
- Baek EB, Yoo HY, Park SJ, Kim HS, Kim SD, Earm YE, and Kim SJ (2008) Luminal ATP-induced contraction of rabbit pulmonary arteries and role of purinoceptors in the regulation of pulmonary arterial pressure. *Pflugers Arch* **457**:281–291.
- Bakand S, Winder C, Khalil C, and Hayes A (2006) An experimental in vitro model for dynamic direct exposure of human cells to airborne contaminants. *Toxicol Lett* **165**:1–10.
- Bakand S, Winder C, and Hayes A (2007) Comparative in vitro cytotoxicity assessment of selected gaseous compounds in human alveolar epithelial cells. *Toxicol In Vitro* **21**:1341–1347.
- Banerjee SK, Young HW, Barczak A, Erle DJ, and Blackburn MR (2004) Abnormal alveolar development associated with elevated adenine nucleosides. *Am J Respir Cell Mol Biol* **30**:38–50.
- Banerjee SK, Young HW, Volmer JB, and Blackburn MR (2002) Gene expression profiling in inflammatory airway disease associated with elevated adenosine. *Am J Physiol Lung Cell Mol Physiol* **282**:L169–L182.
- Bao HF, Chen XJ, Jian L, and Eaton DC (2007) Purinergic regulation of amiloride-sensitive lung sodium channels (Abstract). *FASEB J* **21**:606.20.
- Barker PM and Gatzky JT (1998) Effects of adenosine, ATP, and UTP on chloride secretion by epithelia explanted from fetal rat lung. *Pediatr Res* **43**:652–659.
- Barro-Soria R, Aldehni F, Almacá J, Witzgall R, Schreiber R, and Kunzelmann K (2010) ER-localized bestrophin 1 activates Ca²⁺-dependent ion channels TMEM16A and SK4 possibly by acting as a counterion channel. *Pflugers Arch* **459**:485–497.
- Barros RC and Branco LG (2000) Role of central adenosine in the respiratory and thermoregulatory responses to hypoxia. *Neuroreport* **11**:193–197.
- Barth K and Kasper M (2009) Membrane compartments and purinergic signalling: occurrence and function of P2X receptors in lung. *FEBS J* **276**:341–353.
- Barth K, Weinhold K, Guenther A, Linge A, Gereke M, and Kasper M (2008) Characterization of the molecular interaction between caveolin-1 and the P2X receptors 4 and 7 in E10 mouse lung alveolar epithelial cells. *Int J Biochem Cell Biol* **40**:2230–2239.
- Barth K, Weinhold K, Guenther A, Young MT, Schnittler H, and Kasper M (2007) Caveolin-1 influences P2X₇ receptor expression and localization in mouse lung alveolar epithelial cells. *FEBS J* **274**:3021–3033.
- Basoglu OK, Pelleg A, Essilfie-Quaye S, Brindicci C, Barnes PJ, and Kharitonov SA (2005) Effects of aerosolized adenosine 5'-triphosphate vs adenosine 5'-monophosphate on dyspnea and airway caliber in healthy nonsmokers and patients with asthma. *Chest* **128**:1905–1909.
- Beijer S, Hupperets PS, van den Boorne BE, Eussen SR, van Henten AM, van den Beuken-van Everdingen M, de Graaf A, Ambergen TA, van den Brandt PA, and Dagnelie PC (2009) Effect of adenosine 5'-triphosphate infusions on the nutritional status and survival of preterminal cancer patients. *Anticancer Drugs* **20**:625–633.
- Benali R, Pierrot D, Zahm JM, de Bentzmann S, and Puchelle E (1994) Effect of extracellular ATP and UTP on fluid transport by human nasal epithelial cells in culture. *Am J Respir Cell Mol Biol* **10**:363–368.
- Bergner A and Sanderson MJ (2002) ATP stimulates Ca²⁺ oscillations and contraction in airway smooth muscle cells of mouse lung slices. *Am J Physiol Lung Cell Mol Physiol* **283**:L1271–L1279.
- Biaggioni I, King LS, Enayat N, Robertson D, and Newman JH (1989) Adenosine produces pulmonary vasoconstriction in sheep. Evidence for thromboxane A₂/prostaglandin endoperoxide-receptor activation. *Circ Res* **65**:1516–1525.
- Bianchi AL, Denavit-Saubié M, and Champagnat J (1995) Central control of breathing in mammals: neuronal circuitry, membrane properties, and neurotransmitters. *Physiol Rev* **75**:1–45.
- Bissonnette JM, Hohimer AR, and Knopp SJ (1991) The effect of centrally administered adenosine on fetal breathing movements. *Respir Physiol* **84**:273–285.
- Björck T, Gustafsson LE, and Dahlén SE (1992) Isolated bronchi from asthmatics are hyperresponsive to adenosine, which apparently acts indirectly by liberation of leukotrienes and histamine. *Am Rev Respir Dis* **145**:1087–1091.
- Blackburn MR (2003) Too much of a good thing: adenosine overload in adenosine deaminase-deficient mice. *Trends Pharmacol Sci* **24**:66–70.
- Blackburn MR (2007) A role for neural pathways in adenosine-induced bronchoconstriction. *Am J Physiol Lung Cell Mol Physiol* **293**:L22–L24.
- Blackburn MR, Lee CG, Young HW, Zhu Z, Chunn JL, Kang MJ, Banerjee SK, and Elias JA (2003) Adenosine mediates IL-13-induced inflammation and remodeling in the lung and interacts in an IL-13-adenosine amplification pathway. *J Clin Invest* **112**:332–344.
- Bochenek G, Nizankowska E, Gielicz A, and Szczeklik A (2008) Mast cell activation after adenosine inhalation challenge in patients with bronchial asthma. *Allergy* **63**:140–141.
- Bodin P and Burnstock G (2001) Evidence that release of ATP from endothelial cells during increased shear stress is vesicular. *J Cardiovasc Pharmacol* **38**:900–908.
- Boncoeur E, Tardif V, Tessier MC, Morneau F, Lavoie J, Gendreau-Berthiaume E, Grygorczyk R, Dagenais A, and Berthiaume Y (2010) Modulation of epithelial sodium channel activity by lipopolysaccharide in alveolar type II cells: involvement of purinergic signaling. *Am J Physiol Lung Cell Mol Physiol* **298**:L417–L426.
- Bonneau O, Wyss D, Ferretti S, Blaydon C, Stevenson CS, and Trifilieff A (2006) Effect of adenosine A_{2A} receptor activation in murine models of respiratory disorders. *Am J Physiol Lung Cell Mol Physiol* **290**:L1036–L1043.
- Boots AW, Hristova M, Kasahara DI, Haenen GR, Bast A, and van der Vliet A (2009) ATP-mediated activation of the NADPH oxidase DUOX1 mediates airway epithelial responses to bacterial stimuli. *J Biol Chem* **284**:17858–17867.
- Boucher RC (2002) An overview of the pathogenesis of cystic fibrosis lung disease. *Adv Drug Deliv Rev* **54**:1359–1371.
- Boucher RC (2004) New concepts of the pathogenesis of cystic fibrosis lung disease. *Eur Respir J* **23**:146–158.
- Boucher RC (2007) Airway surface dehydration in cystic fibrosis: pathogenesis and therapy. *Annu Rev Med* **58**:157–170.
- Bowler JW, Bailey RJ, North RA, and Surprenant A (2003) P2X₄, P2Y₁ and P2Y₂ receptors on rat alveolar macrophages. *Br J Pharmacol* **140**:567–575.
- Brackett LE, Shamim MT, and Daly JW (1990) Activities of caffeine, theophylline, and enprofylline analogs as tracheal relaxants. *Biochem Pharmacol* **39**:1897–1904.
- Braiman A, Silberberg SD, and Priel Z (2000) Purinergic stimulation of ciliary activity. *Drug Dev Res* **50**:550–554.
- Breschi MC, Blandizzi C, Fogli S, Martinelli C, Adinolfi B, Calderone V, Camici M, Martinotti E, and Nieri P (2007) In vivo adenosine A_{2B} receptor desensitization in guinea-pig airway smooth muscle: implications for asthma. *Eur J Pharmacol* **575**:149–157.
- Breschi MC, Nieri P, Lazzeri N, and Macchia M (1998) Potentiation by adenosine of histamine-induced bronchospasm in anaesthetized guinea-pig: receptor subtype/s involved. *Drug Dev Res* **45**:17–22.
- Brézillon S, Zahm JM, Pierrot D, Gaillard D, Hinnrasky J, Millart H, Klossek JM, Tümmler B, and Puchelle E (1997) ATP depletion induces a loss of respiratory epithelium functional integrity and down-regulates CFTR (cystic fibrosis transmembrane conductance regulator) expression. *J Biol Chem* **272**:27830–27838.
- Briva A, Santos C, Malacrida L, Rocchiccioli F, Soto J, Angulo M, Batthyany C, Cairoli E, and Piriz H (2011) Adenosine triphosphate-dependent calcium signaling during ventilator-induced lung injury is amplified by hypercapnia. *Exp Lung Res* **37**:471–481.
- Brouns I, Adriaensens D, Burnstock G, and Timmermans JP (2000) Intraepithelial vagal sensory nerve terminals in rat pulmonary neuroepithelial bodies express P2X₃ receptors. *Am J Respir Cell Mol Biol* **23**:52–61.
- Brouns I, De Proost I, Pintelon I, Timmermans JP, and Adriaensens D (2006) Sensory receptors in the airways: neurochemical coding of smooth muscle-associated airway receptors and pulmonary neuroepithelial body innervation. *Auton Neurosci* **126–127**:307–319.
- Brouns I, Oztay F, Pintelon I, De Proost I, Lembrechts R, Timmermans JP, and Adriaensens D (2009) Neurochemical pattern of the complex innervation of neuroepithelial bodies in mouse lungs. *Histochem Cell Biol* **131**:55–74.
- Brouns I, Pintelon I, Timmermans JP, and Adriaensens D (2012) Novel insights in the neurochemistry and function of pulmonary sensory receptors. *Adv Anat Embryol Cell Biol* **211**:1–115, vii.
- Brouns I, Van Genechten J, Burnstock G, Timmermans JP, and Adriaensens D (2003a) Ontogenesis of P2X₃ receptor-expressing nerve fibres in the rat lung, with special reference to neuroepithelial bodies. *Biomed Res* **14**:80–86.
- Brouns I, Van Genechten J, Hayashi H, Gajda M, Gomi T, Burnstock G, Timmermans JP, and Adriaensens D (2003b) Dual sensory innervation of pulmonary neuroepithelial bodies. *Am J Respir Cell Mol Biol* **28**:275–285.
- Brown CM and Collis MG (1982) Evidence for an A₂/R_a adenosine receptor in the guinea-pig trachea. *Br J Pharmacol* **76**:381–387.
- Brown HA, Lazarowski ER, Boucher RC, and Harden TK (1991) Evidence that UTP and ATP regulate phospholipase C through a common extracellular 5'-nucleotide receptor in human airway epithelial cells. *Mol Pharmacol* **40**:648–655.
- Brown RA, Clarke GW, Ledbetter CL, Hurler MJ, Denyer JC, Simcock DE, Coote JE, Savage TJ, Murdoch RD, Page CP, et al. (2008) Elevated expression of adenosine A₁ receptor in bronchial biopsy specimens from asthmatic subjects. *Eur Respir J* **31**:311–319.
- Brown RA, Spina D, and Page CP (2008) Adenosine receptors and asthma. *Br J Pharmacol* **153** (Suppl 1):S446–S456.
- Brugós L, Gesztelyi R, Zsuga J, Cseppento A, Benko I, Galajda Z, Deák G, Sipka S, Roszer T, Kovács P, et al. (2007) Modulation of adenosine-induced responses in the guinea-pig trachea during long-term caffeine treatment: possible role of epithelium. *J Pharmacol Sci* **105**:279–290.
- Bucheimer RE and Linden J (2004) Purinergic regulation of epithelial transport. *J Physiol* **555**:311–321.
- Buell G, Collo G, and Rassendren F (1996) P2X receptors: an emerging channel family. *Eur J Neurosci* **8**:2221–2228.
- Buenestado A, Grassini Delyle S, Arnold I, Besnard F, Naline E, Blouquit-Laye S, Chapelier A, Bellamy JF, and Devillier P (2010) The role of adenosine receptors in regulating production of tumour necrosis factor- α and chemokines by human lung macrophages. *Br J Pharmacol* **159**:1304–1311.
- Burki NK, Alam M, and Lee LY (2006) The pulmonary effects of intravenous adenosine in asthmatic subjects. *Respir Res* **7**:139.
- Burki NK, Dale WJ, and Lee LY (2005) Intravenous adenosine and dyspnea in humans. *J Appl Physiol* **98**:180–185.
- Burnstock G (1972) Purinergic nerves. *Pharmacol Rev* **24**:509–581.
- Burnstock G (1978) A basis for distinguishing two types of purinergic receptor, in *Cell Membrane Receptors for Drugs and Hormones: A Multidisciplinary Approach* (Straub RW and Bolis L eds) pp 107–118, Raven Press, New York.
- Burnstock G (1988) Autonomic neural control mechanisms with special reference to the airways, in *The Airways. Neural Control in Health and Disease* (Kaliner MA and Barnes PJ eds) pp 1–22, Marcel Dekker, New York.
- Burnstock G (1999) Release of vasoactive substances from endothelial cells by shear stress and purinergic mechanosensory transduction. *J Anat* **194**:335–342.
- Burnstock G (2007a) Physiology and pathophysiology of purinergic neurotransmission. *Physiol Rev* **87**:659–797.
- Burnstock G (2007b) Purine and pyrimidine receptors. *Cell Mol Life Sci* **64**:1471–1483.
- Burnstock G (2009a) Autonomic neurotransmission: 60 years since Sir Henry Dale. *Annu Rev Pharmacol Toxicol* **49**:1–30.

- Burnstock G (2009b) Purinergic mechanosensory transduction and visceral pain. *Molecular Pain* **5**:69.
- Burnstock G (2012) Purinergic signalling: its unpopular beginning, its acceptance and its exciting future. *BioEssays* **34**:218–225.
- Burnstock G and Kennedy C (1985) Is there a basis for distinguishing two types of P₂-purinoceptor? *Gen Pharmacol* **16**:433–440.
- Burnstock G and Knight GE (2004) Cellular distribution and functions of P2 receptor subtypes in different systems. *Int Rev Cytol* **240**:31–304.
- Button B, Boucher RC, and University of North Carolina Virtual Lung Group (2008) Role of mechanical stress in regulating airway surface hydration and mucus clearance rates. *Respir Physiol Neurobiol* **163**:189–201.
- Button B, Picher M, and Boucher RC (2007) Differential effects of cyclic and constant stress on ATP release and mucociliary transport by human airway epithelia. *J Physiol* **580**:577–592.
- Cagnina RE, Ramos SI, Marshall MA, Wang G, Frazier CR, and Linden J (2009) Adenosine A_{2B} receptors are highly expressed on murine type II alveolar epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **297**:L467–L474.
- Candenas ML, Devillier P, Naline E, and Advenier C (1992) Contractile effect of α,β -methylene ATP on the guinea-pig isolated trachea. *Fundam Clin Pharmacol* **6**:135–144.
- Cantiello HF, Prat AG, Reisin IL, Ercole LB, Abraham EH, Amara JF, Gregory RJ, and Ausiello DA (1994) External ATP and its analogs activate the cystic fibrosis transmembrane conductance regulator by a cyclic AMP-independent mechanism. *J Biol Chem* **269**:11224–11232.
- Caruso M, Holgate ST, and Polosa R (2006) Adenosine signalling in airways. *Curr Opin Pharmacol* **6**:251–256.
- Caruso M, Varani K, Tringali G, and Polosa R (2009) Adenosine and adenosine receptors: their contribution to airway inflammation and therapeutic potential in asthma. *Curr Med Chem* **16**:3875–3885.
- Chambers LA, Clunes MT, Inglis SK, Olver RE, and Wilson SM (2002) Multiple P2Y receptors present on apical and basolateral membranes of Calu-3 cells. *J Physiol* **544**:95P.
- Chander A, Sen N, Wu AM, and Spitzer AR (1995) Protein kinase C in ATP regulation of lung surfactant secretion in type II cells. *Am J Physiol* **268**:L108–L116.
- Chao AC, Zifferblatt JB, Wagner JA, Dong YJ, Gruenert DC, and Gardner P (1994) Stimulation of chloride secretion by P₁ purinoceptor agonists in cystic fibrosis phenotype airway epithelial cell line CFPEo-. *Br J Pharmacol* **112**:169–175.
- Chen B, Antunes MB, Claire SE, Palmer JN, Chiu AG, Kennedy DW, and Cohen NA (2007) Reversal of chronic rhinosinusitis-associated sinonasal ciliary dysfunction. *Am J Rhinol* **21**:346–353.
- Chen BC, Lee CM, Lee YT, and Lin WW (1996) Characterization of signaling pathways of P_{2Y} and P_{2U} purinoceptors in bovine pulmonary artery endothelial cells. *J Cardiovasc Pharmacol* **28**:192–199.
- Chen BC and Lin WW (1997) Inhibition of ecto-ATPase by the P₂ purinoceptor agonists, ATP γ S, α,β -methylene-ATP, and AMP-PNP, in endothelial cells. *Biochem Biophys Res Commun* **233**:442–446.
- Chen BC and Lin WW (1999) PKC β 1 mediates the inhibition of P2Y receptor-induced inositol phosphate formation in endothelial cells. *Br J Pharmacol* **127**:1908–1914.
- Chen L, Song W, Davis IC, Shrestha K, Schwiebert E, Sullender WM, and Matalon S (2009) Inhibition of Na⁺ transport in lung epithelial cells by respiratory syncytial virus infection. *Am J Respir Cell Mol Biol* **40**:588–600.
- Chen WT, Huang WH, Wang D, Yu FC, Chi YC, Wu JC, Wu K, Perng WC, Wu CP, and Yan HC (2003) The protective effect of adenosine triphosphate-MgCl₂ on ischemia-reperfusion lung injury is leukocyte dependent. *J Biomed Sci* **10**:725–730.
- Chen Y, Zhao YH, and Wu R (2001) Differential regulation of airway mucin gene expression and mucin secretion by extracellular nucleotide triphosphates. *Am J Respir Cell Mol Biol* **25**:409–417.
- Cheng DY, DeWitt BJ, Suzuki F, Neely CF, and Kadowitz PJ (1996) Adenosine A₁ and A₂ receptors mediate tone-dependent responses in feline pulmonary vascular bed. *Am J Physiol* **270**:H200–H207.
- Choi JY, Namkung W, Shin JH, and Yoon JH (2003) Uridine-5'-triphosphate and adenosine triphosphate γ S induce mucin secretion via Ca²⁺-dependent pathways in human nasal epithelial cells. *Acta Otolaryngol* **123**:1080–1086.
- Chootip K, Ness KF, Wang Y, Gurney AM, and Kennedy C (2002) Regional variation in P2 receptor expression in the rat pulmonary arterial circulation. *Br J Pharmacol* **137**:637–646.
- Chuaychoo B, Lee MG, Kollarik M, Pullmann R Jr, and Udem BJ (2006) Evidence for both adenosine A₁ and A_{2A} receptors activating single vagal sensory C-fibres in guinea pig lungs. *J Physiol* **575**:481–490.
- Chunn JL, Molina JG, Mi T, Xia Y, Kellems RE, and Blackburn MR (2005) Adenosine-dependent pulmonary fibrosis in adenosine deaminase-deficient mice. *J Immunol* **175**:1937–1946.
- Cicko S, Lucattelli M, Müller T, Lommatzsch M, De Cunto G, Cardini S, Sundas W, Grimm M, Zeiser R, Dürk T, et al. (2010) Purinergic receptor inhibition prevents the development of smoke-induced lung injury and emphysema. *J Immunol* **185**:688–697.
- Clancy JP, Ruiz FE, and Sorscher EJ (1999) Adenosine and its nucleotides activate wild-type and R117H CFTR through an A_{2B} receptor-coupled pathway. *Am J Physiol* **276**:C361–C369.
- Clarke LL and Boucher RC (1992) Chloride secretory response to extracellular ATP in human normal and cystic fibrosis nasal epithelia. *Am J Physiol* **263**:C348–C356.
- Clarke LL, Chinnet T, and Boucher RC (1997) Extracellular ATP stimulates K⁺ secretion across cultured human airway epithelium. *Am J Physiol* **272**:L1084–L1091.
- Clunes MT and Boucher RC (2008) Front-runners for pharmacotherapeutic correction of the airway ion transport defect in cystic fibrosis. *Curr Opin Pharmacol* **8**:292–299.
- Clunes MT, Collett A, Baines DL, Bovell DL, Murphie H, Inglis SK, McAlroy HL, Olver RE, and Wilson SM (1998) Culture substrate-specific expression of P2Y₂ receptors in distal lung epithelial cells isolated from foetal rats. *Br J Pharmacol* **124**:845–847.
- Clunes MT and Kemp PJ (1996) P_{2U} purinoceptor modulation of intracellular Ca²⁺ in a human lung adenocarcinoma cell line: down-regulation of Ca²⁺ influx by protein kinase C. *Cell Calcium* **20**:339–346.
- Coakley RD, Sun H, Clunes LA, Rasmussen JE, Stackhouse JR, Okada SF, Fricks I, Young SL, and Tarran R (2008) 17 β -Estradiol inhibits Ca²⁺-dependent homeostasis of airway surface liquid volume in human cystic fibrosis airway epithelia. *J Clin Invest* **118**:4025–4035.
- Coburn RF and Tomita T (1973) Evidence for nonadrenergic inhibitory nerves in the guinea pig trachealis muscle. *Am J Physiol* **224**:1072–1080.
- Coleman RA and Levy GP (1974) A non-adrenergic inhibitory nervous pathway in guinea-pig trachea. *Br J Pharmacol* **52**:167–174.
- Collo G, North RA, Kawashima E, Merlo-Pich E, Neidhart S, Surprenant A, and Buell G (1996) Cloning of P2X₅ and P2X₆ receptors and the distribution and properties of an extended family of ATP-gated ion channels. *J Neurosci* **16**:2495–2507.
- Communi D, Paindavoine P, Place GA, Parmentier M, and Boeynaems JM (1999) Expression of P2Y receptors in cell lines derived from the human lung. *Br J Pharmacol* **127**:562–568.
- Conway JD, Bartolotta T, Abdullah LH, and Davis CW (2003) Regulation of mucin secretion from human bronchial epithelial cells grown in murine hosted xenografts. *Am J Physiol Lung Cell Mol Physiol* **284**:L945–L954.
- Cressman VL, Hicks EM, Funkhouser WK, Backlund DC, and Koller BH (1998) The relationship of chronic mucin secretion to airway disease in normal and CFTR-deficient mice. *Am J Respir Cell Mol Biol* **19**:853–866.
- Cressman VL, Lazarowski E, Homolya L, Boucher RC, Koller BH, and Grubb BR (1999) Effect of loss of P2Y₂ receptor gene expression on nucleotide regulation of murine epithelial Cl⁻ transport. *J Biol Chem* **274**:26461–26468.
- Crimi N, Palermo F, Oliveri R, Maccarrone C, Palermo B, Vancheri C, Polosa R, and Mistretta A (1988) Enhancing effect of dipyrindamole inhalation on adenosine-induced bronchospasm in asthmatic patients. *Allergy* **43**:179–183.
- Crimi N, Palermo F, Polosa R, Oliveri R, Maccarrone C, Palermo B, and Mistretta A (1989) Effect of indomethacin on adenosine-induced bronchoconstriction. *J Allergy Clin Immunol* **83**:921–925.
- Cruz CM, Rinna A, Forman HJ, Ventura AL, Persechini PM, and Ojcius DM (2007) ATP activates a reactive oxygen species-dependent oxidative stress response and secretion of proinflammatory cytokines in macrophages. *J Biol Chem* **282**:2871–2879.
- Cushley MJ, Tattersfield AE, and Holgate ST (1983) Inhaled adenosine and guanosine on airway resistance in normal and asthmatic subjects. *Br J Clin Pharmacol* **15**:161–165.
- Cutz E, Fu XW, Yeger H, Peers C, and Kemp PJ (2003) Oxygen sensing in pulmonary neuroepithelial bodies and related tumour cell models. In *Lung Oxygen Sensing* (Lahiri S, Semenza GL and Prabhakar NR eds) pp 567–602, Marcel Dekker, New York.
- Dagnelie PC and Agteresch HJ (2004) Promising effects of adenosine triphosphate infusion on nutritional status and quality of life in advanced non-small-cell lung cancer: a randomized clinical trial. *Drug Dev Res* **59**:146–151.
- Darmani NA and Broadley KJ (1986) Actions and interactions of adenosine, theophylline and enprofylline on the guinea-pig spirally cut trachea. *Eur J Pharmacol* **125**:353–362.
- Davis CW, Dowell ML, Lethem M, and Van Scott M (1992) Goblet cell degranulation in isolated canine tracheal epithelium: response to exogenous ATP, ADP, and adenosine. *Am J Physiol* **262**:C1313–C1323.
- Davis CW and Lazarowski E (2008) Coupling of airway ciliary activity and mucin secretion to mechanical stresses by purinergic signaling. *Respir Physiol Neurobiol* **163**:208–213.
- Davis IC, Sullender WM, Hickman-Davis JM, Lindsey JR, and Matalon S (2004) Nucleotide-mediated inhibition of alveolar fluid clearance in BALB/c mice after respiratory syncytial virus infection. *Am J Physiol Lung Cell Mol Physiol* **286**:L112–L120.
- Dawicki DD, McGowan-Jordan J, Bullard S, Pond S, and Rounds S (1995) Extracellular nucleotides stimulate leukocyte adherence to cultured pulmonary artery endothelial cells. *Am J Physiol* **268**:L666–L673.
- Dawson JM, Cook ND, Coade SB, Baum H, and Peters TJ (1986) Demonstration of plasma-membrane adenosine diphosphatase activity in rat lung. *Biochim Biophys Acta* **856**:566–570.
- De Proost I, Pintelon I, Wilkinson WJ, Goethals S, Brouns I, Van Nassauw L, Riccardi D, Timmermans JP, Kemp PJ, and Adriaenssen D (2009) Purinergic signaling in the pulmonary neuroepithelial body microenvironment unraveled by live cell imaging. *FASEB J* **23**:1153–1160.
- de Souza Alvares T, Huxtable A, Ireland M, Adachi T, Poon B, Lorier A, and Funk G (2007) P2Y receptor modulation of XII inspiratory motor output in neonatal rat. *FASEB J* **21**:A1295–A1296.
- Dellabianca A, Faniglione M, De Angelis S, Tonini S, Balestra B, Colucci M, Cervio M, Clavanzani P, Chiochetti R, De Giorgio R, et al. (2009) Adenosine A₁ and A₃ receptor agonists inhibit nonadrenergic, noncholinergic relaxations in the guinea pig isolated trachea. *Respiration* **78**:75–83.
- Denlinger LC, Shi L, Guadarrama A, Schell K, Green D, Morrin A, Hogan K, Sorkness RL, Busse WW, and Gern JE (2009) Attenuated P2X₇ pore function as a risk factor for virus-induced loss of asthma control. *Am J Respir Crit Care Med* **179**:265–270.
- Diamond L and O'Donnell M (1980) A nonadrenergic vagal inhibitory pathway to feline airways. *Science* **208**:185–188.
- Dieterle Y, Ody C, Ehrensberger A, Stalder H, and Junod AF (1978) Metabolism and uptake of adenosine triphosphate and adenosine by porcine aortic and pulmonary endothelial cells and fibroblasts in culture. *Circ Res* **42**:869–876.
- Dietl P, Haller T, Wirlreitner B, Völk H, Friedrich F, and Striessnig J (1995)

- Activation of L-type Ca^{2+} channels after purinoceptor stimulation by ATP in an alveolar epithelial cell (L2). *Am J Physiol* **269**:L873–L883.
- Dietl P, Liss B, Felder E, Miklavc P, and Wirtz H (2010) Lamellar body exocytosis by cell stretch or purinergic stimulation: possible physiological roles, messengers and mechanisms. *Cell Physiol Biochem* **25**:1–12.
- Dimakou K, Hillas G, and Bakakos P (2009) Adenosine deaminase activity and its isoenzymes in the sputum of patients with pulmonary tuberculosis. *Int J Tuberc Lung Dis* **13**:744–748.
- Dinh Xuan AT, Higenbottam TW, Clelland C, Pepke-Zaba J, Wells FC, and Wallwork J (1990) Acetylcholine and adenosine diphosphate cause endothelium-dependent relaxation of isolated human pulmonary arteries. *Eur Respir J* **3**:633–638.
- Donaldson SH, Picher M, and Boucher RC (2002) Secreted and cell-associated adenylate kinase and nucleoside diphosphokinase contribute to extracellular nucleotide metabolism on human airway surfaces. *Am J Respir Cell Mol Biol* **26**:209–215.
- Douillet CD, Robinson WP 3rd, Milano PM, Boucher RC, and Rich PB (2006) Nucleotides induce IL-6 release from human airway epithelia via P2Y2 and p38 MAPK-dependent pathways. *Am J Physiol Lung Cell Mol Physiol* **291**:L734–L746.
- Drury AN and Szent-Györgyi A (1929) The physiological activity of adenine compounds with special reference to their action upon the mammalian heart. *J Physiol* **68**:213–237.
- Duffy SM, Cruse G, Brightling CE, and Bradding P (2007) Adenosine closes the K^{+} channel KCa3.1 in human lung mast cells and inhibits their migration via the adenosine A_{2A} receptor. *Eur J Immunol* **37**:1653–1662.
- Dulong S, Bernard K, and Ehrenfeld J (2007) Enhancement of P2Y₆-induced Cl^{-} secretion by IL-13 and modulation of SK4 channels activity in human bronchial cells. *Cell Physiol Biochem* **20**:483–494.
- Eckle T, Grenz A, Laucher S, and Eltzhchig HK (2008) A2B adenosine receptor signaling attenuates acute lung injury by enhancing alveolar fluid clearance in mice. *J Clin Invest* **118**:3301–3315.
- Eckle T, Koeppen M, and Eltzhchig HK (2009) Role of extracellular adenosine in acute lung injury. *Physiology (Bethesda)* **24**:298–306.
- el-Hashim A, D'Agostino B, Matera MG, and Page C (1996) Characterization of adenosine receptors involved in adenosine-induced bronchoconstriction in allergic rabbits. *Br J Pharmacol* **119**:1262–1268.
- El-Kashef H, Elmazar MM, Al-Shabanah OA, and Al-Bekairi AM (1999) Effect of adenosine on pulmonary circulation of rabbits. *Gen Pharmacol* **32**:307–313.
- Eldridge FL and Millhorn DE (1987) Role of adenosine in the regulation of breathing, in *Topics and Perspectives in Adenosine Research* (Gerlach E and Becker BF eds) pp 586–596. Springer-Verlag, Berlin.
- Elzein E, Kalla RV, Li X, Perry T, Gimbel A, Zeng D, Lustig D, Leung K, and Zablocki J (2008) Discovery of a novel A2B adenosine receptor antagonist as a clinical candidate for chronic inflammatory airway diseases. *J Med Chem* **51**:2267–2278.
- Esther CR Jr, Alexis NE, Clas ML, Lazarowski ER, Donaldson SH, Ribeiro CM, Moore CG, Davis SD, and Boucher RC (2008) Extracellular purines are biomarkers of neutrophilic airway inflammation. *Eur Respir J* **31**:949–956.
- Esther CR Jr, Peden DB, Alexis NE, and Hernandez ML (2011) Airway purinergic responses in healthy, atopic nonasthmatic, and atopic asthmatic subjects exposed to ozone. *Inhal Toxicol* **23**:324–330.
- Ethier MF and Madison JM (2006) Adenosine A_1 receptors mediate mobilization of calcium in human bronchial smooth muscle cells. *Am J Respir Cell Mol Biol* **35**:496–502.
- Evaldsson C, Rydén I, Rosén A, and Uppugunduri S (2009) 4-thiouridine induces dose-dependent reduction of oedema, leucocyte influx and tumour necrosis factor in lung inflammation. *Clin Exp Immunol* **155**:330–338.
- Evans JH and Sanderson MJ (1999) Intracellular calcium oscillations induced by ATP in airway epithelial cells. *Am J Physiol* **277**:L30–L41.
- Factor P, Mutlu GM, Chen L, Mohamed J, Akhmedov AT, Meng FJ, Jilling T, Lewis ER, Johnson MD, Xu A, et al. (2007) Adenosine regulation of alveolar fluid clearance. *Proc Natl Acad Sci USA* **104**:4083–4088.
- Faffe DS, Costa-Vieira J, Mazzoli-Rocha F, Cagido VR, Moncao-Ribeiro LC, Takiya CM, Coutinho-Silva R, and Zin WA (2009) Role of purinergic P2X₇ receptors in allergic inflammation. *Am J Respir Crit Care Med* **179**:A4255.
- Fan M and Mustafa SJ (2002) Adenosine-mediated bronchoconstriction and lung inflammation in an allergic mouse model. *Pulm Pharmacol Ther* **15**:147–155.
- Fan M, Qin W, and Mustafa SJ (2003) Characterization of adenosine receptor(s) involved in adenosine-induced bronchoconstriction in an allergic mouse model. *Am J Physiol Lung Cell Mol Physiol* **284**:L1012–L1019.
- Farmer JB and Farrar DG (1976) Pharmacological studies with adenine, adenosine and some phosphorylated derivatives on guinea-pig tracheal muscle. *J Pharm Pharmacol* **28**:748–752.
- Fausther M, Pelletier J, Ribeiro CM, Sévigny J, and Picher M (2010) Cystic fibrosis remodels the regulation of purinergic signaling by NTPDase1 (CD39) and NTP-Dase3. *Am J Physiol Lung Cell Mol Physiol* **298**:L804–L818.
- Fedan JS, Belt JJ, Yuan LX, and Frazer DG (1993a) Contractile effects of nucleotides in guinea pig isolated, perfused trachea: involvement of respiratory epithelium, prostanoids and Na^{+} and Cl^{-} channels. *J Pharmacol Exp Ther* **264**:210–216.
- Fedan JS, Belt JJ, Yuan LX, and Frazer DG (1993b) Relaxant effects of nucleotides in guinea pig isolated, perfused trachea: lack of involvement of prostanoids, Cl^{-} channels and adenosine. *J Pharmacol Exp Ther* **264**:217–220.
- Feoktistov I and Biaggioni I (1996) Role of adenosine in asthma. *Drug Dev Res* **39**:333–336.
- Feoktistov I, Goldstein A, Sheller JR, Schwartz LB, and Biaggioni I (2003) Immunological characterization of A_{2B} adenosine receptors in human mast cells. *Drug Dev Res* **58**:461–471.
- Finney MJ, Karlsson JA, and Persson CG (1985) Effects of bronchoconstrictors and bronchodilators on a novel human small airway preparation. *Br J Pharmacol* **85**:29–36.
- Forsythe P and Ennis M (1999) Adenosine, mast cells and asthma. *Inflamm Res* **48**:301–307.
- Forteza R, Salathe M, Miot F, Forteza R, and Conner GE (2005) Regulated hydrogen peroxide production by Duox in human airway epithelial cells. *Am J Respir Cell Mol Biol* **32**:462–469.
- Fortner CN, Breyer RM, and Paul RJ (2001) EP₂ receptors mediate airway relaxation to substance P, ATP, and PGE₂. *Am J Physiol Lung Cell Mol Physiol* **281**:L469–L474.
- Fozard JR (2003) The case for a role for adenosine in asthma: almost convincing? *Curr Opin Pharmacol* **3**:264–269.
- Fozard JR, Ellis KM, Vilella Dantas MF, Tigani B, and Mazzoni L (2002) Effects of CGS 21680, a selective adenosine A_{2A} receptor agonist, on allergic airways inflammation in the rat. *Eur J Pharmacol* **438**:183–188.
- Fozard JR and Hannon JP (2000) Species differences in adenosine receptor-mediated bronchoconstrictor responses. *Clin Exp Allergy* **30**:1213–1220.
- Fozard JR and McCarthy C (2002) Adenosine receptor ligands as potential therapeutics in asthma. *Curr Opin Investig Drugs* **3**:69–77.
- Freem LJ, Escot S, Tannahill D, Druckenbrod NR, Thapar N, and Burns AJ (2010) The intrinsic innervation of the lung is derived from neural crest cells as shown by optical projection tomography in *Wnt1-Cre;YFP* reporter mice. *J Anat* **217**:651–664.
- Fricke E, Esdorn E, Kammeier A, Fricke H, Preuss R, Burchert W, and Lindner O (2008) Respiratory resistance of patients during cardiac stress testing with adenosine: is dyspnea a sign of bronchospasm? *J Nucl Cardiol* **15**:94–99.
- Fu XW, Nurse CA, and Cutz E (2004) Expression of functional purinergic receptors in pulmonary neuroepithelial bodies and their role in hypoxia chemotransmission. *Biol Chem* **385**:275–284.
- Fu Z, Bettge K, Carroll S, Buchholz KR, and Machen TE (2007) Role of Ca^{2+} in responses of airway epithelia to *Pseudomonas aeruginosa*, flagellin, ATP, and thapsigargin. *Am J Physiol Lung Cell Mol Physiol* **292**:L353–L364.
- Funk GD, Huxtable AG, and Lorier AR (2008) ATP in central respiratory control: a three-part signaling system. *Respir Physiol Neurobiol* **164**:131–142.
- Funk GD, Kanjhan R, Walsh C, Lipski J, Comer AM, Parkis MA, and Housley GD (1997) P2 receptor excitation of rodent hypoglossal motoneuron activity in vitro and in vivo: a molecular physiological analysis. *J Neurosci* **17**:6325–6337.
- Gaba SJ, Bourguoin-Karaoui D, Dujols P, Michel FB, and Préfaut C (1986) Effects of adenosine triphosphate on pulmonary circulation in chronic obstructive pulmonary disease. ATP: a pulmonary vasoregulator? *Am Rev Respir Dis* **134**:1140–1144.
- Gaba SJ and Préfaut C (1990) Comparison of pulmonary and systemic effects of adenosine triphosphate in chronic obstructive pulmonary disease—ATP: a pulmonary controlled vasoregulator? *Eur Respir J* **3**:450–455.
- Gallagher CJ and Salter MW (2003) Differential properties of astrocyte calcium waves mediated by P2Y1 and P2Y2 receptors. *J Neurosci* **23**:6728–6739.
- Gayle S and Burnstock G (2005) Immunolocalisation of P2X and P2Y nucleotide receptors in the rat nasal mucosa. *Cell Tissue Res* **319**:27–36.
- Gerasimovskaya EV, Woodward HN, Tucker DA, and Stenmark KR (2008) Extracellular ATP is a pro-angiogenic factor for pulmonary artery vasa vasorum endothelial cells. *Angiogenesis* **11**:169–182.
- Ghai G, Zimmerman MB, and Hopkins MF (1987) Evidence for A_1 and A_2 adenosine receptors in guinea pig trachea. *Life Sci* **41**:1215–1224.
- Gilfillan AM and Rooney SA (1987) Purinoceptor agonists stimulate phosphatidylcholine secretion in primary cultures of adult rat type II pneumocytes. *Biochim Biophys Acta* **917**:18–23.
- Gobran LI and Rooney SA (1990) Adenosine A_1 receptor-mediated inhibition of surfactant secretion in rat type II pneumocytes. *Am J Physiol* **258**:L45–L51.
- Goralski JL, Boucher RC, and Button B (2010) Osmolytes and ion transport modulators: new strategies for airway surface rehydration. *Curr Opin Pharmacol* **10**:294–299.
- Gottlieb JE, Peake MD, and Sylvester JT (1984) Adenosine and hypoxic pulmonary vasodilation. *Am J Physiol* **247**:H541–H547.
- Gourine AV (2005) On the peripheral and central chemoreception and control of breathing: an emerging role of ATP. *J Physiol* **568**:715–724.
- Gourine AV, Atkinson L, Deuchars J, and Spyer KM (2003) Purinergic signalling in the medullary mechanisms of respiratory control in the rat: respiratory neurones express the P2X2 receptor subunit. *J Physiol* **552**:197–211.
- Gourine AV, Dale N, and Spyer KM (2004) Chemosensory control of the respiratory function. Working towards understanding the role of ATP-mediated purinergic signalling. *Adv Exp Med Biol* **551**:31–38.
- Gourine AV, Kasymov V, Marina N, Tang F, Figueiredo MF, Lane S, Teschemacher AG, Spyer KM, Deisseroth K, and Kasparov S (2010) Astrocytes control breathing through pH-dependent release of ATP. *Science* **329**:571–575.
- Gourine AV, Llaudet E, Dale N, and Spyer KM (2005) ATP is a mediator of chemosensory transduction in the central nervous system. *Nature* **436**:108–111.
- Gourine AV, Llaudet E, Thomas T, Dale N, and Spyer KM (2002) Adenosine release in nucleus tractus solitarius does not appear to mediate hypoxia-induced respiratory depression in rats. *J Physiol* **544**:161–170.
- Gourine AV and Spyer KM (2003) Chemosensitivity of medullary respiratory neurones. A role for ionotropic P2X and GABA_A receptors, in *Chemosensation* (Gonzalez G, Prabhakar JM, Pequignot JM, Nurse CA and Dalmaz Y eds) pp 375–387. Kluwer Academic/Plenum Publishers, New York.
- Govindaraju V, Martin JG, Maghni K, Ferraro P, and Michoud MC (2005) The effects of extracellular purines and pyrimidines on human airway smooth muscle cells. *J Pharmacol Exp Ther* **315**:941–948.
- Grantham CJ and Bakhle YS (1988) Effect of acute lung injury on metabolism of adenine nucleotides in rat perfused lung. *Br J Pharmacol* **94**:1029–1036.
- Greenberg B, Rhoden K, and Barnes PJ (1987) Endothelium-dependent relaxation of human pulmonary arteries. *Am J Physiol* **252**:H434–H438.
- Greenberg SS, Zhao X, Wang JF, Hua L, and Ouyang J (1997) cAMP and purinergic P_{2y} receptors upregulate and enhance inducible NO synthase mRNA and protein in vivo. *Am J Physiol* **273**:L967–L979.
- Griese M, Gobran LI, and Rooney SA (1991) A_2 and P₂ purine receptor interactions and surfactant secretion in primary cultures of type II cells. *Am J Physiol* **261**:L140–L147.

- Gross CH, Abdul-Manan N, Fulghum J, Lippke J, Liu X, Prabhakar P, Brennan D, Willis MS, Faerman C, Connelly P, et al (2006) Nucleotide-binding domains of cystic fibrosis transmembrane conductance regulator, an ABC transporter, catalyze adenylate kinase activity but not ATP hydrolysis. *J Biol Chem* **281**:4058–4068.
- Gu Q, Ruan T, Hong JL, Burki N, and Lee LY (2003) Hypersensitivity of pulmonary C fibers induced by adenosine in anesthetized rats. *J Appl Physiol* **95**:1315–1324.
- Gu YY, Zhang HY, Zhang HJ, Li SY, Ni JH, and Jia HT (2006) 8-Chloro-adenosine inhibits growth at least partly by interfering with actin polymerization in cultured human lung cancer cells. *Biochem Pharmacol* **72**:541–550.
- Gui Y, Walsh MP, Jankowski V, Jankowski J, and Zheng XL (2008) U_p4A stimulates endothelium-independent contraction of isolated rat pulmonary artery. *Am J Physiol Lung Cell Mol Physiol* **294**:L733–L738.
- Guibert C, Pacaud P, Loirand G, Marthan R, and Savineau JP (1996) Effect of extracellular ATP on cytosolic Ca^{2+} concentration in rat pulmonary artery myocytes. *Am J Physiol* **271**:L450–L458.
- Gundersen KL and Kopito RR (1994) Effects of pyrophosphate and nucleotide analogs suggest a role for ATP hydrolysis in cystic fibrosis transmembrane regulator channel gating. *J Biol Chem* **269**:19349–19353.
- Guyot A and Hanrahan JW (2002) ATP release from human airway epithelial cells studied using a capillary cell culture system. *J Physiol* **545**:199–206.
- Hachiya H, Hirao K, Takahashi A, Nagata Y, Suzuki K, Maeda S, Sasaki T, Kawabata M, Isobe M, and Iesaka Y (2007) Clinical implications of reconnection between the left atrium and isolated pulmonary veins provoked by adenosine triphosphate after extensive encircling pulmonary vein isolation. *J Cardiovasc Electrophysiol* **18**:392–398.
- Hannon JP, Tigani B, Williams I, Mazzoni L, and Fozard JR (2001) Mechanism of airway hyperresponsiveness to adenosine induced by allergen challenge in actively sensitized Brown Norway rats. *Br J Pharmacol* **132**:1509–1523.
- Hansen M, Boitano S, Dirksen ER, and Sanderson MJ (1993) Intercellular calcium signaling induced by extracellular adenosine 5'-triphosphate and mechanical stimulation in airway epithelial cells. *J Cell Sci* **106**:995–1004.
- Hargitai D, Pataki A, Raffai G, Füzi M, Dankó T, Csernok L, Várnai P, Szigeti GP, and Zsembery A (2010) Calcium entry is regulated by Zn^{2+} in relation to extracellular ionic environment in human airway epithelial cells. *Respir Physiol Neurobiol* **170**:67–75.
- Hartley SA, Kato K, Salter KJ, and Kozłowski RZ (1998) Functional evidence for a novel suramin-insensitive pyrimidine receptor in rat small pulmonary arteries. *Circ Res* **83**:940–946.
- Hartley SA and Kozłowski RZ (1997) Electrophysiological consequences of purinergic receptor stimulation in isolated rat pulmonary arterial myocytes. *Circ Res* **80**:170–178.
- Haskell CM, Mendoza E, Pisters KM, Fossella FV, and Figlin RA (1998) Phase II study of intravenous adenosine 5'-triphosphate in patients with previously untreated stage IIIB and stage IV non-small cell lung cancer. *Invest New Drugs* **16**:81–85.
- Haskó G, Xu DZ, Lu Q, Németh ZH, Jabush J, Berezina TL, Zaets SB, Csóka B, and Deitch EA (2006) Adenosine A2A receptor activation reduces lung injury in trauma/hemorrhagic shock. *Crit Care Med* **34**:1119–1125.
- Hasséssian H, Bodin P, and Burnstock G (1993) Blockade by glibenclamide of the flow-induced endothelial release of ATP that contributes to vasodilatation in the pulmonary vascular bed of the rat. *Br J Pharmacol* **109**:466–472.
- Hasséssian H and Burnstock G (1995) Interacting roles of nitric oxide and ATP in the pulmonary circulation of the rat. *Br J Pharmacol* **114**:846–850.
- Hatta Y, Takahashi M, Enomoto Y, Takahashi N, Sawada U, and Horie T (2004) Adenosine triphosphate (ATP) enhances the antitumor effect of etoposide (VP16) in lung cancer cells. *Oncol Rep* **12**:1139–1142.
- Hedner T, Hedner J, Wessberg P, and Jonason J (1982) Regulation of breathing in the rat: indications for a role of central adenosine mechanisms. *Neurosci Lett* **33**:147–151.
- Heller AR, Rothelmer J, Weigand MA, Plaschke K, Schmeck J, Wendel M, Bardenheuer HJ, and Koch T (2007) Adenosine A1 and A2 receptor agonists reduce endotoxin-induced cellular energy depletion and oedema formation in the lung. *Eur J Anaesthesiol* **24**:258–266.
- Hellewell PG and Pearson JD (1984) Purinoceptor mediated stimulation of prostacyclin release in the porcine pulmonary vasculature. *Br J Pharmacol* **83**:457–462.
- Herlenius E, Adén U, Tang LQ, and Lagercrantz H (2002) Perinatal respiratory control and its modulation by adenosine and caffeine in the rat. *Pediatr Res* **51**:4–12.
- Hinshaw LB, Kuida H, Gilbert RP, and Visscher MB (1957) Influence of perfusate characteristics on pulmonary vascular response to endotoxin. *Am J Physiol* **191**:293–295.
- Holgate ST (2002) Adenosine: a key effector molecule of asthma or just another mediator? *Am J Physiol Lung Cell Mol Physiol* **282**:L167–L168.
- Holgate ST (2005) The Quintiles Prize Lecture 2004. The identification of the adenosine A_{2B} receptor as a novel therapeutic target in asthma. *Br J Pharmacol* **145**:1009–1015.
- Holgate ST, Finnerty JP, and Polosa R (1990) Mechanisms of purine-induced bronchoconstriction in asthma. *Arch Int Pharmacodyn Ther* **303**:122–131.
- Holgate ST, Mann JS, Church MK, and Cusley MJ (1987) Mechanisms and significance of adenosine-induced bronchoconstriction in asthma. *Allergy* **42**:481–484.
- Homolya L, Steinberg TH, and Boucher RC (2000) Cell to cell communication in response to mechanical stress via bilateral release of ATP and UTP in polarized epithelia. *J Cell Biol* **150**:1349–1360.
- Horckmans M, Marcet B, Marteau F, Bulté F, Maho A, Parmentier M, Boeynaems JM, and Communi D (2006) Extracellular adenine nucleotides inhibit the release of major monocyte recruiters by human monocyte-derived dendritic cells. *FEBS Lett* **580**:747–754.
- Hua X, Chason KD, Fredholm BB, Deshpande DA, Penn RB, and Tilley SL (2008) Adenosine induces airway hyperresponsiveness through activation of A_3 receptors on mast cells. *J Allergy Clin Immunol* **122**:107–113, 113.e1–e7.
- Hua X, Erikson CJ, Chason KD, Rosebrock CN, Deshpande DA, Penn RB, and Tilley SL (2007) Involvement of A_1 adenosine receptors and neural pathways in adenosine-induced bronchoconstriction in mice. *Am J Physiol Lung Cell Mol Physiol* **293**:L25–L32.
- Huang P, Lazarowski ER, Tarran R, Milgram SL, Boucher RC, and Stutts MJ (2001) Compartmentalized autocrine signaling to cystic fibrosis transmembrane conductance regulator at the apical membrane of airway epithelial cells. *Proc Natl Acad Sci USA* **98**:14120–14125.
- Huckstepp RT, id Bihi R, Eason R, Spyer KM, Dicke N, Willecke K, Marina N, Gourine AV, and Dale N (2010) Connexin hemichannel-mediated CO_2 -dependent release of ATP in the medulla oblongata contributes to central respiratory chemosensitivity. *J Physiol* **588**:3901–3920.
- Hughes PJ, Holgate ST, and Church MK (1984) Adenosine inhibits and potentiates IgE-dependent histamine release from human lung mast cells by an A_2 -purinoceptor mediated mechanism. *Biochem Pharmacol* **33**:3847–3852.
- Husted SE and Nedergaard OA (1985) Dual inhibitory action of ATP on adrenergic neuroeffector transmission in rabbit pulmonary artery. *Acta Pharmacol Toxicol (Copenh)* **57**:204–213.
- Huszár E, Vass G, Vizi E, Csoma Z, Barát E, Molnár Világos G, Herjavec I, and Horváth I (2002) Adenosine in exhaled breath condensate in healthy volunteers and in patients with asthma. *Eur Respir J* **20**:1393–1398.
- Huxtable A, Parmar M, Poon B, Lorier A, Pagliardini S, Greer J, and Funk G (2007) Purinergic modulation of preBöttinger (PBC) rhythm generating networks during perinatal development (Abstract). *FASEB J* **21**:A1296.
- Huxtable AG, Zwicker JD, Alvares TS, Ruangkittisakul A, Fang X, Hahn LB, Posse de Chaves E, Baker GB, Ballanyi K, and Funk GD (2010) Glia contribute to the purinergic modulation of inspiratory rhythm-generating networks. *J Neurosci* **30**:3947–3958.
- Huxtable AG, Zwicker JD, Poon BY, Pagliardini S, Vrouwe SQ, Greer JJ, and Funk GD (2009) Tripartite purinergic modulation of central respiratory networks during perinatal development: the influence of ATP, ectonucleotidases, and ATP metabolites. *J Neurosci* **29**:14713–14725.
- Hwang TC, Nagel G, Nairn AC, and Gadsby DC (1994) Regulation of the gating of cystic fibrosis transmembrane conductance regulator C1 channels by phosphorylation and ATP hydrolysis. *Proc Natl Acad Sci USA* **91**:4698–4702.
- Hwang TH, Schwiebert EM, and Guggino WB (1996) Apical and basolateral ATP stimulates tracheal epithelial chloride secretion via multiple purinergic receptors. *Am J Physiol* **270**:C1611–C1623.
- Ildzko M, Hammad H, van Nimwegen M, Kool M, Willart MA, Muskens F, Hoogsteden HC, Luttmann W, Ferrari D, Di Virgilio F, et al. (2007) Extracellular ATP triggers and maintains asthmatic airway inflammation by activating dendritic cells. *Nat Med* **13**:913–919.
- Inbar S, Schrader BJ, Kaufmann E, Vestal RE, and Rich S (1993) Effects of adenosine in combination with calcium channel blockers in patients with primary pulmonary hypertension. *J Am Coll Cardiol* **21**:413–418.
- Inglis SK, Collett A, McAlroy HL, Wilson SM, and Olver RE (1999) Effect of luminal nucleotides on Cl^- secretion and Na^+ absorption in distal bronchi. *Pflugers Arch* **438**:621–627.
- Inoue T and Kannan MS (1988) Nonadrenergic and noncholinergic excitatory neurotransmission in rat intrapulmonary artery. *Am J Physiol* **254**:H1142–H1148.
- Irvin C, Martin R, and Macklem PT (1983) Non-adrenergic inhibitory nerves. *Eur J Respir Dis Suppl* **128**:133–138.
- Irvin CG, Boileau R, Tremblay J, Martin RR, and Macklem PT (1980) Bronchodilation: noncholinergic, nonadrenergic mediation demonstrated in vivo in the cat. *Science* **207**:791–792.
- Jankowski V, Tölle M, Vanholder R, Schönfelder G, van der Giet M, Henning L, Schlüter H, Paul M, Zidek W, and Jankowski J (2005) Uridine adenosine tetraphosphate: a novel endothelium-derived vasoconstrictive factor. *Nat Med* **11**:223–227.
- Johnson HG and McNeer ML (1985) Adenosine-induced secretion in the canine trachea: modification by methylxanthines and adenosine derivatives. *Br J Pharmacol* **86**:63–67.
- Jones TR, Lefcoe NM, and Hamilton JT (1980) Pharmacological study of adenosine and related compounds on isolated guinea pig trachea: evidence for more than one type of purine receptor. *Can J Physiol Pharmacol* **58**:1356–1365.
- Junger WG, Chen Y, Corriden R, Yip L, and Hashiguchi N (2006) ATP and adenosine control PMN chemotaxis via autocrine feedback mechanisms—role in shock (Abstract). *Shock* **25**:27.
- Kalenberg S and Satchell DG (1979) The inhibitory innervation of the guinea-pig trachea: a study of its adrenergic and non-adrenergic components. *Clin Exp Pharmacol Physiol* **6**:549–559.
- Kamei J and Takahashi Y (2006) Involvement of ionotropic purinergic receptors in the histamine-induced enhancement of the cough reflex sensitivity in guinea pigs. *Eur J Pharmacol* **547**:160–164.
- Kamikawa Y and Shimo Y (1976) Pharmacological differences of non-adrenergic inhibitory response and of ATP-induced relaxation in guinea-pig tracheal strips. *J Pharm Pharmacol* **28**:854–855.
- Kamikawa Y and Shimo Y (1989) Adenosine selectively inhibits noncholinergic transmission in guinea pig bronchi. *J Appl Physiol* **66**:2084–2091.
- Kanazawa H, Fujiwara H, Shoji S, Fujii T, Kudoh S, Hirata K, Kurihara N, and Yoshikawa J (1997) Adenosine modulates endothelin-induced bronchoconstriction in guinea pig airway. *Int Arch Allergy Immunol* **112**:83–87.
- Karlsson JA and Persson CG (1984) Neither vasoactive intestinal peptide (VIP) nor purine derivatives may mediate non-adrenergic tracheal inhibition. *Acta Physiol Scand* **122**:589–598.
- Katchanov G, Xu J, Schulman ES, and Pelleg A (1998) ATP causes neurogenic bronchoconstriction in the dog. *Drug Dev Res* **45**:342–349.
- Katsuragi T and Su C (1982) Release of purines and noradrenaline by ouabain and potassium chloride from vascular adrenergic nerves. *Br J Pharmacol* **77**:625–629.
- Kaunitz JD and Akiba Y (2009) Regulation of extracellular pH by purinergic signaling. *Physiol News* **77**:22–24.

- Kawakami M, Nagira T, Hayashi T, Shimamoto C, Kubota T, Mori H, Yoshida H, and Nakahari T (2004) Hypo-osmotic potentiation of acetylcholine-stimulated ciliary beat frequency through ATP release in rat tracheal ciliary cells. *Exp Physiol* **89**:739–751.
- Kawasaki H, Cline WH, Jr, and Su C (1983) Possible roles of purinergic modulation in pathogenesis of some diseases: hypertension and asthma, in *Physiology and Pharmacology of Adenosine Derivatives* (Daly JW, Kuroda Y, Phillis JW, Shimizu H and U M eds) pp 189–196, Raven Press, New York.
- Keir S, Boswell-Smith V, Spina D, and Page C (2006) Mechanism of adenosine-induced airways obstruction in allergic guinea pigs. *Br J Pharmacol* **147**:720–728.
- Kellerman D, Evans R, Mathews D, and Shaffer C (2002) Inhaled P2Y₂ receptor agonists as a treatment for patients with Cystic Fibrosis lung disease. *Adv Drug Deliv Rev* **54**:1463–1474.
- Kemp PA, Sugar RA, and Jackson AD (2004) Nucleotide-mediated mucin secretion from differentiated human bronchial epithelial cells. *Am J Respir Cell Mol Biol* **31**:446–455.
- Kessler S, Clauss WG, Günther A, Kummer W, and Fronius M (2011) Expression and functional characterization of P2X receptors in mouse alveolar macrophages. *Pflugers Arch* **462**:419–430.
- Kieffmann R, Islam MN, Lindert J, Parthasarathi K, and Bhattacharya J (2009) Paracrine purinergic signaling determines lung endothelial nitric oxide production. *Am J Physiol Lung Cell Mol Physiol* **296**:L901–L910.
- Kim CH, Kim SS, Choi JY, Shin JH, Kim JY, Namkung W, Lee JG, Lee MG, and Yoon JH (2004) Membrane-specific expression of functional purinergic receptors in normal human nasal epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **287**:L835–L842.
- Kim KC and Lee BC (1991) P₂ purinoceptor regulation of mucin release by airway goblet cells in primary culture. *Br J Pharmacol* **103**:1053–1056.
- Kim KC, Park HR, Shin CY, Akiyama T, and Ko KH (1996) Nucleotide-induced mucin release from primary hamster tracheal surface epithelial cells involves the P_{2U} purinoceptor. *Eur Respir J* **9**:1579.
- Kim SH, Kim YK, Park HW, Kim SH, Kim SH, Ye YM, Min KU, and Park HS (2009) Adenosine deaminase and adenosine receptor polymorphisms in aspirin-intolerant asthma. *Respir Med* **103**:356–363.
- Kim SJ, Min HY, Chung HJ, Park EJ, Hong JY, Kang YJ, Shin DH, Jeong LS, and Lee SK (2008) Inhibition of cell proliferation through cell cycle arrest and apoptosis by thio-Cl-IB-MECA, a novel A₃ adenosine receptor agonist, in human lung cancer cells. *Cancer Lett* **264**:309–315.
- Kishore BK, Ginns SM, Krane CM, Nielsen S, and Knepper MA (2000) Cellular localization of P2Y₂ purinoceptor in rat renal inner medulla and lung. *Am J Physiol Renal Physiol* **278**:F43–F51.
- Knowles MR, Clarke LL, and Boucher RC (1991) Activation by extracellular nucleotides of chloride secretion in the airway epithelia of patients with cystic fibrosis. *N Engl J Med* **325**:533–538.
- Ko YH, Delannoy M, and Pedersen PL (1997) Cystic fibrosis transmembrane conductance regulator: the first nucleotide binding fold targets the membrane with retention of its ATP binding function. *Biochemistry* **36**:5053–5064.
- Koelle GB (1975) Cholinergic and non-cholinergic pharmacology of the bronchioles. *Postgrad Med J* **51**:63–68.
- Kollarik M, Dinh QT, Fischer A, and Undem BJ (2003) Capsaicin-sensitive and -insensitive vagal bronchopulmonary C-fibres in the mouse. *J Physiol* **551**:869–879.
- Kolosova IA, Mirzapozazova T, Moreno-Vinasco L, Sammani S, Garcia JG, and Verin AD (2008) Protective effect of purinergic agonist ATP γ S against acute lung injury. *Am J Physiol Lung Cell Mol Physiol* **294**:L319–L324.
- Kondo M, Kanoh S, Tamaoki J, Shirakawa H, Miyazaki S, and Nagai A (1998) Erythromycin inhibits ATP-induced intracellular calcium responses in bovine tracheal epithelial cells. *Am J Respir Cell Mol Biol* **19**:799–804.
- Konduri GG (1994) Systemic and myocardial effects of ATP and adenosine during hypoxic pulmonary hypertension in lambs. *Pediatr Res* **36**:41–48.
- Konduri GG, Bakhtashvili I, Frenn R, Chandrasekhar I, Jacobs ER, and Khanna AK (2004) P2Y purine receptor responses and expression in the pulmonary circulation of juvenile rabbits. *Am J Physiol Heart Circ Physiol* **287**:H157–H164.
- Konduri GG, Forman K, and Mital S (2000) Characterization of purine receptors in fetal lamb pulmonary circulation. *Pediatr Res* **47**:114–120.
- Konduri GG and Mattei J (2002) Role of oxidative phosphorylation and ATP release in mediating birth-related pulmonary vasodilation in fetal lambs. *Am J Physiol Heart Circ Physiol* **283**:H1600–H1608.
- Konduri GG and Mital S (2000) Adenosine and ATP cause nitric oxide-dependent pulmonary vasodilation in fetal lambs. *Biol Neonate* **78**:220–229.
- Konduri GG, Theodorou AA, Mukhopadhyay A, and Deshmukh DR (1992) Adenosine triphosphate and adenosine increase the pulmonary blood flow to postnatal levels in fetal lambs. *Pediatr Res* **31**:451–457.
- Koos BJ, Chau A, Matsuura M, Punla O, and Kruger L (2000) Thalamic lesions dissociate breathing inhibition by hypoxia and adenosine in fetal sheep. *Am J Physiol Regul Integr Comp Physiol* **278**:R831–R837.
- Kornerup KN, Page CP, and Moffatt JD (2005) Pharmacological characterisation of the adenosine receptor mediating increased ion transport in the mouse isolated trachea and the effect of allergen challenge. *Br J Pharmacol* **144**:1011–1016.
- Korngreen A, Ma W, Priel Z, and Silberberg SD (1998) Extracellular ATP directly gates a cation-selective channel in rabbit airway ciliated epithelial cells. *J Physiol* **508**:703–720.
- Korngreen A and Priel Z (1996) Purinergic stimulation of rabbit ciliated airway epithelia: control by multiple calcium sources. *J Physiol* **497**:53–66.
- Kotsis DH and Spence DM (2003) Detection of ATP-induced nitric oxide in a biomimetic circulatory vessel containing an immobilized endothelium. *Anal Chem* **75**:145–151.
- Kouzaki H, Iijima K, Kobayashi T, O'Grady SM, and Kita H (2011) The danger signal, extracellular ATP, is a sensor for an airborne allergen and triggers IL-33 release and innate Th2-type responses. *J Immunol* **186**:4375–4387.
- Kreda SM, Boucher RC, and Lazarowski ER (2008) Nucleotide release and airway epithelial physiology. *Physiol News* **71**:19–21.
- Kreda SM, Okada SF, van Heusden CA, O'Neal W, Gabriel S, Abdullah L, Davis CW, Boucher RC, and Lazarowski ER (2007) Coordinated release of nucleotides and mucin from human airway epithelial Calu-3 cells. *J Physiol* **584**:245–259.
- Kreda SM, Seminario-Vidal L, van Heusden CA, O'Neal W, Jones L, Boucher RC, and Lazarowski ER (2010) Receptor-promoted exocytosis of airway epithelial mucin granules containing a spectrum of adenine nucleotides. *J Physiol* **588**:2255–2267.
- Kunzelmann K, Bachhuber T, Reeger R, Markovich D, Sun J, and Schreiber R (2005) Purinergic inhibition of the epithelial Na⁺ transport via hydrolysis of PIP₂. *FASEB J* **19**:142–143.
- Kunzelmann K, Scheidt K, Scharf B, Ousingawat J, Schreiber R, Wainwright B, and McMorran B (2006) Flagellin of *Pseudomonas aeruginosa* inhibits Na⁺ transport in airway epithelia. *FASEB J* **20**:545–546.
- Lacroix JS, Ulman LG, and Potter EK (1994) The role of ATP in non-adrenergic sympathetic vascular control of the nasal mucosa in anaesthetized cats and dogs. *J Physiol* **476**:429–435.
- Laubinger W, Streubel G, and Reiser G (2001) Physiological evidence for a P2Y receptor responsive to diadenosine polyphosphates in human lung via Ca²⁺ release studies in bronchial epithelial cells. *Biochem Pharmacol* **61**:623–629.
- Laubinger W, Tulapurkar ME, Schäfer R, and Reiser G (2006) Distinct mono- and dinucleotide-specific P2Y receptors in A549 lung epithelial cells: different control of arachidonic acid release and nitric oxide synthase expression. *Eur J Pharmacol* **543**:1–7.
- Lázár Z, Huszár E, Kullmann T, Barta I, Antus B, Bikov A, Kollai M, and Horváth I (2008) Adenosine triphosphate in exhaled breath condensate of healthy subjects and patients with chronic obstructive pulmonary disease. *Inflamm Res* **57**:367–373.
- Lazarowski ER (2012) Vesicular and conductive mechanisms of nucleotide release. *Purinergic Signal* **8**:359–373.
- Lazarowski ER and Boucher RC (2009) Purinergic receptors in airway epithelia. *Curr Opin Pharmacol* **9**:262–267.
- Lazarowski ER, Boucher RC, and Harden TK (1994) Calcium-dependent release of arachidonic acid in response to purinergic receptor activation in airway epithelium. *Am J Physiol* **266**:C406–C415.
- Lazarowski ER, Paradiso AM, Watt WC, Harden TK, and Boucher RC (1997) UDP activates a mucosal-restricted receptor on human nasal epithelial cells that is distinct from the P2Y₂ receptor. *Proc Natl Acad Sci USA* **94**:2599–2603.
- Lazarowski ER, Tarran R, Grubb BR, van Heusden CA, Okada S, and Boucher RC (2004) Nucleotide release provides a mechanism for airway surface liquid homeostasis. *J Biol Chem* **279**:36855–36864.
- Lazarowski ER, Watt WC, Stutts MJ, Brown HA, Boucher RC, and Harden TK (1996) Enzymatic synthesis of UTP γ S, a potent hydrolysis resistant agonist of P_{2U}-purinoceptors. *Br J Pharmacol* **117**:203–209.
- Lee DK, Gray RD, and Lipworth BJ (2003) Adenosine monophosphate bronchial provocation and the actions of asthma therapy. *Clin Exp Allergy* **33**:287–294.
- Lee MG, Kollarik M, Chuaychoo B, and Undem BJ (2004) Ionotropic and metabotropic receptor mediated airway sensory nerve activation. *Pulm Pharmacol Ther* **17**:355–360.
- Lee SD, Nakano H, and Farkas GA (2005) Adenosinergic modulation of ventilation in obese Zucker rats. *Obes Res* **13**:545–555.
- Leij-Halfwerk S, Ageteresch HJ, Sijens PE, and Dagnelie PC (2002) Adenosine triphosphate infusion increases liver energy status in advanced lung cancer patients: an *in vivo* ³¹P magnetic resonance spectroscopy study. *Hepatology* **35**:421–424.
- Leipzig J (2003) Control of epithelial transport via luminal P2 receptors. *Am J Physiol Renal Physiol* **284**:F419–F432.
- Lemaire I and Leduc N (2004) Purinergic P2X₇ receptor function in lung alveolar macrophages: pharmacologic characterisation and bidirectional regulation by Th1 and Th2 cytokines. *Drug Dev Res* **59**:118–127.
- Li C, Ramjessingh M, and Bear CE (1996) Purified cystic fibrosis transmembrane conductance regulator (CFTR) does not function as an ATP channel. *J Biol Chem* **271**:11623–11626.
- Li WJ, Gu YY, Zhang HJ, Zhou J, and Jia HT (2009) Induction of p14ARF by E2F1 contributes to 8-chloro-adenosine-induced apoptosis in human lung cancer H1299 cells. *Chemotherapy* **55**:335–343.
- Li Y, Wang W, Parker W, and Clancy JP (2006) Adenosine regulation of cystic fibrosis transmembrane conductance regulator through prostenoids in airway epithelia. *Am J Respir Cell Mol Biol* **34**:600–608.
- Liang L, Zsembery A, and Schwiebert EM (2005) RNA interference targeted to multiple P2X receptor subtypes attenuates zinc-induced calcium entry. *Am J Physiol Cell Physiol* **289**:C388–C396.
- Lieb T, Frei CW, Frohock JL, Bookman RJ, and Salathe M (2002) Prolonged increase in ciliary beat frequency after short-term purinergic stimulation in human airway epithelial cells. *J Physiol* **538**:633–646.
- Liesker JJ, Ten Hacken NH, Rutgers SR, Zeinstra-Smith M, Postma DS, and Timens W (2007) Mast cell numbers in airway smooth muscle and PC20AMP in asthma and COPD. *Respir Med* **101**:882–887.
- Linnoila RI (2006) Functional facets of the pulmonary neuroendocrine system. *Lab Invest* **86**:425–444.
- Liu S and Barnes PJ (1994) Role of endothelium in the control of pulmonary vascular tone. *Endothelium* **2**:11–33.
- Liu SF, McCormack DG, Evans TW, and Barnes PJ (1989) Characterization and distribution of P₂-purinoceptor subtypes in rat pulmonary vessels. *J Pharmacol Exp Ther* **251**:1204–1210.
- Livingston M, Heaney LG, and Ennis M (2004) Adenosine, inflammation and asthma—a review. *Inflamm Res* **53**:171–178.
- Lommatzsch M, Cicko S, Müller T, Lucattelli M, Bratke K, Stoll P, Grimm M, Dürk T, Zissel G, Ferrari D, et al. (2010) Extracellular adenosine triphosphate and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* **181**:928–934.

- Lorier AR, Huxtable AG, Robinson DM, Lipski J, Housley GD, and Funk GD (2007) P2Y₁ receptor modulation of the pre-Bötzinger complex inspiratory rhythm generating network in vitro. *J Neurosci* **27**:993–1005.
- Lorier AR, Peebles K, Brosenitsch T, Robinson DM, Housley GD, and Funk GD (2004) P2 receptors modulate respiratory rhythm but do not contribute to central CO₂ sensitivity in vitro. *Respir Physiol Neurobiol* **142**:27–42.
- Losinski A and Alexander SP (1995) Adenosine receptor-mediated relaxation of guinea-pig precontracted, isolated trachea. *Br J Pharmacol* **116**:2425–2428.
- Lossos IS, Izibicki G, Or R, Goldstein RH, and Breuer R (2000) The effect of suramin on bleomycin-induced lung injury. *Life Sci* **67**:2873–2881.
- Lucas R, Verin AD, Black SM, and Catravas JD (2009) Regulators of endothelial and epithelial barrier integrity and function in acute lung injury. *Biochem Pharmacol* **77**:1763–1772.
- Lucattelli M, Cicko S, Müller T, Lommatzsch M, De Cunto G, Cardini S, Sundas W, Grimm M, Zeiser R, Dürk T, et al. (2011) P2X₇ receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol* **44**:423–429.
- Lunell E, Svedmyr N, Andersson KE, and Persson CG (1982) Effects of enprofylline, a xanthine lacking adenosine receptor antagonism, in patients with chronic obstructive lung disease. *Eur J Clin Pharmacol* **22**:395–402.
- Lunell E, Svedmyr N, Andersson KE, and Persson CG (1983) A novel bronchodilator xanthine apparently without adenosine receptor antagonism and tremorogenic effect. *Eur J Respir Dis* **64**:333–339.
- Lustig KD, Erb L, Landis DM, Hicks-Taylor CS, Zhang X, Sportiello MG, and Weisman GA (1992) Mechanisms by which extracellular ATP and UTP stimulate the release of prostacyclin from bovine pulmonary artery endothelial cells. *Biochim Biophys Acta* **1134**:61–72.
- Lustig KD, Shiau AK, Brake AJ, and Julius D (1993) Expression cloning of an ATP receptor from mouse neuroblastoma cells. *Proc Natl Acad Sci USA* **90**:5113–5117.
- Ma W, Korngreen A, Uzlauer N, Priel Z, and Silberberg SD (1999) Extracellular sodium regulates airway ciliary motility by inhibiting a P2X receptor. *Nature* **400**:894–897.
- Ma W, Korngreen A, Weil S, Cohen EB, Priel A, Kuzin L, and Silberberg SD (2006) Pore properties and pharmacological features of the P2X receptor channel in airway ciliated cells. *J Physiol* **571**:503–517.
- Maitre B, Freund M, Hechler B, Léon C, Heim V, Cazenave JP, Hanau D, and Gachet C (2008) Involvement of the P2Y₁ receptor in asthmatic airway inflammation (Abstract) *Purinergic Signal* **4** (Suppl 1):S156.
- Mall M, Wissner A, Gonska T, Calenborn D, Kuehr J, Brandis M, and Kunzelmann K (2000) Inhibition of amiloride-sensitive epithelial Na⁺ absorption by extracellular nucleotides in human normal and cystic fibrosis airways. *Am J Respir Cell Mol Biol* **23**:755–761.
- Mann JS, Cushley MJ, and Holgate ST (1985) Adenosine-induced bronchoconstriction in asthma. Role of parasympathetic stimulation and adrenergic inhibition. *Am Rev Respir Dis* **132**:1–6.
- Mann JS and Holgate ST (1985) Specific antagonism of adenosine-induced bronchoconstriction in asthma by oral theophylline. *Br J Clin Pharmacol* **19**:685–692.
- Mantell SJ, Stephenson PT, Monaghan SM, Maw GN, Trevethick MA, Yeaton D, Keir RF, Walker DK, Jones RM, Selby MD, et al. (2008) Inhaled adenosine A_{2A} receptor agonists for the treatment of chronic obstructive pulmonary disease. *Bioorg Med Chem Lett* **18**:1284–1287.
- Marcet B and Boeynaems JM (2006) Relationships between cystic fibrosis transmembrane conductance regulator, extracellular nucleotides and cystic fibrosis. *Pharmacol Ther* **112**:719–732.
- Marcet B, Horckmans M, Libert F, Hassid S, Boeynaems JM, and Communi D (2007a) Extracellular nucleotides regulate CCL20 release from human primary airway epithelial cells, monocytes and monocyte-derived dendritic cells. *J Cell Physiol* **211**:716–727.
- Marcet B, Libert F, Boeynaems JM, and Communi D (2007b) Extracellular nucleotides induce COX-2 up-regulation and prostaglandin E₂ production in human A549 alveolar type II epithelial cells. *Eur J Pharmacol* **566**:167–171.
- Marino A, Rodrig Y, Métioui M, Lagneaux L, Alzola E, Fernández M, Fogarty DJ, Matute C, Moran A, and Dehaye JP (1999) Regulation by P2 agonists of the intracellular calcium concentration in epithelial cells freshly isolated from rat trachea. *Biochim Biophys Acta* **1439**:395–405.
- Marmo E, Cazzola M, Chieppa S, Lampa E, Lobefalo G, and Filippelli A (1985) Documentation of presence of a purinergic reactivity at the tracheobronchial level in man. *Curr Ther Res* **38**:118–134.
- Masaki E, Ebisawa T, Kondo I, Hayashida K, Matsumoto Y, and Kawamura M (2000) P2-purinergic receptor antagonists reduce the minimum alveolar concentration of inhaled volatile anesthetics. *Brain Res* **864**:130–133.
- Mason SJ, Paradiso AM, and Boucher RC (1991) Regulation of transepithelial ion transport and intracellular calcium by extracellular ATP in human normal and cystic fibrosis airway epithelium. *Br J Pharmacol* **103**:1649–1656.
- Matera MG, De Santis D, D'Agostino B, Pallotta M, Vacca C, Cazzola M, and Rossi F (1995) Role of 5-hydroxytryptamine in mediating adenosine-induced airway contraction. *Immunopharmacology* **29**:73–78.
- Matot I, Weiniger CF, Zeira E, Galun E, Joshi BV, and Jacobson KA (2006) A3 adenosine receptors and mitogen-activated protein kinases in lung injury following in vivo reperfusion. *Crit Care* **10**:R65.
- Matsuyama H, Amaya F, Hashimoto S, Ueno H, Beppu S, Mizuta M, Shime N, Ishizaka A, and Hashimoto S (2008) Acute lung inflammation and ventilator-induced lung injury caused by ATP via the P2Y receptors: an experimental study. *Respir Res* **9**:79.
- McClenahan D, Hillenbrand K, Kapur A, Carlton D, and Czuprynski C (2009) Effects of extracellular ATP on bovine lung endothelial and epithelial cell monolayer morphologies, apoptosis, and permeabilities. *Clin Vaccine Immunol* **16**:43–48.
- McCormack DG, Barnes PJ, and Evans TW (1989) Purinoceptors in the pulmonary circulation of the rat and their role in hypoxic vasoconstriction. *Br J Pharmacol* **98**:367–372.
- McCormack DG, Clarke B, and Barnes PJ (1989) Characterization of adenosine receptors in human pulmonary arteries. *Am J Physiol* **256**:H41–H46.
- McCoy DE, Schwiebert EM, Karlson KH, Spielman WS, and Stanton BA (1995) Identification and function of A₁ adenosine receptors in normal and cystic fibrosis human airway epithelial cells. *Am J Physiol* **268**:C1520–C1527.
- McNamara N, Gallup M, Khong A, Sucher A, Maltseva I, Fahy J, and Basbaum C (2004) Adenosine up-regulation of the mucin gene, *MUC2*, in asthma. *FASEB J* **18**:1770–1772.
- McNamara N, Gallup M, Sucher A, Maltseva I, McKemy D, and Basbaum C (2006) AsialoGM1 and TLR5 cooperate in flagellin-induced nucleotide signaling to activate Erk1/2. *Am J Respir Cell Mol Biol* **34**:653–660.
- McQueen DS, Bond SM, Moores C, Chessell I, Humphrey PP, and Dowd E (1998) Activation of P2X receptors for adenosine triphosphate evokes cardiorespiratory reflexes in anaesthetized rats. *J Physiol* **507**:843–855.
- Meade CJ, Dumont I, and Worrall L (2001) Why do asthmatic subjects respond so strongly to inhaled adenosine? *Life Sci* **69**:1225–1240.
- Meltzer S, Goldberg B, Lad P, and Easton J (1989) Superoxide generation and its modulation by adenosine in the neutrophils of subjects with asthma. *J Allergy Clin Immunol* **83**:960–966.
- Merten MD, Breittmayer JP, Figarella C, and Frelin C (1993) ATP and UTP increase secretion of bronchial inhibitor by human tracheal gland cells in culture. *Am J Physiol* **265**:L479–L484.
- Merten MD, Saleh A, Kammouni W, Marchand S, and Figarella C (1998) Characterization of two distinct P2Y receptors in human tracheal gland cells. *Eur J Biochem* **251**:19–24.
- Miklavc P, Mair N, Wittekindt OH, Haller T, Dietl P, Felder E, Timmler M, and Frick M (2011) Fusion-activated Ca²⁺ entry via vesicular P2X₄ receptors promotes fusion pore opening and exocytotic content release in pneumocytes. *Proc Natl Acad Sci USA* **108**:14503–14508.
- Mikulski Z, Hartmann P, Jositsch G, Zaslona Z, Lips KS, Pfeil U, Kurzen H, Lohmeyer J, Claus WG, Grau V, et al. (2010) Nicotinic receptors on rat alveolar macrophages dampen ATP-induced increase in cytosolic calcium concentration. *Respir Res* **11**:133.
- Mishra A, Chintagari NR, Guo Y, Weng T, Su L, and Liu L (2011) Purinergic P2X₇ receptor regulates lung surfactant secretion in a paracrine manner. *J Cell Sci* **124**:657–668.
- Mizrahi J, Couture R, Caranikas S, and Regoli D (1982) Pharmacological effects of peptides on tracheal smooth muscle. *Pharmacology* **25**:39–50.
- Moerenhout M, Himpens B, and Vereecke J (2001) Intercellular communication upon mechanical stimulation of CPAE- endothelial cells is mediated by nucleotides. *Cell Calcium* **29**:125–136.
- Mohri K, Takeuchi K, Shinozuka K, Bjur RA, and Westfall DP (1993) Simultaneous determination of nerve-induced adenosine nucleotides and nucleosides released from rabbit pulmonary artery. *Anal Biochem* **210**:262–267.
- Mohsenin A and Blackburn MR (2006) Adenosine signaling in asthma and chronic obstructive pulmonary disease. *Curr Opin Pulm Med* **12**:54–59.
- Moon YW, Choi SH, Kim YT, Sohn JH, Chang J, Kim SK, Park MS, Chung KY, Lee HJ, and Kim JH (2007) Adenosine triphosphate-based chemotherapy response assay (ATP-CRA)-guided platinum-based 2-drug chemotherapy for unresectable non-small-cell lung cancer. *Cancer* **109**:1829–1835.
- Moon YW, Sohn JH, Kim YT, Chang H, Jeong JH, Lee YJ, Chang J, Kim SK, Jung M, Hong S, et al. (2009) Adenosine triphosphate-based chemotherapy response assay (ATP-CRA)-guided versus empirical chemotherapy in unresectable non-small cell lung cancer. *Anticancer Res* **29**:4243–4249.
- Moore JP, Hainsworth R, and Drinkhill MJ (2004) Pulmonary arterial distension and vagal afferent nerve activity in anaesthetized dogs. *J Physiol* **555**:805–814.
- Moraes DJ, Bonagamba LG, Zoccal DB, and Machado BH (2011) Modulation of respiratory responses to chemoreflex activation by L-glutamate and ATP in the rostral ventrolateral medulla of awake rats. *Am J Physiol Regul Integr Comp Physiol* **300**:R1476–R1486.
- Morschl E, Molina JG, Volmer JB, Mohsenin A, Pero RS, Hong JS, Kheradmand F, Lee JJ, and Blackburn MR (2008) A3 adenosine receptor signaling influences pulmonary inflammation and fibrosis. *Am J Respir Cell Mol Biol* **39**:697–705.
- Morse DM, Smullen JL, and Davis CW (2001) Differential effects of UTP, ATP, and adenosine on ciliary activity of human nasal epithelial cells. *Am J Physiol Cell Physiol* **280**:C1485–C1497.
- Mortaz E, Braber S, Nazary M, Givi ME, Nijkamp FP, and Folkerts G (2009) ATP in the pathogenesis of lung emphysema. *Eur J Pharmacol* **619**:92–96.
- Mortaz E, Folkerts G, Nijkamp FP, and Henricks PA (2010) ATP and the pathogenesis of COPD. *Eur J Pharmacol* **638**:1–4.
- Mounkaila B, Marthan R, and Roux E (2005) Biphasic effect of extracellular ATP on human and rat airways is due to multiple P2 purinoceptor activation. *Respir Res* **6**:143.
- Mueller RA, Widerlov E, and Breese GR (1984) Attempted antagonism of adenosine analogue induced depression of respiration. *Pharmacol Biochem Behav* **21**:289–296.
- Müller T, Bayer H, Myrtek D, Ferrari D, Sorichter S, Ziegenhagen MW, Zissel G, Virchow JC Jr, Luttmann W, Norgauer J, et al. (2005) The P2Y₁₄ receptor of airway epithelial cells: coupling to intracellular Ca²⁺ and IL-8 secretion. *Am J Respir Cell Mol Biol* **33**:601–609.
- Müller T, Robaye B, Vieira RP, Ferrari D, Grimm M, Jakob T, Martin SF, Di Virgilio F, Boeynaems JM, Virchow JC, et al. (2010) The purinergic receptor P2Y₂ receptor mediates chemotaxis of dendritic cells and eosinophils in allergic lung inflammation. *Allergy* **65**:1545–1553.
- Müller T, Vieira RP, Grimm M, Dürk T, Cicko S, Zeiser R, Jakob T, Martin SF, Blumenthal B, Sorichter S, et al. (2011) A potential role for P2X_{7R} in allergic airway inflammation in mice and humans. *Am J Respir Cell Mol Biol* **44**:456–464.
- Murphy JK, Livingston FR, Gozal E, Torres M, and Forman HJ (1993) Stimulation of the rat alveolar macrophage respiratory burst by extracellular adenosine nucleotides. *Am J Respir Cell Mol Biol* **9**:505–510.
- Musante L, Zegarra-Moran O, Montaldo PG, Ponzoni M, and Galletta LJ (1999)

- Autocrine regulation of volume-sensitive anion channels in airway epithelial cells by adenosine. *J Biol Chem* **274**:11701–11707.
- Mustafa SJ, Nadeem A, Fan M, Zhong H, Belardinelli L, and Zeng D (2007) Effect of a specific and selective A_{2B} adenosine receptor antagonist on adenosine agonist AMP and allergen-induced airway responsiveness and cellular influx in a mouse model of asthma. *J Pharmacol Exp Ther* **320**:1246–1251.
- Myrtek D, Müller T, Geyer V, Derr N, Ferrari D, Zissel G, Dürk T, Sorichter S, Luttmann W, Kuepper M, et al. (2008) Activation of human alveolar macrophages via P2 receptors: coupling to intracellular Ca²⁺ increases and cytokine secretion. *J Immunol* **181**:2181–2188.
- Nadeem A, Ponnath DS, Ansari HR, Batchelor TP, Dey RD, Ledent C, and Mustafa SJ (2009) A2A adenosine receptor deficiency leads to impaired tracheal relaxation via NADPH oxidase pathway in allergic mice. *J Pharmacol Exp Ther* **330**:99–108.
- Nagaoka M, Nara M, Tamada T, Kume H, Oguma T, Kikuchi T, Zaini J, Moriya T, Ichinose M, Tamura G, et al. (2009) Regulation of adenosine 5'-triphosphate (ATP)-gated P2X₄ receptors on tracheal smooth muscle cells. *Respir Physiol Neurobiol* **166**:61–67.
- Nakamura K, Yoshikawa N, Yamaguchi Y, Kagota S, Shinozuka K, and Kunitomo M (2006) Antitumor effect of cordycepin (3'-deoxyadenosine) on mouse melanoma and lung carcinoma cells involves adenosine A₃ receptor stimulation. *Anticancer Res* **26**:43–47.
- Nassenstein C, Taylor-Clark TE, Myers AC, Ru F, Nandigama R, Bettner W, and Undem BJ (2010) Phenotypic distinctions between neural crest and placodal derived vagal C-fibres in mouse lungs. *J Physiol* **588**:4769–4783.
- Neely CF, Haile DM, Cahill BE, and Kadowitz PJ (1991) Adenosine and ATP produce vasoconstriction in the feline pulmonary vasculature by different mechanisms. *J Pharmacol Exp Ther* **258**:753–761.
- Neely CF, Kadowitz PJ, Lippton H, Neiman M, and Hyman AL (1989) Adenosine does not mediate the pulmonary vasodilator response of adenosine 5'-triphosphate in the feline pulmonary vasculature. *J Pharmacol Exp Ther* **250**:170–176.
- Ni D and Lee LY (2008) Effect of increasing temperature on TRPV1-mediated responses in isolated rat pulmonary sensory neurons. *Am J Physiol Lung Cell Mol Physiol* **294**:L563–L571.
- Nieri P, Lazzari N, Martinotti E, and Breschi MC (1997) Histaminic bronchospasm potentiated by adenosine: investigation of the mechanisms. *Eur J Pharmacol* **331**:205–212.
- Nlend MC, Bookman RJ, Conner GE, and Salathe M (2002) Regulator of G-protein signaling protein 2 modulates purinergic calcium and ciliary beat frequency responses in airway epithelia. *Am J Respir Cell Mol Biol* **27**:436–445.
- Nlend MC, Schmid A, Sutto Z, Ransford GA, Conner GE, Fregien N, and Salathe M (2007) Calcium-mediated, purinergic stimulation and polarized localization of calcium-sensitive adenylyl cyclase isoforms in human airway epithelia. *FEBS Lett* **581**:3241–3246.
- Norsted T, Jonzon A, Niklasson F, and Sedin G (1987) Changes in inspiratory activity after injection of adenosine and hypoxanthine in cats. *Ups J Med Sci* **92**:265–275.
- Nyce JW and Metzger WJ (1997) DNA antisense therapy for asthma in an animal model. *Nature* **385**:721–725.
- Oguma T, Ito S, Kondo M, Makino Y, Shimokata K, Honjo H, Kamiya K, and Kume H (2007) Roles of P2X receptors and Ca²⁺ sensitization in extracellular adenosine triphosphate-induced hyperresponsiveness in airway smooth muscle. *Clin Exp Allergy* **37**:893–900.
- Ohata Y, Ogata S, Nakanishi K, Kanazawa F, Uenoyama M, Hiroi S, Tominaga S, and Kawai T (2011) Expression of P2X4R mRNA and protein in rats with hypobaric hypoxia-induced pulmonary hypertension. *Circ J* **75**:945–954.
- Okada SF, Nicholas RA, Kreda SM, Lazarowski ER, and Boucher RC (2006) Physiological regulation of ATP release at the apical surface of human airway epithelia. *J Biol Chem* **281**:22992–23002.
- Okada SF, Zhang L, Kreda SM, Abdullah LH, Davis CW, Pickles RJ, Lazarowski ER, and Boucher RC (2011) Coupled nucleotide and mucin hypersecretion from goblet-cell metaplastic human airway epithelium. *Am J Respir Cell Mol Biol* **45**:253–260.
- Oldenburg PJ and Mustafa SJ (2005) Involvement of mast cells in adenosine-mediated bronchoconstriction and inflammation in an allergic mouse model. *J Pharmacol Exp Ther* **313**:319–324.
- Orphanidou D, Stratakos G, Rasidakis A, Toumbis M, Samara J, Bakakos P, and Jordanoglou J (1998) Adenosine deaminase activity and lysozyme levels in bronchoalveolar lavage fluid in patients with pulmonary tuberculosis. *Int J Tuberc Lung Dis* **2**:147–152.
- Ott I, Lohse MJ, Klotz KN, Vogt-Moykopf I, and Schwabe U (1992) Effects of adenosine on histamine release from human lung fragments. *Int Arch Allergy Immunol* **98**:50–56.
- Paidas CN, Dudgeon DL, Haller JA Jr, and Clemens MG (1988) Adenosine triphosphate: a potential therapy for hypoxic pulmonary hypertension. *J Pediatr Surg* **23**:1154–1160.
- Paradiso AM (1997) ATP-activated basolateral Na⁺/H⁺ exchange in human normal and cystic fibrosis airway epithelium. *Am J Physiol* **273**:L148–L158.
- Paradiso AM, Ribeiro CM, and Boucher RC (2001) Polarized signaling via purinoceptors in normal and cystic fibrosis airway epithelia. *J Gen Physiol* **117**:53–67.
- Parr CE, Sullivan DM, Paradiso AM, Lazarowski ER, Burch LH, Olsen JC, Erb L, Weisman GA, Boucher RC, and Turner JT (1994) Cloning and expression of a human P_{2U} nucleotide receptor, a target for cystic fibrosis pharmacotherapy. *Proc Natl Acad Sci USA* **91**:3275–3279.
- Paruchuri S, Tashimo H, Peng C, Maekawa A, Xing W, Jiang Y, Kanaoka Y, Conley P, and Boyce JA (2009) Leukotriene E₄-induced pulmonary inflammation is mediated by the P2Y₁₂ receptor. *J Exp Med* **206**:2543–2555.
- Patel AS, Reigada D, Mitchell CH, Bates SR, Margulies SS, and Koval M (2005) Paracrine stimulation of surfactant secretion by extracellular ATP in response to mechanical deformation. *Am J Physiol Lung Cell Mol Physiol* **289**:L489–L496.
- Pauwels RA and Joos GF (1995) Characterization of the adenosine receptors in the airways. *Arch Int Pharmacodyn Ther* **329**:151–160.
- Pauwels R, Joos G, Kips J, and Van Der Straeten M (1990) Synergistic mechanisms in the adenosine- and neuropeptide-induced bronchoconstriction. *Arch Int Pharmacodyn Ther* **303**:113–121.
- Pauwels RA and Van der Straeten ME (1987) An animal model for adenosine-induced bronchoconstriction. *Am Rev Respir Dis* **136**:374–378.
- Pearl RG (1994) Adenosine produces pulmonary vasodilation in the perfused rabbit lung via an adenosine A₂ receptor. *Anesth Analg* **79**:46–51.
- Pelleg A and Hurt CM (1996) Mechanism of action of ATP on canine pulmonary vagal C fibre nerve terminals. *J Physiol* **490**:265–275.
- Pelleg A and Schulman ES (2002) Adenosine 5'-triphosphate axis in obstructive airway diseases. *Am J Ther* **9**:454–464.
- Pelletier S, Dubé J, Villeneuve A, Gobeil F Jr, Bernier SG, Battistini B, Guillemette G, and Sirois P (2000) Adenosine induces cyclic-AMP formation and inhibits endothelin-1 production/secretion in guinea-pig tracheal epithelial cells through A_{2B} adenosine receptors. *Br J Pharmacol* **129**:243–250.
- Persson CG, Erjefält I, and Karlsson JA (1981) Adenosine antagonism, a less desirable characteristic of xanthine asthma drugs? *Acta Pharmacol Toxicol (Copenh)* **49**:317–320.
- Persson CG (1982) Xanthines for asthma—present status. *Trends Pharmacol Sci* **3**:312–313.
- Phillis JW, Scislo TJ, and O'Leary DS (1997) Purines and the nucleus tractus solitarius: effects on cardiovascular and respiratory function. *Clin Exp Pharmacol Physiol* **24**:738–742.
- Picher M, Côté YP, Béliveau R, Potier M, and Beaudoin AR (1993) Demonstration of a novel type of ATP-diphosphohydrolase (EC 3.6.1.5) in the bovine lung. *J Biol Chem* **268**:4699–4703.
- Picher M, Béliveau R, Potier M, Savaria D, Rousseau E, and Beaudoin AR (1994) Demonstration of an ectoATP-diphosphohydrolase (E.C. 3.6.1.5) in non-vascular smooth muscles of the bovine trachea. *Biochim Biophys Acta* **1200**:167–174.
- Picher M and Boucher RC (2003) Human airway ecto-adenylate kinase. A mechanism to propagate ATP signaling on airway surfaces. *J Biol Chem* **278**:11256–11264.
- Picher M, Hiesh A, and Burch LH (2006) Imbalance between airway adenosine production and elimination in cystic fibrosis. *Purinergic Signal* **2**:231.
- Pintelon I, Brouns I, De Proost I, Van Meir F, Timmermans JP, and Adriaenssens D (2007) Sensory receptors in the visceral pleura: neurochemical coding and live staining in whole mounts. *Am J Respir Cell Mol Biol* **36**:541–551.
- Polosa R (2002) Adenosine-receptor subtypes: their relevance to adenosine-mediated responses in asthma and chronic obstructive pulmonary disease. *Eur Respir J* **20**:488–496.
- Polosa R and Blackburn MR (2009) Adenosine receptors as targets for therapeutic intervention in asthma and chronic obstructive pulmonary disease. *Trends Pharmacol Sci* **30**:528–535.
- Poulsen AN, Klausen TL, Pedersen PS, Willumsen NJ, and Frederiksen O (2005) Regulation of ion transport via apical purinergic receptors in intact rabbit airway epithelium. *Pflugers Arch* **450**:227–235.
- Poulsen AN, Klausen TL, Pedersen PS, Willumsen NJ, and Frederiksen O (2006) Nucleotide regulation of paracellular Cl⁻ permeability in natural rabbit airway epithelium. *Pflugers Arch* **452**:188–191.
- Polosa R, Rorke S, and Holgate ST (2002) Evolving concepts on the value of adenosine hyperresponsiveness in asthma and chronic obstructive pulmonary disease. *Thorax* **57**:649–654.
- Pratt AD, Clancy G, and Welsh MJ (1986) Mucosal adenosine stimulates chloride secretion in canine tracheal epithelium. *Am J Physiol* **251**:C167–C174.
- Prieto L, Gutiérrez V, Liñana J, and Marin J (2001) Bronchoconstriction induced by inhaled adenosine 5'-monophosphate in subjects with allergic rhinitis. *Eur Respir J* **17**:64–70.
- Qasabian RA, Schyvens C, Owe-Young R, Killen JP, Macdonald PS, Conigrave AD, and Williamson DJ (1997) Characterization of the P₂ receptors in rabbit pulmonary artery. *Br J Pharmacol* **120**:553–558.
- Quinton PM and Reddy MM (1992) Control of CFTR chloride conductance by ATP levels through non-hydrolytic binding. *Nature* **360**:79–81.
- Ralevic V and Burnstock G (1998) Receptors for purines and pyrimidines. *Pharmacol Rev* **50**:413–492.
- Ramminger SJ, Collet A, Baines DL, Murphie H, McAlroy HL, Olver RE, Inglis SK, and Wilson SM (1999) P2Y₂ receptor-mediated inhibition of ion transport in distal lung epithelial cells. *Br J Pharmacol* **128**:293–300.
- Ramsey BW (1996) Management of pulmonary disease in patients with cystic fibrosis. *N Engl J Med* **335**:179–188.
- Ramsingh R, Grygorczyk A, Solecki A, Cherkaoui LS, Berthiaume Y, and Grygorczyk R (2011) Cell deformation at the air-liquid interface induces Ca²⁺-dependent ATP release from lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol* **300**:L587–L595.
- Ransford GA, Fregien N, Qiu F, Dahl G, Conner GE, and Salathe M (2009) Pannexin 1 contributes to ATP release in airway epithelia. *Am J Respir Cell Mol Biol* **41**:525–534.
- Reeves JJ, Jones CA, Sheehan MJ, Vardey CJ, and Whelan CJ (1997) Adenosine A₃ receptors promote degranulation of rat mast cells both in vitro and in vivo. *Inflamm Res* **46**:180–184.
- Remsbury A, Rakhit S, and Wilson SM (1996) P_{2U} receptor agonists do not inhibit forskolin-evoked cAMP accumulation in A549 lung adenocarcinoma cells. *J Physiol* **495**:179P.
- Reuttershan J, Cagnina RE, Chang D, Linden J, and Ley K (2007) Therapeutic anti-inflammatory effects of myeloid cell adenosine receptor A_{2a} stimulation in lipopolysaccharide-induced lung injury. *J Immunol* **179**:1254–1263.
- Reuttershan J, Vollmer I, Stark S, Wagner R, Ngamsri KC, and Eltzschig HK (2009) Adenosine and inflammation: CD39 and CD73 are critical mediators in LPS-induced PMN trafficking into the lungs. *FASEB J* **23**:473–482.
- Reynolds SM, Docherty R, Robbins J, Spina D, and Page CP (2008) Adenosine induces a cholinergic tracheal reflex contraction in guinea pigs in vivo via an adenosine A1 receptor-dependent mechanism. *J Appl Physiol* **105**:187–196.

- Rice WR, Burhans M, and Wispé JR (1989) Effect of oxygen exposure on ATP content of rat bronchoalveolar lavage. *Pediatr Res* **25**:396–398.
- Rice WR, Burton FM, and Fiedeloy DT (1995) Cloning and expression of the alveolar type II cell P_{2U}-purinergic receptor. *Am J Respir Cell Mol Biol* **12**:27–32.
- Rice WR and Singleton FM (1986) P_{2U}-purinoceptors regulate surfactant secretion from rat isolated alveolar type II cells. *Br J Pharmacol* **89**:485–491.
- Rice WR and Singleton FM (1987) P_{2Y}-purinoceptor regulation of surfactant secretion from rat isolated alveolar type II cells is associated with mobilization of intracellular calcium. *Br J Pharmacol* **91**:833–838.
- Rice WR and Singleton FM (1989) Reactive blue 2 selectively inhibits P_{2Y}-purinoceptor-stimulated surfactant phospholipid secretion from rat isolated alveolar type II cells. *Br J Pharmacol* **97**:158–162.
- Rich PB, Douillet CD, Mahler SA, Husain SA, and Boucher RC (2003) Adenosine triphosphate is released during injurious mechanical ventilation and contributes to lung edema. *J Trauma* **55**:290–297.
- Richardson JB and Bouchard T (1975) Demonstration of a nonadrenergic inhibitory nervous system in the trachea of the guinea pig. *J Allergy Clin Immunol* **56**:473–480.
- Richardson J and Béland J (1976) Nonadrenergic inhibitory nervous systems in human airways. *J Appl Physiol* **41**:764–771.
- Riesenman PJ, Douillet CD, and Rich PB (2008) Lipopolysaccharide and mechanical ventilation-induced alterations in purinoceptor expression in the lung and extrapulmonary organ/severity department but does not increase perforation rate. *J Am Coll Surg* **207** (3 Suppl):S46–S47.
- Rimmer J, Peake HL, Santos CM, Lean M, Bardin P, Robson R, Haumann B, Loehrer F, and Handel ML (2007) Targeting adenosine receptors in the treatment of allergic rhinitis: a randomized, double-blind, placebo-controlled study. *Clin Exp Allergy* **37**:8–14.
- Riordan JR, Rommens JM, Kerem B, Alon N, Rozmahel R, Grzelczak Z, Zielenski J, Lok S, Plavsis N, and Chou JL (1989) Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA. *Science* **245**:1066–1073.
- Riteau N, Gasse P, Fauconnier L, Gombault A, Couegnat M, Fick L, Kanellopoulos J, Quesniaux VF, Marchand-Adam S, Crestani B, et al. (2010) Extracellular ATP is a danger signal activating P2X₇ receptor in lung inflammation and fibrosis. *Am J Respir Crit Care Med* **182**:774–783.
- Roman J, Rivera HN, Roser-Page S, Sitarman SV, and Ritzenthaler JD (2006) Adenosine induces fibronectin expression in lung epithelial cells: implications for airway remodeling. *Am J Physiol Lung Cell Mol Physiol* **290**:L317–L325.
- Rooney SA (2001) Regulation of surfactant secretion. *Comp Biochem Physiol A Mol Integr Physiol* **129**:233–243.
- Rorke S, Jennison S, Jeffs JA, Sampson AP, Arshad H, and Holgate ST (2002) Role of cysteinyl leukotrienes in adenosine 5'-monophosphate induced bronchoconstriction in asthma. *Thorax* **57**:323–327.
- Rossi AH, Salmon WC, Chua M, and Davis CW (2007) Calcium signaling in human airway goblet cells following purinergic activation. *Am J Physiol Lung Cell Mol Physiol* **292**:L92–L98.
- Ruan T, Lin YS, Lin KS, and Kou YR (2005) Sensory transduction of pulmonary reactive oxygen species by capsaicin-sensitive vagal lung afferent fibres in rats. *J Physiol* **565**:563–578.
- Rubino A and Burnstock G (1996) Evidence for a P_{2U}-purinoceptor mediating vasoconstriction by UTP, ATP and related nucleotides in the isolated pulmonary vascular bed of the rat. *Br J Pharmacol* **118**:1415–1420.
- Rubino A, Ziabary L, and Burnstock G (1999) Regulation of vascular tone by UTP and UDP in isolated rat intrapulmonary arteries. *Eur J Pharmacol* **370**:139–143.
- Rugolo M, Mastrocola T, Whörle C, Rasola A, Gruenert DC, Romeo G, and Galietta LJ (1993) ATP and A₁ adenosine receptor agonists mobilize intracellular calcium and activate K⁺ and Cl⁻ currents in normal and cystic fibrosis airway epithelial cells. *J Biol Chem* **268**:24779–24784.
- Russell JA (1980) Noradrenergic inhibitory innervation of canine airways. *J Appl Physiol* **48**:16–22.
- Russo C, Arcidiacono G, and Polosa R (2006) Adenosine receptors: promising targets for the development of novel therapeutics and diagnostics for asthma. *Fundam Clin Pharmacol* **20**:9–19.
- Ryzhov S, Novitskiy SV, Zaynagetdinov R, Goldstein AE, Carbone DP, Biaggioni I, Dikov MM, and Feoktistov I (2008) Host A_{2B} adenosine receptors promote carcinoma growth. *Neoplasia* **10**:987–995.
- Saano V, Muranen A, Nuutinen J, Karttunen P, and Silvasti M (1990) Effects of intravenous ATP on tracheal ciliary activity, airway resistance and haemodynamics in rats. *Pharmacol Toxicol* **66**:303–306.
- Saano V, Nuutinen J, Virta P, Joki S, Karttunen P, and Silvasti M (1991) The effect of ATP on the ciliary activity of normal and pathological human respiratory mucosa in vitro. *Acta Otolaryngol* **111**:130–134.
- Sabater JR, Mao YM, Shaffer C, James MK, O'Riordan TG, and Abraham WM (1999) Aerosolization of P2Y₂-receptor agonists enhances mucociliary clearance in sheep. *J Appl Physiol* **87**:2191–2196.
- Sakuma T, Gu X, Sugita M, Sagawa M, Sakuda M, and Toga H (2004) Uridine 5'-triphosphate stimulates alveolar fluid clearance in the isolated rat lungs. *J Pharmacol Sci* **95**:420–425.
- Saleh A, Figarella C, Kammouni W, Marchand-Pinatel S, Lazdunski A, Tubul A, Brun P, and Merten MD (1999) Pseudomonas aeruginosa quorum-sensing signal molecule N-(3-oxododecanoyl)-L-homoserine lactone inhibits expression of P2Y receptors in cystic fibrosis tracheal gland cells. *Infect Immun* **67**:5076–5082.
- Saleh A, Picher M, Kammouni W, Figarella C, and Merten MD (1999) Characterization of a diadenosine tetraphosphate-receptor distinct from the ATP-purinoceptor in human tracheal gland cells. *Eur J Pharmacol* **384**:91–98.
- Salonen RO, Mattila MJ, and Edholm LE (1982) Actions of theophylline, terbutaline, and ipratropium bromide alone and in combinations on the methacholine induced bronchoconstriction in rats. *Acta Pharmacol Toxicol (Copenh)* **51**:309–320.
- Satchell D (1982) Non-adrenergic, non-cholinergic nerves in mammalian airways: their function and the role of purines. *Comp Biochem Physiol C* **72**:189–196.
- Satchell D (1984a) Purine receptors in trachea: studies with adenosine deaminase and pyridamide. *Eur J Pharmacol* **102**:545–548.
- Satchell DG (1984b) Adenosine deaminase antagonizes inhibitory responses to adenosine and non-adrenergic, non-cholinergic inhibitory nerve stimulation in isolated preparations of guinea-pig trachea. *Br J Pharmacol* **83**:323–325.
- Satchell D and Smith R (1984) Adenosine causes contractions in spiral strips and relaxations in transverse strips of guinea-pig trachea: studies on mechanism of action. *Eur J Pharmacol* **101**:243–247.
- Satoh M, Yamaya M, Shimura S, Masuda T, Yamamoto M, Sasaki T, Sasaki H, and Shirato K (1995) Isoproterenol augments ATP-evoked Cl⁻ secretion across canine tracheal epithelium. *Respir Physiol* **99**:13–18.
- Schafer R, Sedehizade F, Welte T, and Reiser G (2003) ATP- and UTP-activated P2Y receptors differently regulate proliferation of human lung epithelial tumor cells. *Am J Physiol Lung Cell Mol Physiol* **285**:L376–L385.
- Schepp CP and Reutershan J (2008) Bench-to-bedside review: adenosine receptors—promising targets in acute lung injury? *Crit Care* **12**:226.
- Schmidt C, Bellingham MC, and Richter DW (1995) Adenosinergic modulation of respiratory neurones and hypoxic responses in the anaesthetized cat. *J Physiol* **483**:769–781.
- Schrader BJ, Inbar S, Kaufmann L, Vestal RE, and Rich S (1992) Comparison of the effects of adenosine and nifedipine in pulmonary hypertension. *J Am Coll Cardiol* **19**:1060–1064.
- Schreiber R, Castro H, and Kunzelmann K (2008) Allergen-induced airway hyper-responsiveness is absent in ecto-5'-nucleotidase (CD73)-deficient mice. *Pflügers Arch* **457**:431–440.
- Schreiber R and Kunzelmann K (2005) Purinergic P2Y₆ receptors induce Ca²⁺ and CFTR dependent Cl⁻ secretion in mouse trachea. *Cell Physiol Biochem* **16**:99–108.
- Schrier DJ, Lesch ME, Wright CD, and Gilbertsen RB (1990) The antiinflammatory effects of adenosine receptor agonists on the carrageenan-induced pleural inflammatory response in rats. *J Immunol* **145**:1874–1879.
- Schulman ES, Glaum MC, Post T, Wang Y, Raible DG, Mohanty J, Butterfield JH, and Pelleg A (1999) ATP modulates anti-IgE-induced release of histamine from human lung mast cells. *Am J Respir Cell Mol Biol* **20**:530–537.
- Schultz BD, Venglarik CJ, Bridges RJ, and Frizzell RA (1995) Regulation of CFTR Cl⁻ channel gating by ADP and ATP analogues. *J Gen Physiol* **105**:329–361.
- Schütz A, von Foerster U, and Bücheler M (2002) Adenosine triphosphatase expression correlates with ciliary activity of respiratory epithelium in vitro. *Cell Tiss Organs* **170**:246–250.
- Schwiebert EM and Zsembery A (2003) Extracellular ATP as a signaling molecule for epithelial cells. *Biochim Biophys Acta* **1615**:7–32.
- Seminario-Vidal L, Kreda S, Jones L, O'Neal W, Trejo J, Boucher RC, and Lazarewicz ER (2009) Thrombin promotes release of ATP from lung epithelial cells through coordinated activation of rho- and Ca²⁺-dependent signaling pathways. *J Biol Chem* **284**:20638–20648.
- Sévigny J, Picher M, Grondin G, and Beaudoin AR (1997) Purification and immunohistochemical localization of the ATP diphosphohydrolase in bovine lungs. *Am J Physiol* **272**:L939–L950.
- Sherwood CL, Lantz RC, Burgess JL, and Boitano S (2011) Arsenic alters ATP-dependent Ca²⁺ signaling in human airway epithelial cell wound response. *Toxicol Sci* **121**:191–206.
- Shimura S, Sasaki T, Nagaki M, Takishima T, and Shirato K (1994) Extracellular ATP regulation of feline tracheal submucosal gland secretion. *Am J Physiol* **267**:L159–L164.
- Shimura S (2001) Extracellular ATP regulation of airway secretion. *Drug Dev Res* **52**:170–177.
- Shin CY, Kim KC, Lee WJ, Jo MJ, Park KH, Dalby R, and Ko KH (2000) Inhaled ATP causes mucin release from goblet cells of intact rats. *Exp Lung Res* **26**:1–11.
- Shin JH, Namkung W, Choi JY, Yoon JH, and Lee MG (2004) Purinergic stimulation induces Ca²⁺-dependent activation of Na⁺-K⁺-2Cl⁻ cotransporter in human nasal epithelia. *J Biol Chem* **279**:18567–18574.
- Small RC (1982) Electrical slow waves and tone of guinea-pig isolated trachealis muscle: effects of drugs and temperature changes. *Br J Pharmacol* **77**:45–54.
- Small RC, Boyle JP, Cortijo J, Curtis-Prior PB, Davies JM, Foster RW, and Hofer P (1988) The relaxant and spasmogenic effects of some xanthine derivatives acting on guinea-pig isolated trachealis muscle. *Br J Pharmacol* **94**:1091–1100.
- Smith N and Broadley KJ (2008) Adenosine receptor subtypes in the airways responses to 5'-adenosine monophosphate inhalation of sensitized guinea-pigs. *Clin Exp Allergy* **38**:1536–1547.
- Smith RA, Alvarez AJ, and Estes DM (2001) The P2X₇ purinergic receptor on bovine macrophages mediates mycobacterial death. *Vet Immunol Immunopathol* **78**:249–262.
- Son M, Ito Y, Sato S, Ishikawa T, Kondo M, Nakayama S, Shimokata K, and Kume H (2004) Apical and basolateral ATP-induced anion secretion in polarized human airway epithelia. *Am J Respir Cell Mol Biol* **30**:411–419.
- Souhreda JF, Melzer E, and Grantham P (1980) Some characteristics of the purinergic nervous system in normal and sensitized airway smooth muscle. *Respir Physiol* **40**:199–209.
- Spicuzza L, Bonfiglio C, and Polosa R (2003) Research applications and implications of adenosine in diseased airways. *Trends Pharmacol Sci* **24**:409–413.
- Spicuzza L, Di Maria G, and Polosa R (2006) Adenosine in the airways: implications and applications. *Eur J Pharmacol* **533**:77–88.
- Sprague RS, Ellsworth ML, Stephenson AH, Kleinhenz ME, and Lonigro AJ (1998) Deformation-induced ATP release from red blood cells requires CFTR activity. *Am J Physiol* **275**:H1726–H1732.
- Sprague RS, Ellsworth ML, Stephenson AH, and Lonigro AJ (1996) ATP: the red blood cell link to NO and local control of the pulmonary circulation. *Am J Physiol* **271**:H2717–H2722.
- Sprague RS, Olearczyk JJ, Spence DM, Stephenson AH, Sprung RW, and Lonigro AJ (2003) Extracellular ATP signaling in the rabbit lung: erythrocytes as determinants of vascular resistance. *Am J Physiol Heart Circ Physiol* **285**:H693–H700.

- Spyer KM (2009) To breathe or not to breathe? That is the question. *Exp Physiol* **94**:1–10.
- Spyer KM and Gourine AV (2009) Chemosensory pathways in the brainstem controlling cardiorespiratory activity. *Philos Trans R Soc Lond B Biol Sci* **364**:2603–2610.
- Spyer KM and Thomas T (2000) A role for adenosine in modulating cardiorespiratory responses: a mini-review. *Brain Res Bull* **53**:121–124.
- Steinhorn RH, Morin FC 3rd, Van Wylen DG, Gugino SF, Giese EC, and Russell JA (1994) Endothelium-dependent relaxations to adenosine in juvenile rabbit pulmonary arteries and veins. *Am J Physiol* **266**:H2001–H2006.
- Stokes L and Surprenant A (2007) Purinergic P2Y₂ receptors induce increased MCP-1/CCL2 synthesis and release from rat alveolar and peritoneal macrophages. *J Immunol* **179**:6016–6023.
- Stokes L and Surprenant A (2009) Dynamic regulation of the P2X₄ receptor in alveolar macrophages by phagocytosis and classical activation. *Eur J Immunol* **39**:986–995.
- Stutts MJ and Boucher RC (1999) Cystic fibrosis gene and functions of CFTR implications of dysfunctional ion transport for pulmonary pathogenesis, in *Cystic Fibrosis in Adults* (Yankaskas JR and Knowles MR eds) pp 3–25, Lippincott-Raven, Philadelphia.
- Stutts MJ, Chinnet TC, Mason SJ, Fullton JM, Clarke LL, and Boucher RC (1992) Regulation of Cl⁻ channels in normal and cystic fibrosis airway epithelial cells by extracellular ATP. *Proc Natl Acad Sci USA* **89**:1621–1625.
- Stutts MJ, Fitz JG, Paradiso AM, and Boucher RC (1994) Multiple modes of regulation of airway epithelial chloride secretion by extracellular ATP. *Am J Physiol* **267**:C1442–C1451.
- Sun CX, Young HW, Molina JG, Volmer JB, Schnermann J, and Blackburn MR (2005) A protective role for the A₁ adenosine receptor in adenosine-dependent pulmonary injury. *J Clin Invest* **115**:35–43.
- Sun Y, Wu F, Sun F, and Huang P (2008) Adenosine promotes IL-6 release in airway epithelia. *J Immunol* **180**:4173–4181.
- Sustiel AM, Joseph B, Rocklin RE, and Borish L (1989) Asthmatic patients have neutrophils that exhibit diminished responsiveness to adenosine. *Am Rev Respir Dis* **140**:1556–1561.
- Swennen EL, Dagnelie PC, Van den Beucken T, and Bast A (2008a) Radioprotective effects of ATP in human blood ex vivo. *Biochem Biophys Res Commun* **367**:383–387.
- Swennen EL, Ummels V, Bast A, and Dagnelie P (2008b) Increased cytotoxicity of cisplatin in a human large cell lung carcinoma cell line by ATP. *Purinergic Signal* **4**:S207.
- Swennen EL, Ummels V, Buss I, Jaehde U, Bast A, and Dagnelie PC (2010) ATP sensitizes H460 lung carcinoma cells to cisplatin-induced apoptosis. *Chem Biol Interact* **184**:338–345.
- Szentmiklósi AJ, Ujfalusi A, Cseppentő A, Nosztray K, Kovács P, and Szabó JZ (1995) Adenosine receptors mediate both contractile and relaxant effects of adenosine in main pulmonary artery of guinea pigs. *Naunyn Schmiedebergs Arch Pharmacol* **351**:417–425.
- Szkotak AJ, Ng AM, Sawicka J, Baldwin SA, Man SF, Cass CE, Young JD, and Duszyk M (2001) Regulation of K⁺ current in human airway epithelial cells by exogenous and autocrine adenosine. *Am J Physiol Cell Physiol* **281**:C1991–C2002.
- Taira M, Tamaoki J, Nishimura K, Nakata J, Kondo M, Takemura H, and Nagai A (2002) Adenosine A₃ receptor-mediated potentiation of mucociliary transport and epithelial ciliary motility. *Am J Physiol Lung Cell Mol Physiol* **282**:L556–L562.
- Takano K, Shigetomi E, and Kato F (2004) P2X receptors mediate central inspiratory facilitation by low frequency vagal afferent inputs in the rabbit. *Soc Neurosci Abstr* **31**:661.10.
- Takeuchi K, Shinozuka K, Ishii R, and Hashimoto T (1995) Regional difference of endogenous ATP release in the pulmonary artery of rabbits. *Clin Exp Pharmacol Physiol* **22**:675–676.
- Tamaoki J, Kondo M, and Takizawa T (1989) Adenosine-mediated cyclic AMP-dependent inhibition of ciliary activity in rabbit tracheal epithelium. *Am Rev Respir Dis* **139**:441–445.
- Tang CM, Hoerning A, Büscher R, O'Connor DT, Ratjen F, Grasemann H, and Insel PA (2005) Human adenosine 2B receptor: SNP discovery and evaluation of expression in patients with cystic fibrosis. *Pharmacogenet Genomics* **15**:321–327.
- Tarran R (2007) Control of CFTR-mediated ASL secretion by adenosine and ENaC. *Pediatr Pulmonol* **42** (Suppl):174–175.
- Tarran R, Button B, and Boucher RC (2006) Regulation of normal and cystic fibrosis airway surface liquid volume by phasic shear stress. *Annu Rev Physiol* **68**:543–561.
- Tarran R, Button B, Picher M, Paradiso AM, Ribeiro CM, Lazarowski ER, Zhang L, Collins PL, Pickles RJ, Fredberg JJ, et al. (2005) Normal and cystic fibrosis airway surface liquid homeostasis. The effects of phasic shear stress and viral infections. *J Biol Chem* **280**:35751–35759.
- Tatur S, Kreda S, Lazarowski E, and Grygorczyk R (2008) Calcium-dependent release of adenosine and uridine nucleotides from A549 cells. *Purinergic Signal* **4**:139–146.
- Taylor AL, Kudlow BA, Marrs KL, Gruenert DC, Guggino WB, and Schwiebert EM (1998) Bioluminescence detection of ATP release mechanisms in epithelia. *Am J Physiol* **275**:C1391–C1406.
- Taylor AL, Schwiebert LM, Smith JJ, King C, Jones JR, Sorscher EJ, and Schwiebert EM (1999) Epithelial P2X purinergic receptor channel expression and function. *J Clin Invest* **104**:875–884.
- Taylor-Clark T and Udem BJ (2006) Transduction mechanisms in airway sensory nerves. *J Appl Physiol* **101**:950–959.
- Théâtre E, Bours V, and Oury C (2009) A P2X ion channel-triggered NF- κ B pathway enhances TNF- α -induced IL-8 expression in airway epithelial cells. *Am J Respir Cell Mol Biol* **41**:705–713.
- Thomas GS, Tammelin BR, Schiffman GL, Marquez R, Rice DL, Milikien D, and Mathur V (2008) Safety of regadenoson, a selective adenosine A_{2A} agonist, in patients with chronic obstructive pulmonary disease: A randomized, double-blind, placebo-controlled trial (RegCOPD trial). *J Nucl Cardiol* **15**:319–328.
- Thomas T, Ralevic V, Bordini M, Burnstock G, and Spyer KM (2001) Evidence for the involvement of purinergic signalling in the control of respiration. *Neuroscience* **107**:481–490.
- Thorne JR and Broadley KJ (1992) Adenosine-induced bronchoconstriction of isolated lung and trachea from sensitized guinea-pigs. *Br J Pharmacol* **106**:978–985.
- Thorne JR and Broadley KJ (2001) Responses to adenosine of isolated transverse or spiral strips of sensitized guinea pig trachea: role of epithelium. *Acta Pharmacol Sin* **22**:481–487.
- Thorne JR, Danahay H, and Broadley KJ (1996) Analysis of the bronchoconstrictor responses to adenosine receptor agonists in sensitized guinea-pig lungs and trachea. *Eur J Pharmacol* **316**:263–271.
- Tilley SL, Tsai M, Williams CM, Wang ZS, Erikson CJ, Galli SJ, and Koller BH (2003) Identification of A₃ receptor- and mast cell-dependent and -independent components of adenosine-mediated airway responsiveness in mice. *J Immunol* **171**:331–337.
- Trevethick MA, Mantell SJ, Stuart EF, Barnard A, Wright KN, and Yeadon M (2008) Treating lung inflammation with agonists of the adenosine A_{2A} receptor: promises, problems and potential solutions. *Br J Pharmacol* **155**:463–474.
- Tritto M, De Ponti R, Salerno-Uriarte JA, Spadacini G, Marazzi R, Moretti P, and Lanzotti M (2004) Adenosine restores atrio-venous conduction after apparently successful ostial isolation of the pulmonary veins. *Eur Heart J* **25**:2155–2163.
- Udem BJ, Chuaychoo B, Lee MG, Weinreich D, Myers AC, and Kollarik M (2004) Subtypes of vagal afferent C-fibres in guinea-pig lungs. *J Physiol* **556**:905–917.
- Udem BJ and Nassenstein C (2009) Airway nerves and dyspnea associated with inflammatory airway disease. *Respir Physiol Neurobiol* **167**:36–44.
- Urbach V, Walsh DE, Mainprice B, Bousquet J, and Harvey BJ (2002) Rapid non-genomic inhibition of ATP-induced Cl⁻ secretion by dexamethasone in human bronchial epithelium. *J Physiol* **545**:869–878.
- Urbach V, Higgins G, Al-Alawai M, Costello RW, McNally PM, Verriere V, Harvey BJ (2012) Regulation of ion transport and ASL height by the anti-inflammatory mediator, lipoxin A₄ in normal and cystic fibrosis bronchial epithelium. *FASEB J* **26**:696.6.
- Uzlaner N and Priel Z (1999) Interplay between the NO pathway and elevated [Ca²⁺]_i enhances ciliary activity in rabbit trachea. *J Physiol* **516**:179–190.
- van den Berge M, Hylkema MN, Versluis M, and Postma DS (2007) Role of adenosine receptors in the treatment of asthma and chronic obstructive pulmonary disease: recent developments. *Drugs R D* **8**:13–23.
- van den Berge M, Polosa R, Kerstjens HA, and Postma DS (2004) The role of endogenous and exogenous AMP in asthma and chronic obstructive pulmonary disease. *J Allergy Clin Immunol* **114**:737–746.
- Van Genechten J, Brouns I, Burnstock G, Timmermans JP, and Adriaensen D (2004) Quantification of neuroepithelial bodies and their innervation in fawn-hooded and Wistar rat lungs. *Am J Respir Cell Mol Biol* **30**:20–30.
- Van Scott MR, Chinnet TC, Burnette AD, and Paradiso AM (1995) Purinergic regulation of ion transport across nonciliated bronchiolar epithelial (Clara) cells. *Am J Physiol* **269**:L30–L37.
- Varani K, Caramori G, Vincenzi F, Adcock I, Casolari P, Leung E, Macleannan S, Gessi S, Morello S, Barnes PJ, et al. (2006) Alteration of adenosine receptors in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* **173**:398–406.
- Varani K, Caramori G, Vincenzi F, Tosi A, Barczyk A, Contoli M, Casolari P, Triggiani M, Hansel T, Leung E, et al. (2010) Oxidative/nitrosative stress selectively altered A_{2B} adenosine receptors in chronic obstructive pulmonary disease. *FASEB J* **24**:1192–1204.
- Vass G and Horváth I (2008) Adenosine and adenosine receptors in the pathomechanism and treatment of respiratory diseases. *Curr Med Chem* **15**:917–922.
- Vaughan RP, Szcwyczyk MT Jr, Lanosa MJ, Desesa CR, Gianutsos G, and Morris JB (2006) Adenosine sensory transduction pathways contribute to activation of the sensory irritation response to inspired irritant vapors. *Toxicol Sci* **93**:411–421.
- Versluis M, ten Hacken N, Postma D, Barroso B, Rutgers B, Geerlings M, Willemsse B, Timens W, and Hylkema M (2009) Adenosine receptors in COPD and asymptomatic smokers: effects of smoking cessation. *Virchows Arch* **454**:273–281.
- Versluis M, van den Berge M, Timens W, Luijk B, Rutgers B, Lammers JW, Postma DS, and Hylkema MN (2008) Allergen inhalation decreases adenosine receptor expression in sputum and blood of asthma patients. *Allergy* **63**:1186–1194.
- Vieira RP, Toledo AC, Ferreira SC, Santos AB, Medeiros MC, Hage M, Mauad T, Martins Mde A, Dolhnikoff M, and Carvalho CR (2011) Airway epithelium mediates the anti-inflammatory effects of exercise on asthma. *Respir Physiol Neurobiol* **175**:383–389.
- Vizi E, Huszár E, Csoma Z, Böszörményi-Nagy G, Barát E, Horváth I, Herjavec I, and Kollai M (2002) Plasma adenosine concentration increases during exercise: a possible contributing factor in exercise-induced bronchoconstriction in asthma. *J Allergy Clin Immunol* **109**:446–448.
- von Euler US and Liljestrand G (1946) Observations on the pulmonary arterial blood pressure in the cat. *Acta Physiol Scand* **12**:301–320.
- Wagner R, Ngamsri KC, Stark S, Vollmer I, and Reutershan J (2010) Adenosine receptor A₃ is a critical mediator in LPS-induced pulmonary inflammation. *Am J Physiol Lung Cell Mol Physiol* **299**:L502–L512.
- Wali A, Beers MF, Dodia C, Feinstein SI, and Fisher AB (1993) ATP and adenosine 3',5'-cyclic monophosphate stimulate the synthesis of surfactant protein A in rat lung. *Am J Physiol* **264**:L431–L437.
- Wang CH, Lin HC, Lin CH, Yu CT, Liu SL, Huang KH, Chung KF, and Kuo HP (2003) Effect of theophylline and specific phosphodiesterase IV inhibition on proliferation and apoptosis of progenitor cells in bronchial asthma. *Br J Pharmacol* **138**:1147–1155.
- Wang D, Sun Y, Zhang W, and Huang P (2008) Apical adenosine regulates basolateral Ca²⁺-activated potassium channels in human airway Calu-3 epithelial cells. *Am J Physiol Cell Physiol* **294**:C1443–C1453.
- Wang JL, Wu ZH, Pan BX, and Li J (2005) Adenosine A₁ receptors modulate the

- discharge activities of inspiratory and biphasic expiratory neurons in the medial region of Nucleus Retrofacialis of neonatal rat in vitro. *Neurosci Lett* **379**:27–31.
- Warburton D, Buckley S, and Cosico L (1989a) P1 and P2 purinergic receptor signal transduction in rat type II pneumocytes. *J Appl Physiol* **66**:901–905.
- Warburton D, Buckley S, Cosico L, and Forman HJ (1989b) Sublethal oxidant injury inhibits signal transduction in rat type II pneumocytes. *Am J Physiol* **257**:L217–L220.
- Wareham K, Vial C, Wykes RC, Bradding P, and Seward EP (2009) Functional evidence for the expression of P2X₁, P2X₄ and P2X₇ receptors in human lung mast cells. *Br J Pharmacol* **157**:1215–1224.
- Wegner CD (2001) P2 receptors in the respiratory system, in *Purinergic and Pyrimidinic Signalling II* (Abbraccio MP and Williams M eds) pp 281–300, Springer, Berlin.
- Weisman GA, Garrad RC, Erb LJ, Otero M, Gonzalez FA, and Clarke LL (1998) Structure and function of P2Y₂ nucleotide receptors in cystic fibrosis (CF) epithelium. *Adv Exp Med Biol* **431**:417–424.
- Welford LA and Anderson WH (1988) Purine receptors and guinea-pig trachea: evidence for a direct action of ATP. *Br J Pharmacol* **95**:689–694.
- Wesley UV, Bove PF, Hristova M, McCarthy S, and van der Vliet A (2007) Airway epithelial cell migration and wound repair by ATP-mediated activation of dual oxidase 1. *J Biol Chem* **282**:3213–3220.
- Wessberg P, Hedner J, Hedner T, Persson B, and Jonason J (1984) Adenosine mechanisms in the regulation of breathing in the rat. *Eur J Pharmacol* **106**:59–67.
- Wiklund NP, Cederqvist B, Matsuda H, and Gustafsson LE (1987) Adenosine can excite pulmonary artery. *Acta Physiol Scand* **131**:477–478.
- Willart MA and Lambrecht BN (2009) The danger within: endogenous danger signals, atopy and asthma. *Clin Exp Allergy* **39**:12–19.
- Wilson CG, Martin RJ, Jaber M, Abu-Shaweeh J, Jafri A, Haxhiu MA, and Zaidi S (2004) Adenosine A_{2A} receptors interact with GABAergic pathways to modulate respiration in neonatal piglets. *Respir Physiol Neurobiol* **141**:201–211.
- Wilson CN (2008) Adenosine receptors and asthma in humans. *Br J Pharmacol* **155**:475–486.
- Wilson CN and Batra VK (2002) Lipopolysaccharide binds to and activates A₁ adenosine receptors on human pulmonary artery endothelial cells. *J Endotoxin Res* **8**:263–271.
- Winters SL, Davis CW, and Boucher RC (2007) Mechanosensitivity of mouse tracheal ciliary beat frequency: roles for Ca²⁺, purinergic signaling, tonicity, and viscosity. *Am J Physiol Lung Cell Mol Physiol* **292**:L614–L624.
- Wolk KE, Lazarowski ER, Traylor ZP, Yu EN, Jewell NA, Durbin RK, Durbin JE, and Davis IC (2008) Influenza A virus inhibits alveolar fluid clearance in BALB/c mice. *Am J Respir Crit Care Med* **178**:969–976.
- Wong AM, Chow AW, Au SC, Wong CC, and Ko WH (2009) Apical versus basolateral P2Y₆ receptor-mediated Cl⁻ secretion in immortalized bronchial epithelia. *Am J Respir Cell Mol Biol* **40**:733–745.
- Wong LB and Yeates DB (1992) Luminal purinergic regulatory mechanisms of tracheal ciliary beat frequency. *Am J Respir Cell Mol Biol* **7**:447–454.
- Woodruff ML, Chaban VV, Worley CM, and Dirksen ER (1999) PKC role in mechanically induced Ca²⁺ waves and ATP-induced Ca²⁺ oscillations in airway epithelial cells. *Am J Physiol* **276**:L669–L678.
- Woodworth BA, Zhang S, Tamashiro E, Bhargava G, Palmer JN, and Cohen NA (2010) Zinc increases ciliary beat frequency in a calcium-dependent manner. *Am J Rhinol Allergy* **24**:6–10.
- Wyatt TA, Forget MA, Adams JM, and Sisson JH (2005) Both cAMP and cGMP are required for maximal ciliary beat stimulation in a cell-free model of bovine ciliary axonemes. *Am J Physiol Lung Cell Mol Physiol* **288**:L546–L551.
- Wyss D, Bonneau O, and Trifilieff A (2005) Mast cell involvement in the adenosine mediated airway hyper-reactivity in a murine model of ovalbumin-induced lung inflammation. *Br J Pharmacol* **145**:845–852.
- Xing H, Ling JX, Chen M, Johnson RD, Tominaga M, Wang CY, and Gu J (2008) TRPM8 mechanism of autonomic nerve response to cold in respiratory airway. *Mol Pain* **4**:22.
- Yamaguchi M, Shibata O, Saito M, Yoshimura M, Nishioka K, Makita T, and Sumikawa K (2007) Effects of propofol and ketamine on ATP-induced contraction of the rat trachea. *J Anesth* **21**:37–41.
- Yamamoto K, Sokabe T, Ohura N, Nakatsuka H, Kamiya A, and Ando J (2003) Endogenously released ATP mediates shear stress-induced Ca²⁺ influx into pulmonary artery endothelial cells. *Am J Physiol Heart Circ Physiol* **285**:H793–H803.
- Yamaya M, Sekizawa K, Kakuta Y, Ohru T, Sawai T, and Sasaki H (1996) P_{2U}-purinoceptor regulation of chloride secretion in cultured human tracheal submucosal glands. *Am J Physiol* **270**:L979–L984.
- Yegutkin GG (2008) Nucleotide- and nucleoside-converting ectoenzymes: Important modulators of purinergic signalling cascade. *Biochim Biophys Acta* **1783**:673–694.
- Yerxa BR (2001) Therapeutic use of nucleotides in respiratory and ophthalmic diseases. *Drug Dev Res* **52**:196–201.
- Yerxa BR, Sabater JR, Davis CW, Stutts MJ, Lang-Furr M, Picher M, Jones AC, Cowlen M, Dougherty R, Boyer J, et al. (2002) Pharmacology of INS37217 [P¹-(uridine 5′)-P⁴-(2′-deoxycytidine 5′)tetraphosphate, tetrasodium salt], a next-generation P2Y₂ receptor agonist for the treatment of cystic fibrosis. *J Pharmacol Exp Ther* **302**:871–880.
- Young HW, Molina JG, Dimina D, Zhong H, Jacobson M, Chan LN, Chan TS, Lee JJ, and Blackburn MR (2004) A₃ adenosine receptor signaling contributes to airway inflammation and mucus production in adenosine deaminase-deficient mice. *J Immunol* **173**:1380–1389.
- Young HW, Sun CX, Evans CM, Dickey BF, and Blackburn MR (2006) A₃ adenosine receptor signaling contributes to airway mucin secretion after allergen challenge. *Am J Respir Cell Mol Biol* **35**:549–558.
- Zhang AL and Roomans GM (1999) Regulation of intracellular calcium by extracellular nucleotides in pig tracheal submucosal gland cells. *Respir Physiol* **118**:237–246.
- Zhang HY, Gu YY, Li ZG, Jia YH, Yuan L, Li SY, An GS, Ni JH, and Jia HT (2004) Exposure of human lung cancer cells to 8-chloro-adenosine induces G₂/M arrest and mitotic catastrophe. *Neoplasia* **6**:802–812.
- Zhang L and Sanderson MJ (2003) Oscillations in ciliary beat frequency and intracellular calcium concentration in rabbit tracheal epithelial cells induced by ATP. *J Physiol* **546**:733–749.
- Zhao DM, Xue HH, Chida K, Suda T, Oki Y, Kanai M, Uchida C, Ichijima A, and Nakamura H (2000) Effect of erythromycin on ATP-induced intracellular calcium response in A549 cells. *Am J Physiol Lung Cell Mol Physiol* **278**:L726–L736.
- Zhong H, Belardinelli L, Maa T, Feoktistov I, Biaggioni I, and Zeng D (2004) A_{2B} adenosine receptors increase cytokine release by bronchial smooth muscle cells. *Am J Respir Cell Mol Biol* **30**:118–125.
- Zhong H, Chunn JL, Volmer JB, Fozard JR, and Blackburn MR (2001) Adenosine-mediated mast cell degranulation in adenosine deaminase-deficient mice. *J Pharmacol Exp Ther* **298**:433–440.
- Zhong H, Shlykov SG, Molina JG, Sanborn BM, Jacobson MA, Tilley SL, and Blackburn MR (2003) Activation of murine lung mast cells by the adenosine A₃ receptor. *J Immunol* **171**:338–345.
- Zhong H, Wu Y, Belardinelli L, and Zeng D (2006) A_{2B} adenosine receptors induce IL-19 from bronchial epithelial cells, resulting in TNF- α increase. *Am J Respir Cell Mol Biol* **35**:587–592.
- Zhou G, Dada LA, and Sznajder JI (2008) Regulation of alveolar epithelial function by hypoxia. *Eur Respir J* **31**:1107–1113.
- Zhou Y, Mohsenin A, Morschl E, Young HW, Molina JG, Ma W, Sun CX, Martinez-Valdez H, and Blackburn MR (2009) Enhanced airway inflammation and remodeling in adenosine deaminase-deficient mice lacking the A_{2B} adenosine receptor. *J Immunol* **182**:8037–8046.
- Zhou Y, Murthy JN, Zeng D, Belardinelli L, and Blackburn MR (2010) Alterations in adenosine metabolism and signaling in patients with chronic obstructive pulmonary disease and idiopathic pulmonary fibrosis. *PLoS One* **5**:e9224.
- Zhou Y, Schneider DJ, and Blackburn MR (2009) Adenosine signaling and the regulation of chronic lung disease. *Pharmacol Ther* **123**:105–116.
- Zhou Z, Wang X, Liu HY, Zou X, Li M, and Hwang TC (2006) The two ATP binding sites of cystic fibrosis transmembrane conductance regulator (CFTR) play distinct roles in gating kinetics and energetics. *J Gen Physiol* **128**:413–422.
- Zsembery A, Boyce AT, Liang L, Peti-Peterdi J, Bell PD, and Schwiebert EM (2003) Sustained calcium entry through P2X nucleotide receptor channels in human airway epithelial cells. *J Biol Chem* **278**:13398–13408.
- Zsembery A, Fortenberry JA, Liang L, Bekob Z, Tucker TA, Boyce AT, Braunstein GM, Welty E, Bell PD, Sorscher EJ, et al. (2004) Extracellular zinc and ATP restore chloride secretion across cystic fibrosis airway epithelia by triggering calcium entry. *J Biol Chem* **279**:10720–10729.
- Zuo P, Picher M, Okada SF, Lazarowski ER, Button B, Boucher RC, and Elston TC (2008) Mathematical model of nucleotide regulation on airway epithelia. Implications for airway homeostasis. *J Biol Chem* **283**:26805–26819.
- Zwicker JD, Rajani V, Hahn LB, and Funk GD (2011) Purinergic modulation of preBotzinger complex inspiratory rhythm in rodents: the interaction between ATP and adenosine. *J Physiol* **589**:4583–4600.