

sigma receptors in mouse brain using (+)-[³H]-SKF10047 as a ligand, and have attempted to compare the relative potencies of various drugs on *sigma* sites *in vivo* and *in vitro*. Mice were injected with 5 μ Ci of (+)-[³H]-SKF10047 into the tail vein. After various time intervals, the mice were decapitated, their brains were rapidly removed, weighed, homogenized and total and particulate (specific and nonspecific) bound radioactivity were determined (detailed methodology will be presented). Specifically bound (+)-[³H]-SKF10047 in the particulate fraction was defined as the difference in total radioactivity in the particulate fraction obtained from vehicle injected mice minus the radioactivity in the particulate fraction from Haldol (2 mg/kg IP) injected mice. Specifically bound (+)-[³H]-SKF10047 in the particulate fraction reached peak levels 30 min after IV injection, declined rapidly over the next 120 min, and constituted 90–95% of the total particulate radioactivity. Labeling of the *sigma* sites could be blocked *in vivo* by injecting mice IP with the drug 30 min before the IV injection of the [³H]-ligand. Under these conditions, the site in brain labeled by [³H]-(+)-SKF-10047 had the following characteristics: (1) naloxone insensitive; (2) stereoselective towards (+)-enantiomers of certain benzomorphan opiates like N-allylnormetazocine; (3) high affinity for haloperidol, (+)-3-PPP, cyclazocine, pentazocine and (+)-SKF-10047, and; (4) weak affinity for NMDA, (–)-3-PPP, PCP, and m-NH₂-PCP. This pharmacological profile would suggest that we are preferentially labeling the high affinity *sigma* site rather than the low affinity *sigma*/PCP site *in vivo* with [³H]-(+)-SKF-10047. Attempts to label the low affinity *sigma*/PCP site *in vivo* with ³H-TCP have failed. Thus, this *in vivo* binding assay should be a useful new technique for studying the effect of drugs on high affinity sites in the intact animal and for correlating these data with behavioral responses elicited over the same dose ranges.

INTERACTIONS OF PCP AND DERIVATIVES WITH THE BINDING OF ³H 5-HT AND ³H-MINAPRINE IN RAT BRAIN. Fillion, G., J. M. Sani, F. Christophe de Lamotte. Unite de Pharmacologie Neuroimmunoendocrinienne, Institut Pasteur, Paris, France.

PCP has been previously described to interact with the serotonergic system at the uptake sites (Smith, 1977) and at the 5-HT₂ receptors (Nabeshima, 1984). The present study was performed to examine the interactions of PCP (GK₁), TCP (GK₀) and GK₁₃ at the high affinity 5-HT sites. One class of sites (5-HT₁) corresponds to a serotonergic receptor able to recognize ³H 5-HT with a high affinity (K_D=3 nM) through distinct subclasses (5-HT_{1A/B/C/D}); one of them is likely related to a high apparent affinity adenylate cyclase activation. The effects of PCP and derivatives are not constantly observed at concentrations close to 10⁻⁵ M; they might induce a modest increase (+16%) (GK₁₃) or decrease (20–40%) (GK₀, GK₁) of the 5-HT₁ binding. The effects of these substances are quite constant and significant on a second population of ³H 5-HT binding sites having an intermediate affinity (K_D=10 nM). GK₀, GK₁ and GK₁₃ decrease the binding of ³H 5-HT with IC 50's close to 10⁻⁵ M. These results show that PCP and derivatives interact with ³H 5-HT binding; the interactions appear as non competitive phenomena. The effects of these substances also have been examined on the binding of an antidepressant, ³H-minaprine (³H-MIN) to hippocampal membranes which likely interacts

with the ³H 5-HT binding. An enhancement of the ³H-MIN binding has been observed which corresponded to an increase in the B_{max} accompanied by a change in Hill coefficient. These results suggest that PCP and derivatives not only may interact with the serotonergic function through complex molecular mechanisms affecting the binding of the amine to its specific sites, but indicate that they could also modify the binding of antidepressants to their specific sites. At the present time it is not known whether these molecular activities of PCP and derivatives correspond to clinical changes.

PHARMACOLOGICAL SPECIFICITY OF THE ELECTROPHYSIOLOGICAL EFFECTS OF PCP AND BENZOMORPHANS ON CEREBELLAR PURKINJE NEURONS. Freedman, R., Y. Wang, M. Kim, E. Moore, B. Hoffer and M. R. Palmer. Departments of Pharmacology and Psychiatry, University of Colorado Health Sciences Center, Denver, CO 80262.

Noradrenergic neurons have a well characterized input to cerebellar Purkinje neurons which we have previously found to be sensitive to presynaptic actions of phencyclidine (PCP). In our previous studies, we found that PCP causes depressions of the firing rates of single Purkinje neurons by potentiating synaptically released norepinephrine (NE). This effect appears to be caused by a blockade of synaptic reuptake of NE as well as, perhaps, by the potentiation of ongoing NE release. More recently, we have characterized the pharmacological specificity of PCP actions in cerebellum using the putative PCP receptor blocker, metaphit. Metaphit irreversibly blocked the electrophysiologically recorded depressions of Purkinje neurons caused by local applications of PCP, but not those caused by the inhibitory neurotransmitters, NE and GABA. Metaphit also blocked the depressions caused by local applications of the specific PCP-receptor agonist, dexodrol, but not the effects of its stereoisomer, levodrol. Furthermore, noncompetitive antagonists of *mu* and *delta* opiate receptors, BIT and FIT respectively, which contain an isothiocyanate moiety identical to that of metaphit, did not antagonize the effects of either dexodrol or PCP. We have also found that cyclazocine, a psychoactive benzomorphan, causes both metaphit-sensitive and metaphit-insensitive responses in cerebellum. The metaphit-insensitive responses were reversed by high doses of naloxone, suggesting a possible *kappa* opiate mechanism in addition to a metaphit-sensitive PCP mechanism. Both the (+) and (–) enantiomers of the benzomorphan *sigma* receptor agonist, SKF 10,047, caused depressions of Purkinje neurons which could be antagonized by metaphit. Unexpectedly, however, high doses of naloxone also partially antagonized the effects of these compounds. The naloxone applications also reversibly blocked the effects of concomitantly applied U-50, 488H, a highly selective *kappa* agonist, suggesting at least a small contribution of *kappa* mechanisms to the responses caused by both SKF 10,047 enantiomers.

PHENCYCLIDINE-INDUCED CHANGES IN A₁₀ DOPAMINE NEURONAL ACTIVITY AND LOCOMOTOR BEHAVIOR IN RATS CHRONICALLY TREATED WITH PCP. French, E. D. Maryland Psychiatric Research Center, University of Maryland, Baltimore, MD 21228.

A number of *in vivo* and *in vitro* studies have examined a