



Serotonergic lesions of the periaqueductal gray, a primary source of serotonin to the nucleus paragigantocellularis, facilitate sexual behavior in male rats

Joseph J. Normandin¹, Anne Z. Murphy^{*}

Department of Biology, Center for Behavioral Neuroscience, Georgia State University, Atlanta, GA 30302–4010, United States
Neuroscience Institute, Georgia State University, Atlanta, GA 30302–5010, United States

ARTICLE INFO

Article history:

Received 29 October 2010

Received in revised form 21 January 2011

Accepted 28 January 2011

Available online 3 February 2011

Keywords:

Genital reflexes

Ejaculation

Copulation

Periaqueductal gray

SSRI

ABSTRACT

While selective serotonin reuptake inhibitors (SSRIs) are widely used to treat anxiety and depression, they also produce profound disruptions of sexual function including delayed orgasm/ejaculation. The nucleus paragigantocellularis (nPGi), a primary source of inhibition of ejaculation in male rats, contains receptors for serotonin (5-HT). The ventrolateral periaqueductal gray (vlPAG) provides serotonin to this region, thus providing an anatomical and neurochemical basis for serotonergic regulation of the nPGi. We hypothesize that 5-HT acting at the nPGi could underlie the SSRI-induced inhibition of ejaculation in rodents. To this end, we produced 5-HT lesions of the source of 5-HT to the nPGi (the vlPAG) and examined sexual behavior. Removing the source of 5-HT to the nPGi facilitated genital reflexes, but not other aspects of sexual behavior, consistent with our hypothesis. Namely, 5-HT lesions produced a significant increase in the mean number of ejaculations and a significant decrease in ejaculation latency as compared to sham lesioned animals, while latency to mating and the post-ejaculatory interval did not differ. These data suggest that the serotonergic vlPAG–nPGi pathway is an important regulatory mechanism for the inhibition of ejaculation in rats and supports the hypothesis that this circuit contributes to SSRI-induced inhibition of ejaculation.

© 2011 Elsevier Inc. All rights reserved.

1. Introduction

Selective serotonin reuptake inhibitors (SSRIs) are the most commonly prescribed class of drugs for individuals with the clinical diagnosis of depression (Arroll et al., 2009; Koenig and Thase, 2009). SSRIs increase synaptic availability of serotonin (5-HT) by blocking reuptake at axon terminals (Hiemke and Hartter, 2000; Richelson, 1994). A well-documented side effect of SSRI treatment is sexual dysfunction (Kennedy and Rizvi, 2009; Schweitzer et al., 2009), with symptoms of decreased libido, premature ejaculation, delayed ejaculation, and anorgasmia most commonly reported.

The interaction between 5-HT and the neural circuitry underlying mammalian sexual behavior is not completely understood, although experiments in the male rat have suggested that serotonin can either inhibit or facilitate sexual behavior, depending on the brain site in question and/or the type of receptor targeted (Bitran and Hull, 1987; de Jong et al., 2006). For example, systemic administration of 5-HT or 5-hydroxytryptophan (a precursor to 5-HT synthesis) to male rats increases

latency to ejaculation (Ahlenius and Larsson, 1991; Gonzales et al., 1982), whereas the 5-HT synthesis inhibitor p-chlorophenylalanine decreases ejaculation latency (McIntosh and Barfield, 1984). In addition, general increases in 5-HT levels through chronic systemic administration of an SSRI produce delayed ejaculation in male rats (Ahlenius et al., 1980; de Jong et al., 2005; Mos et al., 1999; Vega Matuszyk et al., 1998) much like in humans (Hsu and Shen, 1995). In these studies, 5-HT was administered or manipulated systematically; therefore, the specific brain region(s) mediating the effects of 5-HT on sexual behavior are not known.

The nucleus paragigantocellularis (nPGi) of the rostroventrolateral medulla in rats, and the homologous structure in humans, the nucleus paragigantocellularis lateralis (Zec and Kinney, 2001), is the hypothesized site of descending inhibition of genital reflexes in both rats (Marson and McKenna, 1990) and humans (Johnson, 2006). The nPGi receives projections from upstream sites related to sexual behavior, including the paraventricular nucleus of the hypothalamus, medial preoptic area, and periaqueductal gray (Murphy and Hoffman, 2001; Murphy et al., 1999; Normandin and Murphy, 2008), and in turn, projects to the spinal motoneurons (pudendal motoneurons) innervating the bulbospongiosus and ischiocavernosus muscles (Hermann et al., 2003; Marson and Carson, 1999; Marson and McKenna, 1996; Tang et al., 1999), which are critical for ejaculation in male rats (Miura et al., 2001; Pescatori et al., 1993) and humans (Hsu et al., 2004; Shafik et al., 2009).

The rat nPGi and the human homolog contain 5-HT immunoreactive cells (Azmitia and Gannon, 1986; Marson and McKenna, 1992) and application of 5-HT to the spinal targets of the nPGi in male rats

^{*} Corresponding author at: Neuroscience Institute, Georgia State University, PO Box 5030, Atlanta, GA 30302–5030, United States. Tel.: +1 404 413 5332; fax: +1 404 413 5301.

E-mail addresses: j.normandin@nin.knaw.nl (J.J. Normandin), amurphy@gsu.edu (A.Z. Murphy).

¹ Present address: Netherlands Institute for Neuroscience, Neuroendocrinology Group, Meibergdreef 47, 1105 BA, Amsterdam, The Netherlands.

blocks the urethro-genital reflex (Marson and McKenna, 1992). Similarly, Clement et al. (2007) found that, in male rats, systemic administration of the SSRI dapoxetine reduced fictive ejaculatory reflexes, which are reversed by lesions of the nPGi. In addition, lesions of the nPGi in male rats block the inhibitory effects of systemic fluoxetine (an SSRI) on sexual behavior (Yells et al., 1994). These studies make it clear that 5-HT acts at the level of the spinal cord to inhibit genital reflexes and that 5-HT from the nPGi is a necessary antecedent for normal inhibition of genital reflexes. It is unclear, however, whether 5-HT may also be acting at the level of the nPGi to produce the observed effects.

The nPGi contains receptors for 5-HT including the 5-HT_{1A} (Thor et al., 1992, 1990), 5-HT_{1C} (Hoffman and Mezey, 1989), 5-HT_{2A} (Fay and Kubin, 2000; Fonseca et al., 2001), 5-HT_{2C} (Fonseca et al., 2001), and 5-HT₃ (Fonseca et al., 2001) subtypes, although only the 5-HT_{1A} and 5-HT_{2C} subtypes are found in abundance (Fonseca et al., 2001; Thor et al., 1990). The primary source of 5-HT to the nPGi is the ventrolateral periaqueductal gray (vlPAG). vlPAG neurons project directly to the nPGi (Murphy and Hoffman, 2001; Normandin and Murphy, 2008), and these vlPAG–nPGi cells co-localize with 5-HT (Bago et al., 2002) and sexual behavior-induced Fos (Normandin and Murphy, 2008), suggesting a role for the serotonergic vlPAG–nPGi pathway in male sexual behavior.

The present studies were designed to test the hypothesis that the effects of SSRIs on male sexual behavior are mediated, at least in part, via the nPGi. Our working hypothesis is that removal of the source of 5-HT to the nPGi would alter the expression of genital reflexes but not other aspects of sexual behavior.

2. Methods

2.1. Subjects

Thirty male Sprague–Dawley Rats (*Rattus norvegicus*; Charles River; 275–375 g) were same-sex double-housed in a temperature-controlled vivarium in reverse light (lights on 7:00 pm, lights off 7:00 am) with ad libitum access to food and water. All experiments were approved by the Georgia State Institutional Animal Care and Use Committee, with pain and suffering minimized in accordance with the committee's policies.

2.2. Ovariectomy and gonadal-steroid replacement in stimulus females

To induce sexual receptivity, ovariectomized female rats were injected with β -estradiol-3-benzoate (10 μ g/0.1 ml sesame oil s.c.; Sigma-Aldrich) 48 h before testing and progesterone (500 μ g/0.1 ml

sesame oil s.c.; Fluka) 4 h before testing as previously described (Barfield and Lisk, 1970; McEwen et al., 1987; Normandin and Murphy, 2011; Quadagno et al., 1972).

2.3. vlPAG 5-HT lesions

All surgeries were performed under aseptic conditions. Male rats ($n = 30$) were anesthetized by inhalation of isoflurane (2%–5%; Henry Schein) and placed in a stereotaxic frame. The skull was leveled such that bregma and lambda were at the same dorsoventral plane, and a craniotomy was performed dorsal to the PAG. vlPAG serotonergic-specific lesions were produced by bilateral injection of 5,7-dihydroxytryptamine creatine sulfate (5,7-DHT; 3 mg/ml in 0.9% saline 0.1% ascorbic acid; Fluka; $n = 22$ males) using a 1 μ l Hamilton syringe. Sham animals received equivolume injections of vehicle (0.9% saline 0.1% ascorbic acid; $n = 8$ males). Thirty minutes prior to 5,7-DHT administration, all animals received an injection of desipramine hydrochloride (25 mg/kg in 0.9% saline i.p.; Sigma) to block uptake of the 5-HT neurotoxin by noradrenergic transporters on noradrenergic cells (Bjorklund et al., 1975). The coordinates for the vlPAG were (in millimeters): AP, -8.5 bregma, ML, ± 0.75 , DV, -5.75 . 5,7-DHT or vehicle (300 nl/side) was slowly injected over 2 min and the syringe was left in place for 10 min before being removed. A small amount of bone wax was placed in the burr-hole and the animals' skin was wound-clipped. Animals received buprenorphine (0.1 mg/kg s.c.; Henry Schein) for pain relief and baytril (5 mg/kg i.m.; Henry Schein) as a prophylactic antibiotic. Animals recovered in clean heated cages before being returned to the housing facility.

2.4. Sexual behavior tests

Sexual behavior tests were conducted in acrylic aquariums (61 cm \times 30.5 cm \times 30.5 cm). Animals were acclimated to the arena prior to the initiation of the experiment for 10 min on two consecutive days. Three and six days later, males engaged in one-hour mating bouts with stimulus females to gain sexual experience. A third mating bout on day nine served as the baseline (pre-lesion) measure. Experimental animals then underwent lesion (or sham lesion) surgery as described above. Following a seven-day recovery, a final one-hour mating bout served as the post-lesion measure. All sexual behavior bouts were recorded and the number of mounts (attempted intromissions without the stereotyped pelvic thrust associated with an intromission), intromissions, and ejaculations, as well as the latency to begin mating (from placement of female into mating chamber to first mount or intromission), ejaculation latency, and




Fig. 1. Photomicrograph of sham lesion and lesion sites. A photomicrograph of 5-hydroxytryptophan (5-HTP) immunolabeled cells of the ventrolateral periaqueductal gray (vlPAG) and dorsal raphe (DR) is shown on the left, with a diagram (modified from Paxinos and Watson, 2005) at the approximate level of the tissue section shown on the right. Intra-vlPAG injection of vehicle (photomicrograph left) preserved 5-HTP immunoreactivity in the vlPAG. Intra-vlPAG injection of 5,7-dihydroxytryptamine (photomicrograph right) markedly reduced immunoreactivity for 5-HTP in the vlPAG. Lesion sites are indicated by the gray circles in the diagram. Aq indicates cerebral aqueduct.




Fig. 2. Measures of sexual behavior unrelated to genital reflex function. Serotonergic lesions of the vIPAG did not alter the latency to mating (A) or post-ejaculatory interval (B) from baseline to sexual behavior test within either the sham or lesion groups. Error bars indicates standard error of the mean (SEM).

post-ejaculatory interval were scored. Data were tabulated for each ejaculatory series and then compared across series.

2.5. Perfusion/fixation/tissue preparation

Following a euthanizing dose of SleepAway (0.5 ml i.p.; Henry Schein), all animals were transcardially perfused with 250 ml of 0.9% sodium chloride/2% sodium nitrite, followed by 300 ml of 4% paraformaldehyde 2.5% acrolein (Polysciences) in 0.1 M phosphate buffer then 150 ml of the sodium chloride/sodium nitrite solution. Following perfusion/fixation, brains were removed and stored at 4 °C in 30% sucrose solution until sectioned. Brains were cut into 25 μ m coronal sections in a 1:4 series through the PAG with a Leica 2000R freezing microtome and stored free-floating in cryoprotectant-antifreeze solution (Watson et al., 1986) at -20 °C.

2.6. Immunohistochemistry

Midbrain tissue sectioned at 25 μ m in a 1:4 series was processed for NeuN or 5-HTP for lesion verification as previously described (Loyd and Murphy, 2006; Murphy and Hoffman, 2001). Briefly, sections were removed from the cryoprotectant solution, rinsed extensively in potassium phosphate buffered saline (KPBS; pH 7.4), and then reacted for 20 min in 1% sodium borohydride to remove excess aldehydes. Sections were then incubated in primary antibody solution directed against either NeuN (Millipore, MAB377; monoclonal, raised in mouse; 1:70,000) or 5-HTP (Immunostar, 24446; polyclonal, raised in rabbit; 1:1000) in KPBS containing 0.1% Triton X for 1 h at room temperature followed by 48 h at 4 °C. After primary antibody incubation, tissue was rinsed in KPBS, incubated for 1 h in biotinylated goat-anti mouse (mouse anti-NeuN primary antibody) or goat anti-rabbit (rabbit anti-5-HTP primary antibody) IgG (Jackson ImmunoResearch) at a concentration of 1:600, rinsed in KPBS, followed by a 1 h incubation in avidin-biotin peroxidase complex (Vector Labs, ABC Elite Kit PK-6100) at a concentration of 1:10. After rinsing in KPBS and sodium acetate (0.175 M; pH 6.5), NeuN and 5-HTP were visualized as a black reaction product using nickel sulfate intensified 3,3'-diaminobenzidine solution containing 0.08% hydrogen peroxide in sodium acetate buffer. The reaction product was terminated after approximately 15 min by rinsing in sodium acetate buffer. Sections were mounted out of saline onto gelatin-subbed slides, air dried overnight, dehydrated in a series of graded alcohols, cleared in HistoClear, and cover-slipped using Permount.

2.7. Lesion analysis

5,7-DHT lesion and vehicle sham lesion sites were verified by light-microscopic analysis of vIPAG-containing brain sections immunohistochemically stained for NeuN or 5-HTP. The position of the bottom of the injection track was noted in the NeuN stained sections, and the absence or presence of 5-HTP was noted in 5-HTP stained sections. Only those injections centered in the vIPAG and with an absence of 5-HTP immunoreactivity in the vIPAG were considered for analysis.

2.8. Statistical analysis

Planned comparisons of mean sexual behavior within the lesion or sham groups were analyzed with two-tailed paired *t*-tests. Between group (lesion vs. sham) comparisons of means of sexual behavior were conducted using independent sample *t*-tests (two-tailed). All statistical comparisons were made with the alpha value set at 0.05.

3. Results

3.1. Lesion verification

An example of 5-HTP immunoreactivity in a lesioned animal and sham-lesioned animal is provided in Fig. 1. Serotonergic lesions were restricted to the vIPAG in the lesion ($n = 9$) group, with slight spread to the dorsal raphe nucleus, and encompassing an approximate 1 mm³ volume. Syringe paths in our sham-lesion group ($n = 7$) ended just dorsal to or within the vIPAG. The two animals with misplaced lesions (dorsal to the vIPAG) did not exhibit any obvious sexual behavior changes. In addition, no obvious changes in non-sexual behaviors were noted in any group.

3.2. Sexual behavior

Serotonergic lesions of the vIPAG did not affect behaviors not directly related to genital reflex function. For example, there was no

change from baseline in the mean latency to initiate copulation or in the mean post-ejaculatory interval within either group (Fig. 2). By contrast, serotonergic lesions of the vIPAG facilitated male sexual behavior in many measures related to genital reflex functions. While there was no change from baseline in the mean total of mounts or intromissions within either group (Fig. 3A and B), there was an increase from baseline in the mean number of ejaculations within the lesion group ($p=0.005$) but not the sham group ($p=1.000$). In addition, the mean number of ejaculations in the sexual behavior test was significantly greater in the lesion group than in the sham group ($p=0.039$; Fig. 3C). There was a decrease from baseline in the mean number of intromissions required for ejaculation within both groups (lesion, $p=0.004$; sham, $p=0.039$; Fig. 3D). There was a trend approaching statistical significance for a decrease from baseline in the mean ejaculation latency within the lesion group ($p=0.054$; Fig. 3E), and this trend is supported by the significantly shorter mean ejaculation latency in the sexual behavior test in the lesion group versus the sham group ($p=0.026$; Fig. 3E).

4. Discussion

Overall, serotonergic lesions of the vIPAG in male rats produced a facilitation of sexual behavior. In particular, 5-HT lesions of the vIPAG significantly increased the number of ejaculations during the mating bout, and there was a trend for a decrease in the mean latency to ejaculation. 5-HT vIPAG lesioned animals also had significantly shorter ejaculation latencies as compared to sham. There were no differences in the mean latency to mate or in the post-ejaculatory interval for either group, indicating that the effect of serotonergic vIPAG lesions is limited to those behaviors directly associated with genital reflexes.

The facilitation of sexual behavior we observed was specific to those measures associated with genital reflex function. Given the anatomical connections between vIPAG 5-HT cells and the nPGi (Bago et al., 2002; Normandin and Murphy, 2008; Underwood et al., 1999), as well as the concordance of effects in both vIPAG 5-HT lesions and nPGi lesions (Liu and Sachs, 1999; Yells et al., 1992), it is likely that the effects observed here are dependent on vIPAG 5-HT neurotransmission to the nPGi. In the context of a normally behaving animal, one would expect that 5-HT from the vIPAG serves to enhance nPGi activity, thereby increasing descending inhibition of genital reflexes. This serotonergic signal from the vIPAG would be particularly active in behavioral contexts where sexual behavior would not be appropriate (e.g., a “fight-or-flight” response) and likely “de-activated” when mating conditions were optimal (e.g., availability of a sexually receptive conspecific in the absence of a predator). This serotonergic vIPAG–nPGi pathway may effectively act as a gating mechanism for descending inhibition of genital reflexes.

Regulation of nPGi function (and thereby genital reflexes) through vIPAG is an interesting prospect as midbrain periaqueductal gray (PAG) cells are known to be involved in the coordination of cardiovascular responses to anxiety and stress (Johnson et al., 2004; Moraes et al., 2008; Murphy et al., 1995), nociception (Haghighparast and Ahmad-Molaei, 2009; Loyd et al., 2007), as well as social behaviors (Lonstein and Stern, 1998; Pavesi et al., 2007) and could therefore signal the current behavioral context to the nPGi. With respect to sexual behavior, we have previously found that the medial preoptic area (MPOA) of the hypothalamus, a region critical in the expression of male sexual behavior (Arendash and Gorski, 1983;

Malsbury, 1971), including genital reflexes (Giuliano et al., 1996; Marson and McKenna, 1994), provides input to the nPGi both directly (Normandin and Murphy, 2008) and through a PAG relay (Murphy and Hoffman, 2001). This MPOA–PAG pathway may provide necessary input to the nPGi regarding sexual “tone.” Indeed electrolytic lesions of the PAG, including the vIPAG, blocks normal MPOA-elicited bulbospongiosus contractions (Marson, 2004), suggesting that the MPOA–vIPAG–nPGi circuit is required for the elicitation of genital reflexes.

The medially adjacent dorsal raphe nucleus (DR) also contains serotonergic neurons that may have been affected by our lesions. However, results from previous work suggest that our observed findings are primarily the result of vIPAG lesions and not due to the spread of the toxin into the DR. For example, in male rats, lesions of the DR, or application of 5-HT or 5-HT agonists to the DR, have no effect on male ejaculatory behavior (Albinsson et al., 1996; Fernandez-Guasti et al., 1992; Hillegaart, 1991), although one study has reported that 5-HT DR manipulations shortened both the latency to ejaculate and the post-ejaculatory interval (McIntosh and Barfield, 1984). Interestingly, lesions of the DR also have been found to facilitate lordosis in male rats treated with estradiol (Takeyama and Yamanouchi, 1992), leading to speculation that the DR is not involved in male typical sexual behavior per se but rather inhibits female-typical behavior in male rats.






5. Conclusions

Serotonergic lesions of the vIPAG in male rats facilitated genital reflex function in a manner similar to nPGi lesions. This result lends evidence to the functional serotonergic connectivity between the vIPAG and nPGi. This work has implications for the occurrence of SSRI-induced delayed ejaculation in human males, and in particular, suggests that these drugs may be producing an effect by increasing serotonergic neurotransmission within the vIPAG–nPGi pathway. Limiting serotonergic drugs for mood disorders in humans to receptors and loci that influence mood regulation systems, but not sexual behavior systems, is an emerging area of research (Baldwin et al., 2006; Breuer et al., 2008; Kennedy and Rizvi, 2010). While it has been known that SSRIs can inhibit genital reflexes in male rats at the level of the spinal cord (Marson and McKenna, 1992), our work implies that this could also occur through the nPGi via vIPAG 5-HT input. Thus, in developing treatments for depression and anxiety, clinicians must consider both the spinal and supraspinal targets of serotonergic drugs in this genital reflex circuitry. In addition, SSRIs are often prescribed to men experiencing premature ejaculation (Hatzimouratidis et al., 2010; Hellstrom, 2009). A global increase in 5-HT is neither warranted nor desired in these patients, as other side effects may be incurred (Haddad and Dursun, 2008; Hellstrom, 2009) as a result of treatment. Targeting the serotonergic vIPAG pathway to the nPGi with appropriately specific drugs may be one way to circumvent global treatment of these patients with SSRIs. However, it remains to be elucidated which 5-HT receptor subtypes on nPGi cells are responsible for modulating nPGi activity.

Acknowledgments

The authors would like to acknowledge the expert technical contributions of Hila Eichenbaum and Vincent Laufer to this work. This work was supported by National Institutes of Health, grant

Fig. 3 Measures of sexual behaviors related to genital reflex function. Serotonergic lesions of the vIPAG did not alter the mean number of mounts (A) or intromissions (B) from baseline to sexual behavior test within either group. However, such lesions significantly increased the mean number of ejaculations (C) within the lesion group, which was significantly greater than in sham group. The mean number of intromission per ejaculation (D) significantly decreased within both the sham and lesion groups. The mean ejaculation latency (min; E) decreased within the lesion group, approaching significance, which was significantly less than the sham group. * $p<0.05$; error bars indicates SEM.



number DA16272 (to A.Z.M.), and National Science Foundation grant number IBN9876754 (to the Center for Behavioral Neuroscience).

References

- Ahlenius S, Larsson K. Opposite effects of 5-methoxy-N, N-di-methyl-tryptamine and 5-hydroxytryptophan on male rat sexual behavior. *Pharmacol Biochem Behav* 1991;38:201–5.
- Ahlenius S, Larsson K, Svensson L. Further evidence for an inhibitory role of central 5-HT in male rat sexual behavior. *Psychopharmacology (Berl)* 1980;68:217–20.
- Albinsson A, Andersson G, Andersson K, Vega-Matuszczyk J, Larsson K. The effects of lesions in the mesencephalic raphe systems on male rat sexual behavior and locomotor activity. *Behav Brain Res* 1996;80:57–63.
- Arendash GW, Gorski RA. Effects of discrete lesions of the sexually dimorphic nucleus of the preoptic area or other medial preoptic regions on the sexual behavior of male rats. *Brain Res Bull* 1983;10:147–54.
- Arroll B, Elley CR, Fishman T, Goodyear-Smith FA, Kenealy T, Blashki G, et al. Antidepressants versus placebo for depression in primary care. *Cochrane Database Syst Rev* 2009:CD007954.
- Azmitia EC, Gannon PJ. The primate serotonergic system: a review of human and animal studies and a report on *Macaca fascicularis*. *Adv Neurol* 1986;43:407–68.
- Bago M, Marson L, Dean C. Serotonergic projections to the rostroventrolateral medulla from midbrain and raphe nuclei. *Brain Res* 2002;945:249–58.
- Baldwin D, Bridgman K, Buis C. Resolution of sexual dysfunction during double-blind treatment of major depression with reboxetine or paroxetine. *J Psychopharmacol* 2006;20:91–6.
- Barfield MA, Lisk RD. Advancement of behavioral estrus by subcutaneous injection of progesterone in the 4-day cyclic rat. *Endocrinology* 1970;87:1096–8.
- Bitran D, Hull EM. Pharmacological analysis of male rat sexual behavior. *Neurosci Biobehav Rev* 1987;11:365–89.
- Bjorklund A, Baumgarten HG, Rensch A. 5,7-Dihydroxytryptamine: improvement of its selectivity for serotonin neurons in the CNS by pretreatment with desipramine. *J Neurochem* 1975;24:833–5.
- Breuer ME, Chan JS, Oosting RS, Groenink L, Korte SM, Campbell U, et al. The triple monoaminergic reuptake inhibitor DOV 216, 303 has antidepressant effects in the rat olfactory bulbectomy model and lacks sexual side effects. *Eur Neuropsychopharmacol* 2008;18:908–16.
- Clement P, Bernabe J, Gengo P, Denys P, Laurin M, Alexandre L, et al. Supraspinal site of action for the inhibition of ejaculatory reflex by dapoxetine. *Eur Urol* 2007;51:825–32.
- de Jong TR, Pattij T, Veening JG, Dederen PJ, Waldinger MD, Cools AR, et al. Citalopram combined with WAY 100635 inhibits ejaculation and ejaculation-related Fos immunoreactivity. *Eur J Pharmacol* 2005;509:49–59.
- de Jong TR, Veening JG, Waldinger MD, Cools AR, Olivier B. Serotonin and the neurobiology of the ejaculatory threshold. *Neurosci Biobehav Rev* 2006;30:893–907.
- Fay R, Kubin L. Pontomedullary distribution of 5-HT_{2A} receptor-like protein in the rat. *J Comp Neurol* 2000;418:323–45.
- Fernandez-Guasti A, Escalante AL, Ahlenius S, Hillegaart V, Larsson K. Stimulation of 5-HT_{1A} and 5-HT_{1B} receptors in brain regions and its effects on male rat sexual behaviour. *Eur J Pharmacol* 1992;210:121–9.
- Fonseca MI, Ni YG, Dunning DD, Mileti R. Distribution of serotonin 2A, 2C and 3 receptor mRNA in spinal cord and medulla oblongata. *Brain Res Mol Brain Res* 2001;89:11–9.
- Giuliano F, Rampin O, Brown K, Courtois F, Benoit G, Jardin A. Stimulation of the medial preoptic area of the hypothalamus in the rat elicits increases in intracavernous pressure. *Neurosci Lett* 1996;209:1–4.
- Gonzales G, Mendoza L, Ruiz J, Torrejon J. A demonstration that 5-hydroxytryptamine administered peripherally can affect sexual behavior in male rats. *Life Sci* 1982;31:2775–81.
- Haddad PM, Dursun SM. Neurological complications of psychiatric drugs: clinical features and management. *Hum Psychopharmacol* 2008;23(Suppl. 1):15–26.
- Haghighparast A, Ahmad-Molaei L. Effects of electrolytic lesion of dorsolateral periaqueductal gray on analgesic response of morphine microinjected into the nucleus cuneiformis in rat. *Neurosci Lett* 2009;451:165–9.
- Hatzimouratidis K, Amar E, Eardley I, Giuliano F, Hatzichristou D, Montorsi F, et al. Guidelines on male sexual dysfunction: erectile dysfunction and premature ejaculation. *Eur Urol* 2010;57(5):804–14.
- Hellstrom WJ. Emerging treatments for premature ejaculation: focus on dapoxetine. *Neuropsychiatr Dis Treat* 2009;5:37–46.
- Hermann GE, Holmes GM, Rogers RC, Beattie MS, Bresnahan JC. Descending spinal projections from the rostral gigantocellular reticular nuclei complex. *J Comp Neurol* 2003;455:210–21.
- Hiemke C, Hartter S. Pharmacokinetics of selective serotonin reuptake inhibitors. *Pharmacol Ther* 2000;85:11–28.
- Hillegaart V. Effects of local application of 5-HT and 8-OH-DPAT into the dorsal and median raphe nuclei on core temperature in the rat. *Psychopharmacology (Berl)* 1991;103:291–6.
- Hoffman BJ, Mezey E. Distribution of serotonin 5-HT_{1C} receptor mRNA in adult rat brain. *FEBS Lett* 1989;247:453–62.
- Hsu GL, Hsieh CH, Wen HS, Hsu WL, Wu CH, Fong TH, et al. Anatomy of the human penis: the relationship of the architecture between skeletal and smooth muscles. *J Androl* 2004;25:426–31.
- Hsu JH, Shen WW. Male sexual side effects associated with antidepressants: a descriptive clinical study of 32 patients. *Int J Psychiatry Med* 1995;25:191–201.
- Johnson PL, Lightman SL, Lowry CA. A functional subset of serotonergic neurons in the rat ventrolateral periaqueductal gray implicated in the inhibition of sympathoexcitation and panic. *Ann NY Acad Sci* 2004;1018:58–64.
- Johnson RD. Descending pathways modulating the spinal circuitry for ejaculation: effects of chronic spinal cord injury. *Prog Brain Res* 2006;152:415–26.
- Kakeyama M, Yamanouchi K. Lordosis in male rats: the facilitatory effect of mesencephalic dorsal raphe nucleus lesion. *Physiol Behav* 1992;51:181–4.
- Kennedy SH, Rizvi S. Sexual dysfunction, depression, and the impact of antidepressants. *J Clin Psychopharmacol* 2009;29:157–64.
- Kennedy SH, Rizvi SJ. Agomelatine in the treatment of major depressive disorder: potential for clinical effectiveness. *CNS Drugs* 2010;24(6):479–99.
- Koenig AM, Thase ME. First-line pharmacotherapies for depression—what is the best choice? *Pol Arch Med Wewn* 2009;119:478–86.
- Liu YC, Sachs BD. Erectile function in male rats after lesions in the lateral paraventricular nucleus. *Neurosci Lett* 1999;262:203–6.
- Lonstein JS, Stern JM. Site and behavioral specificity of periaqueductal gray lesions on postpartum sexual, maternal, and aggressive behaviors in rats. *Brain Res* 1998;804:21–35.
- Loyd DR, Morgan MM, Murphy AZ. Morphine preferentially activates the periaqueductal gray–rostral ventromedial medullary pathway in the male rat: a potential mechanism for sex differences in antinociception. *Neuroscience* 2007;147:456–68.
- Loyd DR, Murphy AZ. Sex differences in the anatomical and functional organization of the periaqueductal gray–rostral ventromedial medullary pathway in the rat: a potential circuit mediating the sexually dimorphic actions of morphine. *J Comp Neurol* 2006;496:723–38.
- Malsbury CW. Facilitation of male rat copulatory behavior by electrical stimulation of the medial preoptic area. *Physiol Behav* 1971;7:797–805.
- Marson L. Lesions of the periaqueductal gray block the medial preoptic area-induced activation of the urethro-genital reflex in male rats. *Neurosci Lett* 2004;367:278–82.
- Marson L, Carson III CC. Central nervous system innervation of the penis, prostate, and perineal muscles: a transneuronal tracing study. *Mol Urol* 1999;3:43–50.
- Marson L, McKenna KE. CNS cell groups involved in the control of the ischiocavernosus and bulbospongiosus muscles: a transneuronal tracing study using pseudorabies virus. *J Comp Neurol* 1996;374:161–79.
- Marson L, McKenna KE. The identification of a brainstem site controlling spinal sexual reflexes in male rats. *Brain Res* 1990;515:303–8.
- Marson L, McKenna KE. A role for 5-hydroxytryptamine in descending inhibition of spinal sexual reflexes. *Exp Brain Res* 1992;88:313–20.
- Marson L, McKenna KE. Stimulation of the hypothalamus initiates the urethro-genital reflex in male rats. *Brain Res* 1994;638:103–8.
- McEwen BS, Jones KJ, Pfaff DW. Hormonal control of sexual behavior in the female rat: molecular, cellular and neurochemical studies. *Biol Reprod* 1987;36:37–45.
- McIntosh TK, Barfield RJ. Brain monoaminergic control of male reproductive behavior. I. Serotonin and the post-ejaculatory refractory period. *Behav Brain Res* 1984;12:255–65.
- Miura T, Kondo Y, Akimoto M, Sakuma Y. Electromyography of male rat perineal musculature during copulatory behavior. *Urol Int* 2001;67:240–5.
- Moraes CL, Bertoglio LJ, Carobrez AP. Interplay between glutamate and serotonin within the dorsal periaqueductal gray modulates anxiety-related behavior of rats exposed to the elevated plus-maze. *Behav Brain Res* 2008;194:181–6.
- Mos J, Mollet I, Tolboom JT, Waldinger MD, Olivier B. A comparison of the effects of different serotonin reuptake blockers on sexual behaviour of the male rat. *Eur Neuropsychopharmacol* 1999;9:123–35.
- Murphy AZ, Ennis M, Rizvi TA, Behbehani MM, Shipley MT. Fos expression induced by changes in arterial pressure is localized in distinct, longitudinally organized columns of neurons in the rat midbrain periaqueductal gray. *J Comp Neurol* 1995;360:286–300.
- Murphy AZ, Hoffman GE. Distribution of gonadal steroid receptor-containing neurons in the preoptic–periaqueductal gray–brainstem pathway: a potential circuit for the initiation of male sexual behavior. *J Comp Neurol* 2001;438:191–212.
- Murphy AZ, Rizvi TA, Ennis M, Shipley MT. The organization of preoptic–medullary circuits in the male rat: evidence for interconnectivity of neural structures involved in reproductive behavior, antinociception and cardiovascular regulation. *Neuroscience* 1999;91:1103–16.
- Normandin JJ, Murphy AZ. Excitotoxic lesions of the nucleus paragigantocellularis facilitate male sexual behavior but attenuate female sexual behavior in rats. *Neuroscience* 2011;175:212–223.
- Normandin JJ, Murphy AZ. Nucleus paragigantocellularis afferents in male and female rats: organization, gonadal steroid receptor expression, and activation during sexual behavior. *J Comp Neurol* 2008;508:771–94.
- Pavesi E, Enck MC, De Toledo CA, Terenzi MG. Disruption of maternal behaviour by acute specific interaction induces selective activation of the lateral periaqueductal grey. *Eur J Neurosci* 2007;26:2055–65.
- Paxinos G, Watson C. The rat brain in stereotaxic coordinates. Amsterdam; Boston: Elsevier Academic Press; 2005.
- Pescatori ES, Calabro A, Artibani W, Pagano F, Triban C, Italiano G. Electrical stimulation of the dorsal nerve of the penis evokes reflex tonic erections of the penile body and reflex ejaculatory responses in the spinal rat. *J Urol* 1993;149:627–32.
- Quadagno DM, McCullough J, Langan R. The effect of varying amounts of exogenous estradiol benzoate on estrous behavior in the rat. *Horm Behav* 1972;3:175–9.
- Richelson E. The pharmacology of antidepressants at the synapse: focus on newer compounds. *J Clin Psychiatry* 1994;55(Suppl. A):34–9 discussion 40–31, 98–100.
- Schweitzer I, Maguire K, Ng C. Sexual side-effects of contemporary antidepressants: review. *Aust N Z J Psychiatry* 2009;43:795–808.

- Shafik A, Shafik AA, El Sibai O, Shafik IA. Electromyographic study of ejaculatory mechanism. *Int J Androl* 2009;32:212–7.
- Tang Y, Rampin O, Giuliano F, Ugolini G. Spinal and brain circuits to motoneurons of the bulbospongiosus muscle: retrograde transneuronal tracing with rabies virus. *J Comp Neurol* 1999;414:167–92.
- Thor KB, Blitz-Siebert A, Helke CJ. Autoradiographic localization of 5HT1 binding sites in the medulla oblongata of the rat. *Synapse* 1992;10:185–205.
- Thor KB, Blitz-Siebert A, Helke CJ. Discrete localization of high-density 5-HT1A binding sites in the midline raphe and parapyramidal region of the ventral medulla oblongata of the rat. *Neurosci Lett* 1990;108:249–54.
- Underwood MD, Arango V, Bakalian MJ, Ruggiero DA, Mann JJ. Dorsal raphe nucleus serotonergic neurons innervate the rostral ventrolateral medulla in rat. *Brain Res* 1999;824:45–55.
- Vega Matuszczyk J, Larsson K, Eriksson E. The selective serotonin reuptake inhibitor fluoxetine reduces sexual motivation in male rats. *Pharmacol Biochem Behav* 1998;60:527–32.
- Watson Jr RE, Wiegand SJ, Clough RW, Hoffman GE. Use of cryoprotectant to maintain long-term peptide immunoreactivity and tissue morphology. *Peptides* 1986;7:155–9.
- Yells DP, Hendricks SE, Prendergast MA. Lesions of the nucleus paragigantocellularis: effects on mating behavior in male rats. *Brain Res* 1992;596:73–9.
- Yells DP, Prendergast MA, Hendricks SE, Nakamura M. Fluoxetine-induced inhibition of male rat copulatory behavior: modification by lesions of the nucleus paragigantocellularis. *Pharmacol Biochem Behav* 1994;49:121–7.
- Zec N, Kinney HC. Anatomic relationships of the human nucleus paragigantocellularis lateralis: a Dil labeling study. *Auton Neurosci* 2001;89:110–24.