

Nuclear Magnetic Resonance Spectroscopy in Solids

Rex Richards and K. J. Packer

Phil. Trans. R. Soc. Lond. A 1981 **299**, 475-476
doi: 10.1098/rsta.1981.0028

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Phil. Trans. R. Soc. Lond. A* go to: <http://rsta.royalsocietypublishing.org/subscriptions>

Phil. Trans. R. Soc. Lond. A **299**, 475–686 (1981) [475]

Printed in Great Britain

NUCLEAR MAGNETIC RESONANCE SPECTROSCOPY IN SOLIDS

A DISCUSSION ORGANIZED BY SIR REX RICHARDS, F.R.S., AND K. J. PACKER

(Discussion held 18 and 19 June 1980 – Typescripts received 17 September 1980)

CONTENTS

	PAGE
SIR REX RICHARDS, F.R.S.	
Introduction	477
P. MANSFIELD	
Multiple-pulse n.m.r. experiments in solids: an introduction to symmetrized pulse sequences	479
<i>Discussion: J. G. Powles</i>	495
U. HAEBERLEN	
Achievements of multiple-pulse n.m.r.	497
<i>Discussion: R. K. HARRIS</i>	503
E. R. ANDREW	
Magic angle spinning in solid state n.m.r. spectroscopy	505
B. C. GERSTEIN	
High-resolution n.m.r. in solids with strong homonuclear dipolar broadening: combined multiple-pulse decoupling and magic angle spinning	521
<i>Discussion: K. J. PACKER</i>	545
R. G. GRIFFIN, G. BODENHAUSEN, R. A. HABERKORN, T. H. HUANG, M. MUNOWITZ, R. OSREDKAR, D. J. RUBEN, R. E. STARK AND H. VAN WILLIGEN	
High-resolution dipolar n.m.r. spectra in solids	547
M. E. STOLL	
Dipolar oscillations and phase interferometry in solid state n.m.r.	565
G. DROBNY, A. PINES, S. SINTON, W. S. WARREN AND D. P. WEITEKAMP	
Selectivity in multiple-quantum spectroscopy	585
J. SCHAEFER, E. O. STEJSKAL, M. D. SEFCIK AND R. A. MCKAY	
Applications of high-resolution ^{13}C and ^{15}N n.m.r. of solids	593
<i>Discussion: G. C. LEVY</i>	608

A. N. GARROWAY, D. L. VANDERHART AND W. L. EARL	
¹³ C n.m.r. in organic solids: limits to spectral resolution and to determination of molecular motion	609
Discussion: W. S. VEEMAN	628
W. S. VEEMAN	
¹³ C chemical shift tensors in organic single crystals	629
Discussion: D. J. GREENSLADE	641
G. E. BALIMANN, C. J. GROOMBRIDGE, R. K. HARRIS, K. J. PACKER, B. J. SAY AND S. F. TANNER	
Chemical applications of high-resolution ¹³ C n.m.r. spectra for solids	643
Discussion: W. DERBYSHIRE, B. C. GERSTEIN, J. E. PAGE	662
S. J. OPELLA, J. G. HEXEM, M. H. FREY AND T. A. CROSS	
Solid state n.m.r. of biopolymers	665
Discussion: W. DERBYSHIRE	682
R. FREEMAN, F.R.S.	
Concluding remarks	685

A point of notation

In the absence of an internationally approved and formally correct symbol, in this issue the notion 'parts per million' is represented by ' $\times 10^{-6}$ ' or '/ 10^6 '. This is in accordance with the recommendation contained in *Quantities, Units and Symbols* (The Royal Society, 1975). Where no numerical value is given, the expression 'millionths' is used.