

Corrigenda: The Electrical Response of Isolated Salivary Glands during Stimulation with 5-Hydroxytryptamine and Cyclic AMP

The Royal Society

Phil. Trans. R. Soc. Lond. B 1971 **262**, 458 doi: 10.1098/rstb.1971.0107

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click **here**

To subscribe to Phil. Trans. R. Soc. Lond. B go to: http://rstb.royalsocietypublishing.org/subscriptions

458

CORRIGENDA

G. A. Harrison and others. Phil. Trans. R. Soc. Lond. B 256, 147-182 (1969)

Page 173, footnote to table 18, line 5, middle of line:

R. A. Beatty. Phil. Trans. R. Soc. Lond. B 259, 3-13 (1970)

Page 4, lines 2 and 4 from the bottom: interchange isopods and amphipod

Page 14, the middle of the last paragraph should read: ...for Drosophila was that two types of sperm exist, one of which is physiologically abnormal and the other physiologically normal. Certain gene segregations are postulated to be preferential during meiosis in heterozygous males relative to the physiological classes. Thus distorted ratios...

(In some copies this paragraph already reads correctly.)

M. J. Berridge and W. T. Prince. Phil. Trans. R. Soc. Lond. B 262, 111-120 (1971)

Page 117, figure 8, ordinate scale:

$$for -10 -20 -30$$

 $read +10 +20 +30$