
Xwnt11* and the regulation of gastrulation in *Xenopus

J.C. Smith, Frank L. Conlon, Yasushi Saka and Masazumi Tada

Phil. Trans. R. Soc. Lond. B 2000 **355**, 923-930
doi: 10.1098/rstb.2000.0627

References

Article cited in:

<http://rstb.royalsocietypublishing.org/content/355/1399/923#related-urls>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

To subscribe to *Phil. Trans. R. Soc. Lond. B* go to: <http://rstb.royalsocietypublishing.org/subscriptions>

Xwnt11 and the regulation of gastrulation in *Xenopus*

J. C. Smith*, Frank L. Conlon, Yasushi Saka and Masazumi Tada

Division of Developmental Biology, National Institute for Medical Research, The Ridgeway, Mill Hill, London NW7 1AA, UK

The molecular basis of gastrulation is poorly understood. In this paper we address this problem by taking advantage of the observation that the transcription activator Brachyury is essential for gastrulation movements in *Xenopus* and mouse embryos. We infer from this observation that amongst the target genes of Brachyury are some that are involved in the regulation of gastrulation. In the course of a screen for Brachyury targets we identified *Xwnt11*. Use of a dominant-negative *Xwnt11* construct confirms that signalling by this class of Wnts is essential for normal gastrulation movements, and further investigation suggests that *Xwnt11* signals not through the canonical Wnt signalling pathway involving GSK-3 and β -catenin but through another route, which may require small GTPases such as Rho and Rac. Future work will concentrate on elucidating the *Xwnt11* signal transduction pathway and on investigating its influence on cell shape and polarity during *Xenopus* gastrulation.

Keywords: morphogenesis; gastrulation; Brachyury; Wnt signalling

1. INTRODUCTION

Although the gastrulation movements of *Xenopus laevis* have been described in great detail and with great insight by Keller and his colleagues (this issue), it remains true that we understand very little about the molecular basis of gastrulation in this species or, indeed, in any other animal. What are the molecular changes that cause bottle cell formation? How do cells change their adhesion properties? Which molecules drive intercalation? What confers polarity? Our ignorance of this fundamental morphogenetic process is quite profound. In this paper we address the question by taking advantage of our knowledge of the transcription factor Brachyury (Herrmann *et al.* 1990; Papaioannou 1997; Smith 1997). Previous work has shown that Brachyury function is required for normal gastrulation movements in both *Xenopus* and mouse embryos (Conlon *et al.* 1996; Conlon & Smith 1999; Wilson *et al.* 1995). Brachyury is a transcription activator (Conlon *et al.* 1996; Kispert *et al.* 1995a), and this suggests that its downstream targets include genes that are involved in the control of gastrulation. A search for such targets has identified *Xwnt11*, a member of the Wnt family of secreted signalling molecules (Ku & Melton 1993). Our experiments show that *Xwnt11*, like Brachyury, is required for normal gastrulation movements in *Xenopus*, and they suggest that *Xwnt11* does not act through the canonical Wnt signalling pathway involving GSK-3 and β -catenin (Cadigan & Nusse 1997) but through a pathway similar to that involved in planar polarity signalling in *Drosophila* (Axelrod *et al.* 1998; Boutros & Mlodzik 1999; Boutros *et al.* 1998). It is possible, therefore, that *Xwnt11* signalling functions to control cell polarity during gastrulation in *Xenopus*,

perhaps by regulating the function of small GTPases of the Rho family.

2. BRACHYURY

The *Brachyury* (or *T*) gene was first identified over 70 years ago. Mice that are heterozygous for the *Brachyury* mutation have a short tail, while homozygous mutant embryos do not form a proper allantois or notochord and lack mesoderm posterior to somite 7 (Chesley 1935; Dobrovolskaia-Zavadskaja 1927; Gluecksohn-Schoenheimer 1944). This dramatic mutant phenotype indicated that *Brachyury* plays a key role in the formation of mesodermal structures, but significant progress in the analysis of *Brachyury* function only came with the cloning of the gene by Herrmann *et al.* (1990). This work revealed that *Brachyury* encodes a protein of 436 amino acids and that the gene is expressed at the highest levels, and for the longest times, in those structures that are absent in homozygous mutant embryos (Wilkinson *et al.* 1990). Thus transcripts are present initially throughout the primitive streak of the embryo, and expression is maintained in the notochord and tailbud.

Homologues of *Brachyury* were identified in other vertebrate species including chicken, *Xenopus* and zebrafish (Kispert *et al.* 1995b; Schulte-Merker *et al.* 1992; Smith *et al.* 1991). All proved to have similar expression patterns, and experiments in these species emphasized the importance of *Brachyury* in mesoderm formation. The *no tail* mutation in the zebrafish, for example, proved to encode a fish homologue of *Brachyury* (Schulte-Merker *et al.* 1994), and misexpression of *Xenopus Brachyury* (*Xbra*) in prospective ectoderm of the early gastrula proved to be sufficient to induce ectopic mesoderm of ventral and posterior character (Cunliffe & Smith 1992, 1994; O'Reilly *et al.* 1995). *Brachyury* is therefore both necessary

* Author for correspondence (jim@nimr.mrc.ac.uk).




Figure 1. Comparison of the expression patterns of *Xwnt11* and *Xbra* at the early gastrula stage. (a) *In situ* hybridization showing expression of *Xwnt11* in the early gastrula. (b, c) Flat-mount preparations showing *Xwnt11* RNA expression (blue), revealed by *in situ* hybridization, and *Xbra* protein (brown) revealed by antibody staining. (b) View showing whole marginal zone (prospective mesoderm). (c) Dorsal view at higher power. (d) Ventral view at higher power.




Figure 2. Specificity of dominant-negative *Xwnt11*. (a) Schematic illustrations of (i) wild-type *Xwnt11* and (ii) *dn-wnt11*, which carries a C-terminal truncation. (b–d) *Xenopus* embryos were injected with 2pg *Xwnt8* RNA either alone (b) or in the presence of a 20-fold excess of *dn-wnt8* RNA (c) or *dn-wnt11* RNA (d). Note that axis duplication caused by *Xwnt8* is inhibited by *dn-wnt8* but not by *dn-wnt11*. (e) A hyperphosphorylated form of myc-Dsh (triangle) present in uninjected animal caps is downregulated by *dn-wnt11* and rescued by overexpression of *Xwnt11* and *Xwnt5A* RNA.

and sufficient for the formation at least of posterior mesoderm.

Investigation of the Brachyury protein showed that the N-terminal half of the protein has sequence-specific DNA-binding activity, with polymerase chain reaction-based binding-site selection experiments identifying the partially palindromic sequence T[G/C]ACACCTAGG TGTGAAATT (Kispert & Herrmann 1993). Crystallographic studies indicated that Brachyury binds such a sequence as a dimer, and that it binds in a novel manner in which a C-terminal helix becomes embedded into an enlarged minor groove of DNA without causing bending

(Muller & Herrmann 1997). More recent experiments, however, show that Brachyury can also bind a half-palindromic sequence such as [T/C]TTCACACCT (Casey *et al.* 1998, 1999; Tada *et al.* 1998), and it will be interesting to investigate the mode of Brachyury binding under these conditions.

The C-terminal half of Brachyury functions as a transcription activator (Conlon *et al.* 1996; Kispert *et al.* 1995a), and experiments in *Xenopus* show that the main function of Brachyury is indeed to activate the expression of downstream genes; expression of an interfering construct in which the activation domain of *Xbra* is

replaced by the repressor domain of *Drosophila* Engrailed (Xbra-En^R) causes embryos to lose posterior structures and to exhibit deficiencies in notochord differentiation (Conlon *et al.* 1996). This phenotype resembles that of *Brachyury* and *no tail* homozygous mutant embryos.

We now know that *Brachyury* is the founder member of a family of proteins that share sequence homology with the N-terminal DNA-binding domain described above—the so-called ‘T-box’ (reviewed by Papaioannou & Silver 1998; Smith 1997, 1999). The first protein to be recognized as being homologous to *Brachyury* was *Drosophila* optomotor-blind (Pflugfelder *et al.* 1992); since then the family has expanded to include at least seven T-box proteins in the mouse (not including *Brachyury* itself) as well as members in vertebrates, ascidians, sea urchin, *Drosophila* and *Caenorhabditis elegans*. The genes are expressed at various stages of development and in various tissues, and they have diverse functions. *Tbx5*, for example, plays a role in heart and forelimb development, while *Tbx4* regulates hindlimb development (Basson *et al.* 1997; Li *et al.* 1997; Logan & Tabin 1999; Rodriguez-Esteban *et al.* 1999; Takeuchi *et al.* 1999).

3. BRACHYURY IS REQUIRED FOR NORMAL GASTRULATION MOVEMENTS

In mouse, fish and frog embryos, lack of *Brachyury* function results in loss of posterior mesodermal tissues and lack of a properly differentiated notochord. At least in mouse and *Xenopus* embryos, this phenotype is presaged by abnormal gastrulation movements. In the mouse, for example, morphogenesis of the primitive streak and notochord is abnormal and there is a failure of the axis to elongate properly. The generation of chimeras containing both wild-type cells and cells lacking *Brachyury* function reveals that the mutant cells accumulate in the primitive streak during gastrulation, leading to their eventual accumulation in the tailbud (Wilson *et al.* 1995). This suggests that one role of *Brachyury* might be to alter the adhesion properties of cells as they pass through the primitive streak.

Similar defects in gastrulation movements are observed in *Xenopus* embryos in which *Brachyury* function is inhibited by the interfering construct Xbra-En^R. In particular, the blastopore is misshapen and slow to close, as if convergent extension movements are compromised (Conlon *et al.* 1996). The clearest illustration of the requirement for *Brachyury* during gastrulation, however, comes from studies on *Xenopus* animal pole regions. The mesoderm of the amphibian embryo arises through an inductive interaction in which blastomeres of the vegetal hemisphere of the embryo act on overlying equatorial cells (reviewed by Harland & Gerhart 1997). Candidates for the mesoderm-inducing signals include members of the transforming growth factor- β family of growth and differentiation factors, such as activin, *derrière* and the nodal-related genes (Harland & Gerhart 1997; Osada & Wright 1999; Sun *et al.* 1999). For example, treatment of prospective ectodermal tissue of the animal pole region with activin causes those cells to form mesodermal cell types. An earlier response to activin, however, is that the animal pole cells undergo coordinated convergent extension movements such that the isolated tissue extends in a

dramatic fashion (Symes & Smith 1987). Untreated animal pole regions, by contrast, form spheres of ectodermal cells. The elongation of animal pole regions in response to activin provides a powerful model system for the analysis of gastrulation and convergent extension, and significantly, we find that this elongation does not occur in animal caps derived from embryos injected with RNA encoding Xbra-En^R (Conlon & Smith 1999). Together with the phenotypes of intact *Xenopus* embryos expressing Xbra-En^R, in which gastrulation is impaired, and the results obtained with mouse embryos, these experiments indicate that *Brachyury* function is essential for gastrulation movements and in particular for convergent extension. Since *Brachyury* functions as a transcription activator, it is likely that its target genes include some which are required for convergent extension.

4. A SCREEN FOR BRACHYURY TARGETS IDENTIFIES *Xwnt11*

In an effort to identify the target genes of *Brachyury*, we have screened cDNA libraries in which *Brachyury*-inducible genes are highly enriched (Saka *et al.* 2000; Tada *et al.* 1998). Among other genes, this screen has allowed us to isolate *Xwnt11*, which was originally identified as a maternally expressed *Wnt* gene whose transcripts are restricted to the vegetal hemisphere of the oocyte and early embryo (Ku & Melton 1993). Zygotic expression of *Xwnt11* proved to resemble closely that of *Xbra*, with expression commencing throughout the marginal zone and persisting in posterior circumblastoporal tissue throughout gastrula stages to the early neurula stage (figure 1) (Saka *et al.* 2000; Tada & Smith 2000).

Two experiments suggest that *Xwnt11* is a direct target of Xbra and that expression of *Xwnt11* in the early embryo requires Xbra function. First, induction of *Xwnt11* by the hormone-inducible construct *Xbra-GR* can occur in the presence of cycloheximide, an inhibitor of protein synthesis (Saka *et al.* 2000). This indicates that induction of *Xwnt11* by Xbra does not require intervening protein synthesis, suggesting that Xbra acts directly on the *Xwnt11* promoter. This issue is now under investigation. Second, expression of the interfering construct Xbra-En^R causes an almost complete downregulation of *Xwnt11* in the *Xenopus* embryo, arguing that Xbra function is required for expression of *Xwnt11* (Tada & Smith 2000).

5. *Xwnt11*, LIKE *Xbra*, IS REQUIRED FOR GASTRULATION MOVEMENTS

(a) *Specificity of a dominant-negative Wnt11 construct*

To investigate the function of *Xwnt11* during gastrulation, we constructed a C-terminally truncated form of the protein (dn-wnt11) which, by analogy with a similar *Xwnt8* construct (Hoppler *et al.* 1996), might be expected to act in a dominant-negative fashion (figure 2a). Wnt11 and Wnt8 belong to different classes of the Wnt family (Du *et al.* 1995), and it is important to demonstrate first that our dn-wnt11 construct does indeed inhibit the function of Wnt11 (and perhaps other members of the same class such as Wnt5) and second that it has no effect on the activity of the Wnt8 class.




Figure 3. Xwnt11, like Xbra, is required for gastrulation movements in *Xenopus*. (*a–c*) *Xenopus* embryos at the late gastrula–early neurula stage following injection with RNA encoding (*a*) β -galactosidase as a control, (*b*) Xbra-En^R, and (*c*) dn-wnt11. Note the disruption of gastrulation in (*b*) and (*c*). (*d–f*) *Xenopus* embryos at tadpole stages after injection with RNA encoding (*d*) β -galactosidase, (*e*) Xbra-En^R, and (*f*) dn-wnt11. Note loss of posterior structures in (*e*) and (*f*). (*g–i*) Activin-treated animal pole regions derived from uninjected embryos (*g*) or embryos injected with RNA encoding Xbra-En^R (*h*) or dn-wnt11 (*i*). Note inhibition of elongation in (*h*) and (*i*).

To confirm that dn-wnt11 does indeed inhibit the function of wild-type members of this Wnt class, we isolated animal caps from embryos expressing a myc-tagged form of Dishevelled (Dsh), for review see Boutros & Mlodzik 1999). Western blotting revealed the presence of a hyperphosphorylated form of Dsh in such animal caps, which may be due to high levels of signalling by Xwnt5A, which is expressed in this region (Morgan *et al.* 1999). This hyperphosphorylated form of Dsh was downregulated by dn-wnt11, but could be ‘rescued’ by co-expression of wild-type Xwnt11 or Xwnt5A (figure 2*e*). Together, these experiments indicate that dn-wnt11 inhibits the function of the class of Wnts which includes Wnt5A and Wnt11.

By contrast, dn-wnt11 appears not to inhibit the function of the other class of Wnts, which includes Wnt1 and Wnt8. This was examined using an axis-induction assay, in which injection of RNA encoding Xwnt8 into the *Xenopus* egg causes complete axis duplication (McMahon & Moon 1989; Smith & Harland 1991; Sokol *et al.* 1991). This duplication can be inhibited by dominant-negative Xwnt8, but not by similar concentrations of dn-wnt11 (figure 2*b–d*).

Together, these experiments suggest that dn-wnt11 inhibits the function of Wnt11 itself, and of Wnt5A, but does not affect the action of Wnt1 or Wnt8. With this information in hand, we went on to investigate the effects of dn-wnt11 on gastrulation.

(b) dn-wnt11 blocks gastrulation movements

Expression of dn-wnt11 throughout the *Xenopus* embryo caused embryos to lack posterior structures, but differentiation of muscle and notochord in anterior regions

occurred normally (figure 3*d–f* and data not shown). At earlier stages, expression of dn-wnt11 resulted in delayed and incomplete gastrulation movements, reminiscent of those caused by Xbra-En^R (figure 3*a–c*). However, the expression of a variety of mesodermal markers, including Xbra, gooseoid and Myf-5, were unaffected, suggesting that dn-wnt11 inhibits gastrulation without affecting mesodermal specification or differentiation (Tada & Smith 2000).

These conclusions were confirmed by experiments in activin-treated animal caps; dn-wnt11 prevented activin-induced elongation of isolated animal pole regions (figure 3*g–i*), but had no effect on the ability of activin to induce expression of the mesodermal markers Xbra, gooseoid, Xwnt8 or Bix1 (Tada & Smith 2000). Thus, inhibition of signalling by the class of Wnt signalling molecules that includes Wnt11 prevents normal gastrulation movements, but does not affect mesodermal specification.

6. Xwnt11 DOES NOT ACT THROUGH THE CANONICAL Wnt SIGNALLING PATHWAY

What is the intracellular signalling pathway employed by Xwnt11? The ability of Xwnt11 to regulate the hyperphosphorylation of Dsh suggests that it might act through the canonical Wnt pathway involving Dsh, GSK-3 and β -catenin (Cadigan & Nusse 1997), and in support of this idea we find that Dsh can ‘rescue’ the inhibition of activin-induced elongation of animal caps that is caused by dn-wnt11 (figure 4*e*). To investigate this question further, we made use of Δ N-Tcf3, a dominant-negative version of Tcf3 (T-cell factor 3) which can block the





Figure 4. Mapping of Dsh domains required for convergent extension movements. (a) Different Dsh constructs used in our experiments. DN, dominant-negative. Effects of each construct are based on experiments described in this paper, and by Tada & Smith (2000), Axelrod *et al.* (1998), Boutros & Mlodzik (1999) and Boutros *et al.* (1998). (b-e) Dsh and Dsh-ΔN can rescue the inhibition of activin-induced elongation of animal caps caused by *dn-wnt11*. (b) Activin induces elongation of untreated animal caps. (c) Elongation is inhibited by *dn-wnt11*. (d, e) Rescue of elongation by (d) Dsh-ΔN (activin+ *dnwnt11*+ Dsh-ΔN) and (e) Dsh (activin+ *dnwnt11*+ Dsh).

transcriptional function of β -catenin (Molenaar *et al.* 1996). Surprisingly, Δ N-Tcf3 did not block activin-induced elongation of animal caps (Tada & Smith 2000), suggesting that in the regulation of gastrulation, Xwnt11 signalling 'branches' from the canonical Wnt signalling pathway somewhere downstream of Dsh.

(a) Mapping domains of Dsh

The idea that separate signalling pathways derive from Dsh is consistent with genetic evidence in *Drosophila*. The first *dishevelled* mutant to be isolated, *dsh¹*, causes defects in the polarity of bristles on the wing and thorax, a phenotype that gave the mutant its name (Fahmy & Fahmy 1959). Additional genetic screens, however, isolated null alleles of *dsh* the phenotypes of which were completely different, and resembled those of *wingless* (a *Drosophila* Wnt family member) and *armadillo* (*Drosophila* β -catenin) (Nüsslein-Volhard & Wieschaus 1980; Perrimon & Mahowald 1987). Molecular characterization of Dsh

suggests that the different phenotypes are the consequences of mutations occurring in different domains of the protein (Axelrod *et al.* 1998; Boutros & Mlodzik 1999; Boutros *et al.* 1998; and see figure 4a). Thus, the DEP domain of Dsh proves to be essential for polarity signalling, whereas the DIX domain is dispensable. In contrast, the DIX domain is essential for signalling through β -catenin, but the DEP region is not required. Consistent with these conclusions, the original *dsh¹* allele has a single amino-acid substitution in a conserved region of the DEP domain.

(b) Xwnt11 acts through the 'polarity' pathway

We have taken advantage of this information to ask whether Xwnt11 signals through the β -catenin pathway or the polarity pathway. Consistent with the suggestion above that Xwnt11 signalling branches from the canonical Wnt signalling pathway, we find that Dsh constructs lacking just the DIX domain are able to rescue the

inhibition of activin-induced elongation of animal caps that is caused by dn-wnt11 (figure 4d). This result suggests that *Xwnt11* regulates gastrulation through a pathway analogous to that used in polarity signalling in the *Drosophila* embryo. This is discussed below (§7(c, d)).

7. DISCUSSION AND CONCLUSIONS

In an attempt to understand the molecular basis of gastrulation, we have taken advantage of the observation that loss of *Brachyury* function causes a severe disruption of convergent extension. Since *Brachyury* is a transcription activator, and the ability of *Brachyury* to activate transcription is essential for its function, we deduced that amongst its target genes there must be some involved in the regulation of gastrulation. *Xwnt11* proved to be such a gene.

(a) *Xwnt11* as a target of *Brachyury*

Our experiments suggest that *Xwnt11* is a direct target of *Xbra* and that *Xbra* function is essential for expression of *Xwnt11* during early *Xenopus* development. Confirmation that *Xwnt11* is a bona fide target of *Xbra*, however, awaits analysis of the *Xwnt11* 5' regulatory region, where we hope at least to identify *Brachyury* 'half-sites', as are present in the *eFGF* and *Bix4* promoters, both of which are direct targets of T-box genes (Casey *et al.* 1998, 1999; Tada *et al.* 1998).

(b) *Xwnt11* function is required for normal gastrulation movements

The only practical way to study gene function in *Xenopus*, at least for now, is to make use of antisense technology or dominant-negative constructs. To this end we have made a dominant-negative version of *Xwnt11*, and demonstrated that it inhibits the function of the Wnt5A/Wnt11 class of Wnts, but not the Wnt1/Wnt8 class. Expression of dn-wnt11 in the *Xenopus* embryo interferes with gastrulation movements, and it also interferes with the elongation of isolated dorsal marginal zone regions and activin-treated animal caps.

These experiments are consistent with the idea that *Xwnt11* function is required for normal gastrulation movements, but it is possible, of course, that dn-wnt11 also interferes with the function of other members of the Wnt11 class of Wnts, such as *Xwnt5A*. Of the known members of this class, however, only *Wnt11* is expressed in the mesoderm of the embryo during gastrulation (Du *et al.* 1995; Moon *et al.* 1993; Morgan *et al.* 1999), suggesting that the effects of dn-wnt11 are indeed due to interference of the function of this family member.

(c) *Wnt11* does not signal through the canonical Wnt pathway

Experiments using a dominant-negative version of *Tcf3*, and others using different domains of *Dsh*, suggest that *Xwnt11* does not signal through the canonical Wnt signalling pathway involving *Dsh*, *GSK-3* and β -catenin. Rather, it may use a pathway more reminiscent of that used in polarity signalling during *Drosophila* development. This pathway is less well understood than the canonical pathway, but involves small GTPases such as *RhoA* and *Rac* followed by the activation of *JNK/SAPK*-like kinases

(Boutros & Mlodzik 1999; Boutros *et al.* 1998; Strutt *et al.* 1997). In this regard it is interesting that these components have been implicated in migration and cell shape changes occurring during gastrulation and dorsal closure in *Drosophila* (Barrett *et al.* 1997; Noselli & Agnes 1999), and it is an intriguing possibility that the same intracellular signalling pathways regulate gastrulation in both vertebrates and invertebrates.

(d) *The future*

Our work raises more questions than it answers. One task, as suggested above, is to investigate further the *Xwnt11* signalling pathway. As well as the possible involvement of *Rho*- and *Rac*-like molecules, it is important to note that *Xwnt5A*, but not *Xwnt8*, can cause the release of intracellular calcium via G-protein-linked phosphatidylinositol signalling (Slusarski *et al.* 1997), and thereby activate protein kinase C (Sheldahl *et al.* 1999). It is possible that *Xwnt11* employs a similar pathway. One should also remember that thinking of Wnt signalling pathways as being linear may be a mistake—a model based on a signalling network may be more appropriate (Martinez Arias *et al.* 1999).

At the embryological level it may be helpful to extend the analogy with polarity signalling in *Drosophila* to look at the cell polarity during *Xenopus* gastrulation, and to ask if polarity is disrupted by inhibition of *Xwnt11* signalling. Along these lines, it may be significant that gastrulation movements in *Xenopus* are not only disrupted by inhibiting signalling by this class of Wnt—overexpression of *Xwnt5A* also blocks convergent extension (Moon *et al.* 1993). This may reflect the fact that cell polarity can in principle be disrupted both by removing localized Wnt signalling and also by completely flooding the system. We note that in *Drosophila*, overexpression of the Wnt receptor frizzled 1 causes a polarity phenotype similar to that observed in a loss-of-function mutation (Krasnow & Adler 1994).

REFERENCES

- Axelrod, J. D., Miller, J. R., Shulman, J. M., Moon, R. T. & Perrimon, N. 1998 Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and Wingless signaling pathways. *Genes Dev.* **12**, 2610–2622.
- Barrett, K., Leptin, M. & Settleman, J. 1997 The *Rho* GTPase and a putative *Rho*GEF mediate a signaling pathway for the cell shape changes in *Drosophila* gastrulation. *Cell* **91**, 905–915.
- Basson, C. T. (and 13 others) 1997 Mutations in human cause limb and cardiac malformation in Holt–Oram syndrome. *Nature Genet.* **15**, 30–35.
- Boutros, M. & Mlodzik, M. 1999 Dishevelled: at the crossroads of divergent intracellular signaling pathways. *Mech. Dev.* **83**, 27–37.
- Boutros, M., Paricio, N., Strutt, D. I. & Mlodzik, M. 1998 Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and *wingless* signaling. *Cell* **94**, 109–118.
- Cadigan, K. M. & Nusse, R. 1997 Wnt signaling: a common theme in animal development. *Genes Dev.* **11**, 3286–3305.
- Casey, E. S., O'Reilly, M. A., Conlon, F. L. & Smith, J. C. 1998 The T-box transcription factor *Brachyury* regulates expression of *eFGF* through binding to a non-palindromic response element. *Development* **125**, 3887–3894.
- Casey, E. S., Tada, M., Fairclough, L., Wylie, C. C., Heasman, J. & Smith, J. C. 1999 *Bix4* is activated by *VegT* and mediates

- endoderm formation in *Xenopus* development. *Development* **126**, 4193–4200.
- Chesley, P. 1935 Development of the short-tailed mutant in the house mouse. *J. Exp. Zool.* **70**, 429–459.
- Conlon, F. L. & Smith, J. C. 1999 Interference with *Brachyury* function inhibits convergent extension, causes apoptosis, and reveals separate requirements in the FGF and activin signaling pathways. *Dev. Biol.* **213**, 85–100.
- Conlon, F. L., Sedgwick, S. G., Weston, K. M. & Smith, J. C. 1996 Inhibition of *Xbra* transcription activation causes defects in mesodermal patterning and reveals autoregulation of *Xbra* in dorsal mesoderm. *Development* **122**, 2427–2435.
- Cunliffe, V. & Smith, J. C. 1992 Ectopic mesoderm formation in *Xenopus* embryos caused by widespread expression of a *Brachyury* homologue. *Nature* **358**, 427–430.
- Cunliffe, V. & Smith, J. C. 1994 Specification of mesodermal pattern in *Xenopus laevis* by interactions between *Brachyury*, *noggin* and *Xwnt-8*. *EMBO J.* **13**, 349–359.
- Dobrovolskaia-Zavadskaia, N. 1927 Sur la mortification spontanée de la queue chez la souris nouveau-née et sur l'existence d'un caractère héréditaire 'non-viable'. *C. R. Soc. Biol.* **97**, 114–116.
- Du, J. S., Purcell, S. M., Christian, J. L., McGrew, L. L. & Moon, R. T. 1995 Identification of distinct classes and functional domains of Wnts through expression of wild-type and chimeric proteins in *Xenopus* embryos. *Mol. Cell. Biol.* **15**, 2625–2634.
- Fahmy, O. G. & Fahmy, M. 1959 New mutants report. *Drosophila Info. Serv.* **33**, 83–94.
- Gluecksohn-Schoenheimer, S. 1944 The development of normal and homozygous *brachy* (*T/T*) mouse embryos in the extra-embryonic coelom of the chick. *Proc. Natl Acad. Sci. USA* **30**, 134–140.
- Harland, R. & Gerhart, J. 1997 Formation and function of Spemann's organizer. *A. Rev. Cell Devl Biol.* **13**, 611–667.
- Herrmann, B. G., Labeit, S., Poutska, A., King, T. R. & Leirach, H. 1990 Cloning of the T gene required in mesoderm formation in the mouse. *Nature* **343**, 617–622.
- Hoppler, S., Brown, J. D. & Moon, R. T. 1996 Expression of a dominant-negative Wnt blocks induction of MyoD in *Xenopus* embryos. *Genes Dev.* **10**, 2805–2817.
- Kispert, A. & Herrmann, B. G. 1993 The *Brachyury* gene encodes a novel DNA binding protein. *EMBO J.* **12**, 3211–3220.
- Kispert, A., Korschorz, B. & Herrmann, B. G. 1995a The T protein encoded by *Brachyury* is a tissue-specific transcription factor. *EMBO J.* **14**, 4763–4772.
- Kispert, A., Ortner, H., Cooke, J. & Herrmann, B. G. 1995b The chick *Brachyury* gene: developmental expression pattern and response to axial induction by localized activin. *Devl Biol.* **168**, 406–415.
- Krasnow, R. E. & Adler, P. N. 1994 A single frizzled protein has a dual function in tissue polarity. *Development* **120**, 1883–1893.
- Ku, M. & Melton, D. A. 1993 *Xwnt-II*, a maternally expressed *Xenopus wnt* gene. *Development* **119**, 1161–1173.
- Li, Q. Y. (and 16 others) 1997 Holt–Oram syndrome is caused by mutations in TBX5, a member of the *Brachyury* (T) gene family. *Nature Genet.* **15**, 21–29.
- Logan, M. & Tabin, C. J. 1999 Role of Pitx1 upstream of *Tbx4* in specification of hindlimb identity. *Science* **283**, 1736–1739.
- McMahon, A. P. & Moon, R. T. 1989 Ectopic expression of the proto-oncogene int-1 in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* **58**, 1075–1084.
- Martinez Arias, A., Brown, A. M. & Brennan, K. 1999 Wnt signalling: pathway or network? *Curr. Opin. Genet. Dev.* **9**, 447–454.
- Molenaar, M., Van de Wetering, M., Oosterwegel, M., Peter-Maduro, J., Godsave, S., Korinek, V., Roose, J., Destree, O. & Clevers, H. 1996 XTcf-3 transcription factor mediates β -catenin-induced axis formation in *Xenopus* embryos. *Cell* **86**, 391–399.
- Moon, R. T., Campbell, R. M., Christian, J. L., McGrew, L. L., Shih, J. & Fraser, S. 1993 Xwnt-5A: a maternal Wnt that affects morphogenetic movements after overexpression in embryos of *Xenopus laevis*. *Development* **119**, 97–111.
- Morgan, R., Hooiveld, M. H. W., In der Rieden, P. & Durston, A. J. 1999 A conserved 30 base pair element in the *Wnt-5a* promoter is sufficient both to drive its early embryonic expression and to mediate its repression by *otx2*. *Mech. Dev.* **85**, 97–102.
- Muller, C. W. & Herrmann, B. G. 1997 Crystallographic structure of the T domain–DNA complex of the *Brachyury* transcription factor. *Nature* **389**, 884–888.
- Noselli, S. & Agnes, F. 1999 Roles of the JNK signaling pathway in *Drosophila* morphogenesis. *Curr. Opin. Genet. Dev.* **9**, 466–472.
- Nüsslein-Volhard, C. & Wieschaus, E. 1980 Mutations affecting segment number and polarity in *Drosophila*. *Nature* **287**, 795–801.
- O'Reilly, M.-A. J., Smith, J. C. & Cunliffe, V. 1995 Patterning of the mesoderm in *Xenopus*: dose-dependent and synergistic effects of *Brachyury* and *Pintallavis*. *Development* **121**, 1351–1359.
- Osada, S. & Wright, C. V. E. 1999 *Xenopus nodal*-related signal is essential for mesendodermal patterning during early embryogenesis. *Development* **126**, 3229–3240.
- Papaioannou, V. E. 1997 T-box family reunion. *Trends Genet.* **13**, 212–213.
- Papaioannou, V. E. & Silver, L. M. 1998 The T-box gene family. *BioEssays* **20**, 9–19.
- Perrimon, N. & Mahowald, A. P. 1987 Multiple functions of segment polarity genes in *Drosophila*. *Devl Biol.* **119**, 587–600.
- Pflugfelder, G. O., Roth, H. & Poeck, B. 1992 A homology domain shared between I optomotor-blind and mouse *Brachyury* is involved in DNA binding. *Biochem. Biophys. Res. Commun.* **186**, 918–925.
- Rodriguez-Esteban, C., Tsukui, T., Yonei, S., Magallon, J., Tamura, K. & Izpisua Belmonte, J. C. 1999 The T-box genes *Tbx4* and *Tbx5* regulate limb outgrowth and identity. *Nature* **398**, 814–818.
- Saka, Y., Tada, M. & Smith, J. C. 2000 A screen for targets of the *Xenopus* T-box gene *Xbra*. *Mech. Dev.* (In the press.)
- Schulte-Merker, S., Ho, R. K., Herrmann, B. G. & Nüsslein-Volhard, C. 1992 The protein product of the zebrafish homologue of the mouse T gene is expressed in nuclei of the germ ring and the notochord of the early embryo. *Development* **116**, 1021–1032.
- Schulte-Merker, S., Van Eeden, F. M., Halpern, M. E., Kimmel, C. B. & Nüsslein-Volhard, C. 1994 No tail (ntl) is the zebrafish homologue of the mouse T (*Brachyury*) gene. *Development* **120**, 1009–1015.
- Sheldahl, L. C., Park, M., Malbon, C. C. & Moon, R. T. 1999 Protein kinase C is differentially stimulated by Wnt and Frizzled homologs in a G-protein-dependent manner. *Curr. Biol.* **9**, 695–698.
- Slusarski, D. C., Corces, V. G. & Moon, R. T. 1997 Interaction of Wnt and a Frizzled homologue triggers G-protein-linked phosphatidylinositol signalling. *Nature* **390**, 410–413.
- Smith, J. 1997 *Brachyury* and the T-box genes. *Curr. Opin. Genet. Dev.* **7**, 474–480.
- Smith, J. 1999 T-box genes: what they do and how they do it. *Trends Genet.* **15**, 154–158.
- Smith, J. C., Price, B. M. J., Green, J. B. A., Weigel, D. & Herrmann, B. G. 1991 Expression of a *Xenopus* homolog of *Brachyury* (*T*) is an immediate-early response to mesoderm induction. *Cell* **67**, 79–87.

- Smith, W. C. & Harland, R. M. 1991 Injected *Xwnt-8* RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* **67**, 753–765.
- Sokol, S., Christian, J. L., Moon, R. T. & Melton, D. A. 1991 Injected *wnt* RNA induces a complete body axis in *Xenopus* embryos. *Cell* **67**, 741–752.
- Strutt, D. I., Weber, U. & Mlodzik, M. 1997 The role of RhoA in tissue polarity and Frizzled signalling. *Nature* **387**, 292–295.
- Sun, B. I., Bush, S. M., Collins-Racie, L. A., LaVallie, E. R., DiBlasio-Smith, E. A., Wolfman, N. M., McCoy, J. M. & Sive, H. L. 1999 *derrière*: a TGF-beta family member required for posterior development in *Xenopus*. *Development* **126**, 1467–1482.
- Symes, K. & Smith, J. C. 1987 Gastrulation movements provide an early marker of mesoderm induction in *Xenopus*. *Development* **101**, 339–349.
- Tada, M. & Smith, J. C. 2000 *Xwnt11* is a target of *Xenopus* Brachyury: regulation of gastrulation movements via Dishevelled, but not through the canonical Wnt pathway. *Development*. (In the press.)
- Tada, M., Casey, E., Fairclough, L. & Smith, J. C. 1998 *Bixl*, a direct target of *Xenopus* T-box genes, causes formation of ventral mesoderm and endoderm. *Development* **125**, 3997–4006.
- Takeuchi, J. K., Koshiba-Takeuchi, K., Matsumoto, K., Vogel-Hopker, A., Naitoh-Matsuo, M., Ogura, K., Takahashi, N., Yasuda, K. & Ogura, T. 1999 *Tbx5* and *Tbx4* genes determine the wing/leg identity of limb buds. *Nature* **398**, 810–814.
- Wilkinson, D. G., Bhatt, S. & Herrmann, B. G. 1990 Expression pattern of the mouse *T* gene and its role in mesoderm formation. *Nature* **343**, 657–659.
- Wilson, V., Manson, L., Skarnes, W. C. & Beddington, R. S. P. 1995 The *T* gene is necessary for normal mesodermal morphogenetic cell movements during gastrulation. *Development* **121**, 877–886.