

I have computed this Table so far, that the Reader may see in what manner this Method Approximates; this whole Work, as it appears, costing a little more than three Hours time.

V. *Proprietates quædam simplices Sectionum Conicarum ex natura Focorum deductæ; cum Theoremate generali de Viribus Centripetis; quorum ope Lex Virium Centripetarum ad Focos Sectionum tendentium, Velocitates Corporum in illis revolvantium, & Descriptio Orbium facillime determinantur. Per Abr. de Moivre. R. S. Soc.*

Sit  $DE$  Axis Transversus Ellipseos,  $AO$  Axis alter, &  $C$  centrum Sectionis. Sit  $P$  punctum quodvis in circumferentia ejus;  $PQ$  Tangens curvæ ad  $P$ , occurrens Axi Transverso ad  $Q$ ; puncta  $S, F$  Foci;  $CP$ ,  $CK$  semidiametri Conjugatæ;  $PH$  Semilatus rectum ad diametrum  $PC$ ;  $PG$  normalis ad Tangentem, cui occurrat  $HG$ , perpendicularis ipsi  $PC$ , in puncto  $G$ , ut fiat  $PG$  radius Curvaturæ Ellipseos in puncto  $P$ : sint etiam  $ST$ ,  $CR$ ,  $FP$  perpendiculares in Tangentem  $PQ$  demissæ: Jungatur  $SO$ , & demittatur in Axem normalis  $PL$ . His positis, Dico quod,

I. *Rectangulum sub distantis ab utroque Ellipseos Foco, five  $SP \times PF$  æquale est quadrato Semidiametri  $CK$ .*

*Demonstratio.*

$$PSq = PCq + CSq - 2CS \times CL \text{ per 13. II. Elem.}$$

$$PFq = PCq + CSq + 2CS \times CL \text{ per 12. II. Elem.}$$

$$\text{Unde } PSq + PFq = 2PCq + 2CSq.$$

$$\text{Jam } PS + PF = DE = 2CD; \text{ ac propterea}$$

$$PSq + PFq + 2PS \times PF = 4CDq. \quad \text{Quare}$$

Quare transponendo,  $2PS \times PF = 4CDq - 2PCq - 2CSq$ .

Ac Dimidiando  $PS \times PF = 2CDq - PCq - CSq$ .

Est autem  $CS \text{ quad.} = CD \text{ quad.} - CO \text{ quad.}$ , atque adeo

$PS \times PF = 2CDq + COq - PCq$ .

Sed  $CDq + COq = PCq + CKq$ . per 12. VII. Conic. Apollonii.

Quare  $PS \times PF = CKq$ . Q. E. D.

II. Distantia à Foco  $SP$  est ad perpendicularem in Tangentem demissam, ut Semidiameter Conjugata  $CK$  ad Semiaxem minorem  $CO$ .

*Demonstratio.*

Ob similia Triangula  $SPT$ ,  $FPV$ , erit  $PS : PF :: ST : FV$ ; ac componendo  $PS + PF$  erit ad  $ST + FV$ , & earundem dimidia  $CD$  ad  $CR$ , ut  $PS$  ad  $ST$ . Unde  $CD \times CK$  erit ad  $CR \times CK$  ut  $PS$  ad  $ST$ . Sed  $CR \times CK$  æquale est rectangulo sub Semiaxibus  $CD$  in  $CO$ , per 31. VII. Conic. Proinde  $PS$  est ad  $ST$  ut  $CD$  in  $CK$  ad  $CD \times CO$ , sive ut  $CK$  ad  $CO$ . Ac pari argumento demonstrabitur  $PF$  esse ad  $FV$  in eadem ratione. Q. E. D.

III. In eadem etiam est ratione Semiaxis Transversus  $CD$  ad normalem à centro  $C$  ad Tangentem demissam, sive ad  $CR$ .

Etenim cum rectangulum  $CR \times CK$  æquale sit rectangulo  $CD \times CO$ , uti jam dictum est, erit  $\alpha\nu\epsilon\lambda\omicron\gamma\omicron\nu$   $CD$  ad  $CR$  ut  $CK$  ad  $CO$ . Q. E. D.

IV. Semidiameter quævis  $PC$  est ad distantiam puncti  $P$  à foco  $S$ , sive ad  $SP$ , ut distantia ab altero Foco  $FP$  ad dimidium lateris recti ad Verticem  $P$  pertinentis, sive ad  $PH$ .

Hoc autem manifestum est ob Propr. I. cum nempe quadratum ex  $CK$  æquale sit rectangulo sub  $SP \times PF$ .

V. Rectangulum Semiaxium  $CD \times CO$  est ad quadratum semidiametri conjugatæ  $CK$ , ut  $CK$  ad Radium Curvature in puncto  $P$ , sive ad  $PG$ .

Sunt

Sunt enim Triangula  $PCR$ ,  $PGH$  inter se similia,  
unde  $CR$  est ad  $PC$ , ut semilatus rectum  $PH$  ad  $PG$ :  
hoc est, per præmissam Proprietatem III,  $\frac{CD \times CO}{CK} =$   
 $CR$  est ad  $PC$  ut  $\frac{CK^2}{PC} = PH$  ad  $\frac{CK^3}{CD \times CO} = PG$ .  
proinde ἀνάλογον  $CD \times CO : CK^2 :: CK : PG$ . Q.E.D.

## THEOREMA GENERALE I.

*Vis centripeta ad idem punctum S tendens, in Curvis omnibus, est semper proportionalis Quantitati  $\frac{SP}{PG \times ST^3}$*

Hoc Theorema ante plures annos à me investigatum & cum amicis communicatum, propriis demonstrationibus firmavere Geometræ Clarissimi D. J. Bernoullius in *Act. Lipsæ*; D. J. Keillius in harum *Transact.* N. 317. & D. Jac. Hermannus in *Phoronomiâ* suâ pag. 70. quos vide.

Scribendo autem  $CK^3$  pro  $PG$ , per *Prop. V*; &  $\frac{SP}{CK}$  juxta *Prop. II*, pro  $ST$ ; (ob datas scilicet  $CD, CO$ ) erit *Vis centripeta* tendens ad focum Ellipseos  $S$ , semper ut  $\frac{SP \times CK^3}{CK^3 \times ST^3}$ , hoc est ut  $\frac{SP}{SP^3}$  vel  $\frac{1}{SP^2}$ , nempe reciprocè ut quadratum ex  $SP$ . Unde patet quod si Sectio fuerit Ellipsis motu corporis descripta, erit *Vis Centripeta* ut quadratum distantiae à centro Virium reciprocè. Ex his Proprietatibus consequuntur Corollaria nonnulla notatu non indigna.

Coroll. 1. *Velocitas Corporis in Ellipti revolvantis, ad punctum quodlibet P, est ad Velocitatem revolvantis in circulo ad eandem distantiam SP à centro Virium, in subduple ratione distantia ab altero foco PF, ad Semiaxem transversum Sectionis, sive ut media proportionalis inter PF & CD ad CD.*

Est enim velocitas revolventis in Ellipfi ad distantiam  
 $SP_0$  ad Velocitatem revolventis in Circulo vel Ellipfi ad  
 dist.

distantiam Semiaxis  $CD$  vel  $SO$ , ut  $CO$  ad  $ST$ ; hoc est per *Prop. II.* ut  $\sqrt{PF}$  ad  $\sqrt{SP}$ . Velocitas autem revolvantis in Circulo ad distantiam  $CD$  est ad velocitatem revolvantis in Circulo ad distantiam  $SP$ , ut  $\sqrt{SP}$  ad  $\sqrt{CD}$ . Ex æquo igitur, Velocitas revolvantis in Ellipsi ad distantiam  $SP$ , est ad Velocitatem revolvantis in Circulo ad eandem distantiam ut  $\sqrt{PF}$  ad  $\sqrt{CD}$ .

*Coroll. 2. Ex datis Velocitate in Ellipsi, positione Tangentis, & centro Virium seu Foco, facile est determinare Focum alterum.*

Sit enim Velocitas Data  $R$ ; ea autem Velocitas quâ describeretur Circulus ad datam à centro distantiam  $SP$  sit  $Q$ ; ac per *Coroll. præcedens*,  $R$  est ad  $Q$  ut  $\sqrt{PF}$  ad  $\sqrt{CD}$ , adeoque  $QQ$  est ad  $RR$  ut  $CD$  ad  $PF$ , &  $2QQ - RR$  erit ad  $RR$  ut  $SP$  ad  $PF$ : Datur autem  $SP$ ; data est igitur  $PF$  magnitudine. Datur etiam positione, ob angulum  $VPF$  angulo  $SPT$  æqualem. Datur igitur punctum  $F$  alter Focorum: Quo invento prout est Sectionem describere.

Si vero  $\frac{1}{2} RR$  majus fuerit quadrato ex  $Q$ ,  $2QQ - RR$  fit quantitas Negativa, & loco Ellipseos Trajectoria describenda in Hyperbolam transit. Eritque  $RR - 2QQ$  ad  $RR$  ut  $SP$  ad  $PF$  distantiam alterius Foci, ad alterum Tangentis latus ponendam, ut habeatur Focus  $F$ . Proprietates autem omnes quas in Ellipsi demonstravimus; mutatis mutandis etiam Hyperbolæ competunt. *Fig. II.*

Quod si acciderit  $QQ$  æquale esse dimidio quadrati ex  $R$ ; evanescente quantitate  $2QQ - RR = 0$ , quarta proportionalis  $PF$  fit infinita: proinde Trajectoria describenda Parabolica est, Foco scilicet altero in infinitum abeunte. Axis autem Trajectoriæ positione datur; est enim ipsi  $PF$  parallelus, existente scilicet angulo  $FPV$  angulo dato  $SPT$  æquali.

*Coroll. 3. Velocitas revolvantis in data Sectione Conica ad distantiam  $SP$  est ad Velocitatem ejusdem ad distantiam aliquam  $SX$ , ut media proportionalis inter  $FP$  &  $SX$  ad mediam proportionalem inter  $SP$  &  $FX$ .* Velo-

Velocitas enim in  $P$  est ut  $\sqrt{\frac{FP}{SP}}$  (per propr. II.) & per eandem, Velocitas in  $X$  est ut  $\sqrt{\frac{FX}{SX}}$ . Unde manifesta est propositio.

Coroll. 4. *Ratio etiam Velocitatum duorum Corporum in eodem Systemate, sed in datis Confectionibus diversis, revolventium, datis utriusque à communi Orbium Foco distantis, ope Corollarii 1<sup>mi</sup>. statim obtinebitur.*

Cum enim Velocitas corporis in  $P$  sit ad Velocitatem in Circulo ad eandem distantiam  $SP$ , ut  $\sqrt{PF}$  ad  $\sqrt{CD}$ ; & in alia supposita Confectione, cujus Semiaxis  $cd$  & Foci  $S, f$ , ad distantiam  $Sp$  Velocitates illæ sint ut  $\sqrt{pf}$  ad  $\sqrt{cd}$ : Velocitas autem revolventis in circulo ad distantiam  $SP$  sit ad Velocitatem in Circulo ad distantiam  $Sp$  ut  $\sqrt{Sp}$  ad  $\sqrt{SP}$ ; Compositis rationibus, erit Velocitas in  $P$  ad Velocitatem in  $p$ , ut  $\sqrt{PF} \times cd \times Sp$  ad  $\sqrt{pf \times CD \times SP}$ . Quod si Sectio illa altera fuerit Parabola, erunt  $cd, pf$  infinitæ, sed in ratione 1 ad 2; proinde ratio Velocitatum erit ut  $\sqrt{PF} \times Sp$  ad  $\sqrt{2CD \times SP}$ .

Coroll. 5. *Quod si in Hyperbola punctum  $P$  abeat in infinitum, ex præcedentibus manifestum est, Velocitatem ultimam ac minimam, qua cum corpus in æternum ascenderet, æqualem esse ei qua, ad distantiam  $CD$  Semiaxi transverso æqualem, Circulum describeret.*

Coroll. 6. *Ex data distantia à Foco, datur quoque Positio Tangentis, sive angulus  $SPT$ , sub distantia  $SP$  & Tangente  $PT$  contentus.*

Est enim (per propr. II.)  $PS$  ad  $ST$  ut  $CK$  ad  $CO$  sive ut  $\sqrt{SP} \times PF$  ad  $CO$ , atque ita Radius ad Sinum anguli  $SPT$ . At in Ellipsis Circulis affinis præstaret angulum  $PST$ , ejusdem complementum ad quadrantem, inquirere: Hujus autem Sinus est ad Radium ut  $\sqrt{SP \times PF} - CO$  q ad  $\sqrt{SP \times PF}$ .

Coroll.

*Coroll. 7. Atque hinc consequuntur Velocitates quibuscum distantia  $SP$  crescunt vel decrescunt.*

Nam cum, ex Corollario præcedente,  $\sqrt{SP \times PF}$  sit ad  $\sqrt{SP \times PF - CO} q$  ut Radius ad sinum anguli  $PST$ , ac in eadem sit ratione Velocitas Corporis in  $P$  ad Velocitatem momenti ipsius  $SP$ ; Velocitas autem illa in  $P$  sit (per propr. II.) ut  $\sqrt{\frac{PF}{SP}}$ ; elisis superfluis, erit  $\sqrt{\frac{SP \times PF - CO}{SP} q}$  Velocitati, qua crescit vel decrescit distantia  $SP$ , semper proportionalis.

## THEOREMA GENERALE II.

*In omni Trajectoria Curvilinea Velocitates angulares circa centrum Virium sunt reciproce proportionales quadratis distantiarum à centro.*

Nam ob Sectorum minimorum Areas æquales, arcus angulis minimis subtensi sive Bases, sunt reciproce ut Radii: Anguli autem minimi quibus Bases æquales subtenduntur sunt etiam reciproce ut Radii. Proinde anguli Sectorum minimorum Area æqualium, sunt inter se reciproce in dupla ratione Radiorum, sive ut quadrata distantiarum.

*Coroll. 8. Hinc Velocitates angulares revolvantium in diversis Ellipsis datis comparantur inter se.*

Velocitates enim angulares quibuscum ad distantias Semiaxibus Transversis æquales circuli describerentur, sunt reciproce in ratione sesquialtera Axium, sive ut  $\frac{1}{CD \sqrt{CD}}$ .

Velocitates autem angulares has medias habent Corpora revolvantia, cum quadrata distantiarum æquantur rectangulis sub semiaxibus Ellipseôn. Ideo (per Theor. II.) erit  $SP q$  ad  $CD \times CO$  ut  $\frac{1}{CD \sqrt{CD}}$  ad  $\frac{CO}{SP q \times \sqrt{CD}}$ : quæ quidem Quantitas est ut Velocitas anguli ad centrum  $S$ , motu rectæ  $SP$ , tempore quam minimo dato, descripti.

*Coroll. 9. Velocitas angularis qua circumgyratur Tangens  $PT$ , sive recta in Tangentem perpendicularis  $ST$ , est ad Velocitatem*

locitatem angularem rectæ  $SP$ , ut Semiaxis transversus  $CD$  ad distantiam ab altero Foco  $PF$ .

*Demonstratio.*

In Fig. III. Sint puncta  $P, p$ , quamproxima inter se; ductisque  $SP, Sp$ , sint  $PT, pt$  duæ Tangentes, ad quas demittantur normales  $ST, St$ ; iisque parallelæ ducantur radii Curvaturæ  $PG, pG$  coeuntes in  $G$ : ac describatur, centro  $S$  & radio  $SP$ , arcus minimus  $PE$  occurrens ipsi  $Sp$  in  $E$ . Manifestum est angulum  $PGp$  æqualem esse angulo  $TSt$ , sive angulari Velocitati normalis  $ST$ . Est autem angulus  $PSp$  angularis velocitas rectæ  $SP$ ; quare angulus  $PGp$  est ad angulum  $PSp$  ut angularis Velocitas ipsius  $ST$  ad angularem velocitatem rectæ  $SP$ ; hoc est, ut  $\frac{Pp}{PG}$  ad  $\frac{PE}{PS}$ . Sed  $Pp \cdot PE :: SP \cdot ST :: CK : CO$

(per propr. II). Hæ igitur Velocitates sunt ut  $\frac{CK}{PG}$  ad  $\frac{CO}{PS}$ .

Pro  $PG$  scribe  $\frac{CK^3}{CD \times CO}$  (per propr. V.) ac  $\frac{CK}{PG}$  fiet  $\frac{CD \times CO}{CKq} = \frac{CD \times CO}{PS \times PF}$ . Hinc  $\frac{CD \times CO}{PS \times PF}$  erit ad  $\frac{CO}{PS}$ , sive, deletis superfluis,  $CD$  ad  $PF$ , ut angulus  $TSt$  ad angulum  $PSp$ , sive Velocitas angularis Tangentis ad angularem Velocitatem distantiae  $SP$ : proinde Velocitas qua circumgyratur Tangens, semper proportionalis est quantitati  $\frac{CO \times \sqrt{CD}}{PF \times SPq}$ .

Pleraque horum Corollariorum ex aliis Conicarum Sectionum Proprietatibus deducta, vel facile deducenda, inveniet Lector in Sect. III. Lib. I. Princip. Nat. Philosophiæ.

F I N I S.

