II. Observations on the Growth of Trees: By Robert Marsham, of Stratton in Norfolk, Efg; communicated by the Rev. Steph. Hales, D. D. F. R. S.

Measures of Trees, taken in April 1743, before they began to shoot; and again in Autumn 1758, after the Tear's Growth was completed. The Measure taken at 5 Feet from the Earth.

			_
	rease sars.	2 2 2 4 4 7 4 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	00
	linc 6 ye		>
	Solidincreafe in 16 years.	Cubic feet.	721
	Content in 1758.	173 6 0 0 4 1 7 7 0 00 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1666
		1	
	ပိ	2 + 4 1 1 1 1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2	264
	it in	39 11 10 10 10 10 10 10 10 10 10 10 10 10	
,	Content in 1743.	n n n n n n m m n n n n n n n n)
		1991 of 1 2 2 2 2 2 8 Cubicfeet.	617
	Circumf. Circumf. Increase in in Spring in Autumn 16 years.	nl 10 sdis 4 4 0 u u w 0 0 u 0 u v	
		I - Son Ono w u Inches.	
•		0 1 0 - Feet.	
	Circumf. in Autumn 1758.	In I to sdits o o w o w w + u o - u 4	
		com σ = + σ n σ σ = Inches.	
	E.C	Feet. 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	
	ing ing 3.	In I to state 4 4 wo - 00 40 - 0 r	
,	Circumf. in Spring 1743.	.esdoni G 4 2 4 2 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2	
	Ö.s	. 1994 000 000 4 4 4 4 4 4 4 4 4 4 4 4 4 4	_
	FIRST TABLE.	Read Jan. 11, 1759. 1. Ash, planted since 1647 2. Oak, pass thriving, but sound 3. Oak, about 80 years old 4. Scotch Fir, seed in 1698 5. Oak, planted above 60 years — 6. Spanish Chefnut, near 60 years old 7. Another, 45 or 46 years old 8. Oak, planted by me in 1720 9. Scotch Fir, planted in 1734 or 1735 11. Oak, set an acorn in spring 1719 12. Oak, planted in 1720 or 1721	

Now

Now as the twelve trees above, contained 213 cubic feet 300 inches of timber in spring 1743, and have increased to 322 cubic feet 333 inches in autumn 1758; that is, 109 cubic feet 33 inches in 16 years growth; if all the trees were of the same kind, 109 feet pays 3 per cent. for standing: and the fix oaks pay near the same interest, although one of them, No 2. appeared past thriving in 1743; for the increase of the fix oaks is from 112 feet 1 quarter 171 inches of timber, to 167 feet 138 inches, i. e. 54 feet 2 quarters 399 inches; which is above 3 per But if you take only the five thriving oaks, then their content is, from 57 feet 3 quarters 267 inches, to 103 feet 2 quarters 58 inches; i. e. 45 feet 2 quarters 223 inches of timber; or near 5 per cent. And the increase of the most thriving oak, No 8. appears, by the above table, to pay above 12 ½ per cent. and the Scotch fir, No 9. being under 2 feet and half of timber in spring 1743, and 10 feet in autumn 1758, pays above 18 3 per cent. Besides it should be considered, altho' I measured the largest and most thriving oak and Scotch fir in 1743, yet feveral others of the same age, both oaks and Scotch firs, have greatly exceeded the measured trees for many years past; e.g. the oak No 11. appears by the table two feet 8 inches 2-8ths in circumference: and another just by it is 2 feet 11 inches 6-8ths; and an oak transplanted from this grove, is 3 feet 9 inches 5-8ths round; yet this last tree was considerably less than the first when removed, and not planted in a better foil, and yet is I foot I inch 3-8ths larger than the original tree. The first contains 4 feet 1 quarter 336 inches, and has gained 2 feet 3 quarters 220 inches

[9]

inches in fixteen years: the last contains 8 feet 3 quarters 68 inches; and, supposing them equal in 1743, gains 7 feet 384 inches; i. e. above 2 \frac{1}{2} the increase of the first tree. But notwithstanding the transplanted oak is thus much larger than the original oaks in the grove, yet as the transplanted tree does not run half the height of the trees in the grove before it heads, they differ but little in their quantity of timber.

The following table shews the monthly increase of trees in the years 1757 and 1758. As I endeavoured to take the measures with as much exactness as was in my power, I cut three, four, or more notches in the bark of each tree, that my line might always be confined exactly to the same place. I observed, if I measured soon after a rain, whilst the bark was saturate with water, the tree would be $\frac{1}{8}$ of an inch larger than after a day or two of dry weather. I may here add, that all the measures of circumferences of trees are taken at 5 feet from the earth: and consequently the solid measures must include 10 feet in length. I generally made use of Keay's Tables in the solid measures, which go no lower than quarters of inches in girts: which is not so exact as it ought to be.

		[10]
Kinds,	29 Sept. 1758.	H u 4 H 4 G ∞ 040 1000 has 19w visw fish month si .gs .1gs?
veral	29 Aug. 1758.	August 29, 1758. last month hot, and rather to 200 20 20 1758. last month hot, and rather with 200 20 20 20 20 20 20 20 20 20 20 20 20
ing ser	31 July 1758.	0 H 4 H 4 Q V 4 W 0. 1001, but hot nonth very rainty, but hot 4 0 0 0 4 4 0 0 0 1001.
follows 1758.	30 June 1758.	June 30, 1758. laft month rather dry.
f the	r June 1758.	4 + 4 4 4 4 6 4 6 7 4 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
bad, c	Nov. 1757.	4 4 4 4 4 4 4 4 4 6 1 1 1 1 1 1 1 1 1 1
rees I be Year	4 Sept. 1757.	September 4, 1757, from 3d Aug, to 10th , 10 to 0 t
ving T id in th	3 Aug. 1757.	August 3, 1757. the last of the last last last last last last last last
st thri	3 July 1757.	July 3, 1757. the last month was very dry.
the mo	I June 1757.	And to these trees were 1-8th of an includent in the holds are their includent in the beginning of April 1757, such 1,577, with a partition of the section includes and the section of the
rease of 5 Feet fr	Feet. Inches. 8ths of In.	Messiured October 6, 1756; and on the 6th of November they were rather less.
Measures of the monthly Increase of the most thriving Trees I had, of the following several Kinds, taken at 5 Feet from the Ground in the Years 1757 and 1758.	TABLE II.	1. An Oak planted in 1720, No. 8. 7 in the other table ————————————————————————————————————

TABLE III. Shews the Increase in Circumference, and in solid Measure, of each Tree in 1758.

					-	,						
Per Cent.	7	36	4	$12\frac{1}{2}$	124	4	21	93	284	19	19	36
	or near	above	about	above	apove	near	near	apove	apove	apove	above	14 about
Inches.	72	390	340	112	981	340	198	426	48	162	258	14
Quarters.	0	0	-	-	13	-	=	(4	-	0	-	19
Cubic feet.	-	0	0	-	0	0	0	0	0	-	0	0
Inches.	260	174		408	216	408	174	264	∞	48	0	284 0
Quarters.	0	'n	14	14	7	7	0	-	-	60	7	'n
				Ξ			14	∞	-	17	7	_
Inches.	188	216	89	296	30	89	408	270	392	318	174	270
Quarters.	0	8		H	0	H	7	13	3	N	0	-
Cubic feet.	15	0	Ξ	01	2	Ξ	-	7	0	91	8	1-4
Sths.	14	"	9	0	9	4	"	0	Ħ	3	5	0
Inches.	m	14	4	4	11	4	0	00	٠,	4	0	6
Feet.	٧.	_	+	4	7	4	_	3	-	20	7	~
.al to sd18	8	4	9	4	64	4	4	7	-	"	4	71
Inches.	Ξ	0	10		0	60	00	9	"	64	01	9
Feet.	4	_	4	4	7	4	-	3	-	2	-	_
	1	1	1	٠ <u>-</u>	. [1	I	l	1	1	1
	1. The Oak	2. The Beech	3. The Scotch Fir		4. The Oak	5. The Spanish Chefnut	6. The Elm	7. The Spruce Fir -	8. The Larch	9. The Willow -	10. The Beech	A Weymouth Pine
	Feet. Inches. Sths of In. Feet. Inches. Sths. Cubic feet. Inches. Ouarters. Inches. Cubic feet.	Feet. Inches. Reta of In. Reta. Inches. Inches. Reta. Reta. Marters. Cubicfeet. Cubicfeet. Cubicfeet. Cubicfeet. Marters. Cubicfeet. Marters. Ouarters. Ouarters.	Feet. Feet.	4 1 Feet. 3 0 1 Inches. 4 1 2 Feet. 4 2 Inches. 5 6 4 2 Sths of In. 6 4 1 5 Feet. 7 1 Cubic feet. 6 2 1 Cubic feet. 7 2 Inches. 1 2 0 Quarters. 1 2 0 Quarters. 1 2 188 Inches. 1 2 18 Inches. 1 2 18 17 6 Inches. 1 3 17 6 Inches. 1 2 18 17 6 Inches. 1 3 17 6 Inches. 1 2 18 17 6 Inches. 1 3 17 6 Inches. 1 2 18 18 Inches. 1 3 17 6 Inches. 1 2 18 18 Inches. 1 3 17 6 Inches. 1 3 17 6 Inches. 1 3 17 6 Inches. 1 2 18 18 18 Inches. 1 3 18 18 18 18 Inches. 1 3 18 18 18 18 Inches. 1 5 18 18 18 18 Inches. 1 6 18 18 18 18 18 Inches. 1 7 18 18 18 18 18 Inches. 1 8 18 18 18 18 18 18 18 18 18 18 18 18	2d	2d	2d	2d	2d	2d	2d	2d

N. B. I measure the oak No 4. as three feet round, as it wants only 1-4th of an inch of that measure; and the Weymouth pine as I foot and 6 inches, tho' it is I-4th of an inch more.

As the Scotch fir, N° 3. has been fickly for two years paft; therefore I add another Scotch fir (one year younger) to flow the growth of that kind of tree; and the extraordinary increase of the Weymouth pine induced me to put that in also, tho' I had not measured it monthly.

The great Lord Bacon fays, "the improvement of the ground " is the most natural way of obtaining riches." What great fortunes might be raifed, by those that have property, in the vast heaths and downs, or fields of poor land, in this kingdom, by planting parts of them? which would also add great beauty to the country, and render the dwelling much more comfortable to the neighbourhood, by the shade in summer, and warmth in winter. Some parts of these great wastes would produce good oak; and where the foil is moist, poplar, alder, and other aquatics, would be very profitable to the planter. chalky foil feems the least promising; yet beeches sometimes thrive well upon it. The fir kind, especially the Scotch fir, will grow furprifingly upon poor fandy land; but woods of fir should be guarded with an out-line of birch and beech, to break the force of strong winds. Birch, being the quickest grower, will best protect the young fir; but as birch, after a few years, is easily blown down, so beech will be wanted to defend the firs as they become large: for I have feen broad glades made by the wind through great woods of fir in Switzerland: which, perhaps, might have been prevented, at least in part, by an out-line of beech.

I know some think, that poor land cannot produce large trees; yet the oak at Northall in Hertsordshire, whose beautiful head spreads a circle of above 40 yards diameter, stands on a dry and deep sand; and the fine chesnuts and beeches by Mr. Naylor's grand castle of Herst Monceux in Sussex, grow in a light sandy soil: and I have sound, by experience, the Weymouth, Scotch, spruce, and silver firs, which I planted in a poor sandy soil, are larger and siner trees, than others set at the same time in much better land. Perhaps it may require a rich clay to produce such trees as the noble grove of oak in the Earl of Powis's park by Ludlow, or Lord Ducie's vast chesnut at Tortworth, in Gloucestershire, which I measured 46½ feet in

circumference at near 6 feet from the ground.

Although these slight observations are not so deserving the attention of the Royal Society as I could wish; yet they may possibly be the means of producing better; and for my own part, I shall always esteem it a great honour that they were communicated by Dr. Hales.

R. Marsham.